


UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



CHEMISTRY 9701/04

Paper 4 Structured Questions

October/November 2008

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid

DO NOT WRITE IN ANY BARCODES.

Section A

Answer all questions.

Section B

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
Total		

This document consists of 18 printed pages and 2 blank pages.



[4]

Section A

Answer all questions in the spaces provided.

1 (a) Natural bromine consists of the two isotopes ⁷⁹Br and ⁸¹Br in roughly equal proportions.

The mass spectrum of bromine consists of 5 peaks.

(i)	Suggest the mass numbers for the 5 peaks and the identities of the species responsible for them.
(ii)	Suggest the ratios of the relative abundances of
	the three lines with the highest mass numbers,
	the two lines with the lowest mass numbers.

Esters of 2,3-dibromopropan-1-ol with phosphoric acid are useful flame retardants used in plastics and fibres.

2,3-dibromopropan-1-ol can be made from propenal by the following two-stage process.

[3]

(b) (i)	Draw the structure of the intermediate A in the box opposite.
(ii)	Suggest reagents and conditions for
	• reaction I,
	reaction II.

(c) The mass spectrum of 2,3-dibromopropan-1-ol includes the following peaks.

mass number	relative abundance
31	100
106	44
108	45
185	0.3
187	0.6
189	0.3

(i) At what mass number would you expect the molecular ion to occur?

(ii) Identify the molecular formula (including isotopic composition where relevant) of these 6 peaks.

mass number	molecular formula
31	
106	
108	
185	
187	
189	

[5]

[Turn over

2 In the late 19th century the two pioneers of the study of reaction kinetics, Vernon Harcourt and William Esson, studied the rate of the reaction between hydrogen peroxide and iodide ions in acidic solution.

$$H_2O_2 + 2I^- + 2H^+ \rightarrow 2H_2O + I_2$$

This reaction is considered to go by the following steps.

step 1
$$H_2O_2 + I^- \rightarrow IO^- + H_2O$$

step 2
$$IO^- + H^+ \rightarrow HOI$$

step 3
$$HOI + H^+ + I^- \rightarrow I_2 + H_2O$$

The general form of the rate equation is as follows.

rate =
$$k[H_2O_2]^a[I^-]^b[H^+]^c$$

(a) Suggest how the appearance of the solution might change as the reaction takes place.


.....[1

(b) Suggest values for the orders *a*, *b* and *c* in the rate equation for each of the following cases.

0000	numerical value				
case	а	b	С		
step 1 is the slowest overall					
step 2 is the slowest overall					
step 3 is the slowest overall					

[3]

A study was carried out in which both $[H_2O_2]$ and $[H^+]$ were kept constant at 0.05 mol dm⁻³, and $[I^-]$ was plotted against time. The following curve was obtained.



To gain full marks for the following answers you will need to draw relevant construction lines on the graph opposite to show your working. Draw them using a pencil and ruler.

	0	•	o .	0 ,	
(c)	Calculate th	ne initial rate of this i	eaction and state	its units.	
			rate =	uni	its[2]
(d)	Use half-life	e data calculated fro			tion is first order with
()	respect to []		u.o g.ap to o.		
					[2]
(e)	Use the following your reason		ce the orders with	respect to [H ₂ O ₂]	and [H ⁺], explaining
	•	_			1
		$[\mathrm{H_2O_2}]/\mathrm{moldm^{-3}}$	[H ⁺]/moldm ⁻³	relative rate	
		0.05	0.05	1.0	
		0.07	0.05	1.4	
		0.09	0.07	1.8	
		order wit	h respect to [H ₂ O ₂	1 =	
		order v	viiii respect to [i i]	[2]
(f)	From your step.	results, deduce whi	ch of the three st	teps is the slowe	est (rate determining)
					[1]
					[Total: 11]

3	(a)	(i)	Describe and explain the trend observed in the thermal stability of the carbonates of the Group II elements.
		(ii)	By quoting suitable data from the <i>Data Booklet</i> suggest how the thermal stabilities of
			zinc carbonate andlead carbonate
			might compare to that of calcium carbonate.
			[6]
	(b)	Mal	achite is an ore of copper. It contains the following percentages by mass.
			copper 57.7% oxygen 36.2%
			carbon 5.4% hydrogen 0.9%
			achite reacts with dilute $\rm H_2SO_4$ producing a gas B that turns limewater milky and ving a blue solution C .
			en heated in the absence of air, malachite produces gas B and steam, and leaves a ck solid D . D reacts with dilute H ₂ SO ₄ to produce the same blue solution C .
		Add	ling iron filings to C produces a pink solid E and a pale green solution F .
		(i)	Calculate the empirical formula of malachite.


(ii)	Suggest the formula of the ion responsible for the blue colour of solution C.
(iii)	Identify the black solid D and calculate the mass of D that could be obtained by heating 10 g of malachite.
(iv)	Use data from the <i>Data Booklet</i> to identify the pink solid E and the solution F , and suggest an equation for the reaction producing them.
(v)	What type of reaction is the reaction that produces E and F ?
(vi)	Describe and explain what you would see happen when dilute NH ₃ (aq) is added slowly to the solution C until it is in an excess.
	[13]

[Total: 19]

4	(a)		viscosity of engine oil can be improved by the addition of certain medium chainth polymers.
		Арс	ortion of the chain of one such polymer is shown below.
			$-CH_2CH(CH_2CH_2CH_3)CH_2CH(CH_2CH_2CH_3)CH_2-$
		On	average, the molecules of the medium-chain polymer contain 40 carbon atoms.
		(i)	Suggest the structure of the monomer.
		(ii)	How many monomer units are incorporated into the average molecule of the polymer?
			[2]
	(b)		d car engine oil can be recycled for use as a fuel by the processes of distillation and king.
		(i)	Assuming a typical molecule of engine oil has the formula $C_{40}H_{82}$, suggest an equation for a cracking reaction that could produce diesel fuel with the formula $C_{16}H_{34}$ and other hydrocarbons only.
		(ii)	What conditions are needed for this cracking reaction?
		(iii)	Considering only the bonds broken and the bonds formed during the reaction, use the <i>Data Booklet</i> to calculate the enthalpy change for the reaction you wrote in (b)(i) .
		(iv)	Comment on how the conditions you described in (b)(ii) relate to the enthalpy change you calculated in (b)(iii) .
			[4]
			[Total: 6]

5 (4-aminophenyl)ethanoic acid (4-APEA) and its derivatives are being investigated as possible drugs to treat chronic inflammation of the intestines.

The synthesis of 4-APEA from methylbenzene is shown in the following scheme.



- (a) Draw the structures of the compounds **G** and **H** in the boxes above. [2]
- (b) Suggest reagents and conditions for the following steps.
 - step II

step V

step III

[3]

[Total: 5]

[2]

		1	0		
6		a test or simple reaction you could carry out on each of the following pairs of nds to enable them to be distinguished.			
	(a)	CH ₃ COCH ₂ CH ₂ CH ₃	CH ₃ CH ₂ COCH ₂ CH ₃		
		J	K		
	(i)	description of test or reaction			
	(ii)	observation with compound J			
	(iii)	observation with compound ${\bf K}$			
			[2]		
	(b)		[-]		
	(2)	\sim NH ₂	\sim NH ₂		
		L	М		
	(i)	description of test or reaction			
	<i>/</i> ····	ahaamattan sitti a			
	(ii)	observation with compound L			

(iii) observation with compound ${\bf M}$

(c)	CH ₃ CH ₂ COC <i>l</i>	$\mathrm{CH_3CH_2CH_2C}\mathit{l}$
	N	Р
(i)	description of test or reaction	
(ii)	observation with compound N	
(/		
(iii)	observation with compound P	
(111)	observation with compound F	
		[2]
(d)	CH ₃ CH ₂ CONH ₂	CH ₃ CH ₂ CH ₂ NH ₂
	Q	R
(i)	description of test or reaction	
.,	·	
(ii)	observation with compound Q	
(11)	observation with compound &	
<i>,</i> ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
(iii)	observation with compound R	
		[2]
		[Total: 8]
		11

7	(a)	Explain briefly what is meant by the word <i>protein</i> .
	(b)	Describe how peptide bonds are formed between amino acids during the formation of a tripeptide. Include diagrams and displayed formulae in your answer.
		[3
	(c)	Describe how proteins can be broken down into amino acids in the laboratory withou the aid of enzymes.
	(d)	
	S	3 NH ₂ CH ₂ CO ₂ H + 2 NH ₂ CHCO ₂ H + 2 NH ₂ CHCO ₂ H CH ₃ CH ₂
		$M_{\rm r}=75 \qquad \qquad M_{\rm r}=89$
		$M_{\rm r} = 165$ (i) How many peptide bonds were broken during this reaction?
		(ii) Calculate the $M_{\rm r}$ of the polypeptide S .
		M —
		$M_{\rm r} = \dots$ [3

[Total: 9]


BLANK PAGE

9701/04/O/N/08 **[Turn over**

Section B – Applications of Chemistry

Answer all questions in the spaces provided.


- **8 (a)** Enzymes play a vital role in all living organisms, helping chemical reactions to take place at body temperature.
 - (i) The diagram below shows the reaction pathway of an enzyme-catalysed reaction without an enzyme present. On the diagram sketch the pathway if the enzyme was present.



(ii)	What type of molecule are most enzymes?
(iii)	Why do many enzymes lose their catalytic effectiveness above 40°C?
	[3]
(b) (i)	Explain the difference between competitive and non-competitive inhibition of an enzyme.

(ii) The graph below shows how the rate of an enzyme-catalysed reaction varies with substrate concentration in the absence of an inhibitor.

For a given amount of enzyme, $V_{\rm max}$ represents the rate when all of the active sites on the enzyme are being used.



Sketch on the diagram curves to show the effect on the rate of reaction of:

I a competitive inhibitor;

II a non-competitive inhibitor.

(i) Suggest to what atom or group Hg²⁺ ions bind.

Clearly label your curves.

[4]


(c) Heavy metal ions like ${\rm Hg^{2+}}$ can bind irreversibly to enzymes and this can result in poisoning.

[Total: 10]

9


	hnology of DNA fingerprinting has enormously advanced scientific identification es in medicine, crime detection and archaeology in recent years.
(a) (i)	In order to prepare a DNA sample for analysis, the DNA is treated with restriction enzymes. What do restriction enzymes do?
(ii)	What is the next stage in DNA analysis, after the treatment with restriction enzymes?
(iii)	How are the DNA fragments made visible?
	[3]
(b) NM	R and X-ray crystallography have made significant contributions to our knowledge of
the	structure of proteins and, in the pharmaceutical industry, how drugs react with target reins.
(i)	Suggest an advantage of each technique in helping to determine protein structure.
(ii)	MRI scanning is a medical technique based on NMR spectroscopy. It is particularly useful for looking for tumours in healthy tissue.
	Suggest how this technique can distinguish tumour tissue from healthy tissue.
	[3]

- (c) A saturated molecule of formula C_xH_yNO was subjected to analysis by mass spectrometry and NMR spectroscopy. In the mass spectrum of the compound, the M peak was at m/e 73 and the ratio of the heights of the M:M+1 peak was 48:1.7.
 - (i) Using the data from the mass spectrum, determine the values of x and y in the formula of the compound.
 - (ii) Use the data from (i) together with the NMR spectrum below to deduce a structure for the compound, explaining how you arrive at your answer.



[Turn over

10 (a) Silk from silkworms, used as a fabric shows a different secondary structure to that produced by spiders.



(i)	What sort of bonding would you expect to occur between adjacent parts of the protein chains in each form of silk?
	silkworm
	spider
(ii)	Suggest two differences in properties that these forms of silk could have. Explain your answer.
(iii)	Spider dragline silk contains large amounts of the amino acid glycine. How does this affect the properties of the silk?
	[5]

(b)	Botl	n forms of silk are condensation polymers.
	(i)	Explain what is meant by a condensation polymer.
	<i>(</i>)	
	(ii)	Another type of polymer is called an addition polymer. Name an example of an addition polymer.
	(iii)	Suggest why condensation polymers such as proteins show a wider range of properties than addition polymers.
		[5]
		[Total: 10]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.