



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
General Certificate of Education  
Advanced Subsidiary Level and Advanced Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

Advanced Practical Skills 2

**9701/34**

**October/November 2011**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
Give details of the practical session and laboratory where appropriate, in the boxes provided.  
Write in dark blue or black pen.  
You may use a soft pencil for any diagrams, graphs or rough working.  
Do not use staples, paper clips, highlighters, glue or correction fluid.  
**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.

<b>Session</b>
<b>Laboratory</b>

<b>For Examiner's Use</b>	
<b>1</b>	
<b>2</b>	
<b>Total</b>	

This document consists of **11** printed pages and **1** blank page.



- 1 You are to determine the enthalpy change of neutralisation of sodium hydroxide by an acid and also the concentration of hydrogen ions in the acid. These can be found by measuring the temperature change when solutions of the acid and alkali are mixed.

For  
Examiner's  
Use

**FB 1** is  $1.50 \text{ mol dm}^{-3}$  sodium hydroxide, NaOH.

**FB 2** is an aqueous solution of an acid.

**(a) Method**

- Fill a burette with **FB 1**. [**Care: FA1 is corrosive**]
- Support the plastic cup in a  $250 \text{ cm}^3$  beaker.
- Run  $10.0 \text{ cm}^3$  of **FB 1** from the burette into the plastic cup.
- Measure and record, in the table below, the temperature of the **FB 1** in the cup. You may need to tilt the beaker to ensure that the bulb of the thermometer is covered.
- Measure  $40 \text{ cm}^3$  of **FB 2** using the measuring cylinder.
- Pour this volume of **FB 2** into the plastic cup containing **FB 1**. Stir carefully and measure the highest temperature obtained.
- Record this temperature in the table.
- Rinse the plastic cup with water.
- Repeat the experiment using  $15.0 \text{ cm}^3$  of **FB 1** and  $35 \text{ cm}^3$  of **FB 2** as shown for experiment **2** in the table.
- Carry out experiments **3** to **7** in the same way.
- Complete the table for each experiment.

**Results**

<i>experiment number</i>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>		
volume of <b>FB 1</b> / $\text{cm}^3$	10.0	15.0	20.0	25.0	30.0	35.0	40.0	50.0	0.0
volume of <b>FB 2</b> / $\text{cm}^3$	40	35	30	25	20	15	10	0	50
initial temperature <b>FB 1</b> /°C									
highest temperature/°C									
temperature change/°C								0.0	0.0


[8]

I	
II	
III	
IV	
V	
VI	
VII	
VIII	

- (b) On the grid below plot the temperature **change** (y-axis) against the volume of **FB 1** (x-axis). Use **all** the results in the table including those provided in the final two columns.

For  
Examiner's  
Use

Using these points, draw two straight lines that intersect.



[4]

I	
II	
III	
IV	

- largest temperature change = ..... °C

- volume of **FB 1** required is = ..... cm<sup>3</sup>

- moles of sodium hydroxide present = ..... mol

- [Assume that 4.3 J are required to raise the temperature of 1 cm<sup>3</sup> of any solution by 1°C]

heat energy produced = ..... J

- Give your answer in  $\text{kJ mol}^{-1}$  and include the relevant sign.

enthalpy change of neutralisation = ..... kJ mol<sup>-1</sup>  
sign value

[7]

I	
II	
III	
IV	
V	
VI	
VII	

- (d) Identify a source of error, other than heat loss, in the experimental method. Suggest an improvement which would reduce this source of error.

For  
Examiner's  
Use

.....

.....

.....

..... [2]

- (e) (i) Calculate the concentration of hydrogen ions,  $\text{H}^+$ , in  $\text{mol dm}^{-3}$ , present in **FB 2**.

concentration of hydrogen ions in **FB 2** = .....  $\text{mol dm}^{-3}$

- (ii) If the acid present in **FB 2** is sulfuric acid, calculate its concentration.

concentration of sulfuric acid = .....  $\text{mol dm}^{-3}$

- (iii) Describe a chemical test by which you could prove that the acid in **FB 2** is sulfuric acid.

**Do not carry out the test.**

.....

.....

.....

.....

[4]

[Total: 25]

I	
II	
III	
IV	

## 2 Qualitative Analysis

For  
Examiner's  
Use

At each stage of any test you are to record the details of the following

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

**No additional tests for ions present should be attempted.**

**If any solution is warmed, a boiling tube MUST be used.**

Rinse and reuse test-tubes and boiling tubes where possible.

**Where reagents are selected for use in a test, the full name or correct formula of the reagents must be given.**

- (a) (i) **FB 3, FB 4 and FB 5** are aqueous solutions of sodium compounds. None of these compounds contains sulfur.

To about 1 cm depth of each of the solutions in separate test-tubes add the same depth of dilute sulfuric acid.

To another 1 cm depth of each of the solutions in separate test-tubes add a few drops of aqueous lead(II) nitrate.

Record your observations for these tests in an appropriate form in the space below.

I	
II	
III	
IV	
V	
VI	

Using the Qualitative Analysis Notes printed on pages 10 and 11 and your observations identify the anions in **FB 3** and **FB 4**.

For  
Examiner's  
Use

**FB 3** contains .....

**FB 4** contains .....

[6]

- (ii) Select a reagent or pair of reagents that would enable you to determine the identity of the anion in **FB 5**.

reagent(s) .....

Carry out a test on **FB 5** using the reagent(s) given above. Record your observations below.

.....

.....

.....

I	
II	
III	

The anion present in **FB5** is .....

[3]

(b) You are provided with solid **FB 6**. Complete the following table.

For  
Examiner's  
Use

<i>test</i>		<i>observations</i>
(i)	To 1 cm depth of aqueous hydrogen peroxide in a test-tube, add a very small spatula measure of <b>FB 6</b> .	
(ii)	To 1 cm depth of aqueous iron(II) sulfate in a boiling tube, add the same depth of dilute sulfuric acid. Add a very small spatula measure of <b>FB 6</b> to the tube. Warm the mixture gently using a Bunsen burner for about 20 seconds, then filter the warm mixture and collect the filtrate.	
	To the filtrate, add aqueous sodium hydroxide.	
(iii)	To 2 cm depth of aqueous potassium manganate(VII) in a test-tube, add the same depth of aqueous sodium hydroxide and then a small spatula measure of <b>FB 6</b> . Stir the contents of the test-tube for about 20 seconds. Filter the mixture and collect the filtrate.	
	To the filtrate, add dilute sulfuric acid.	



Suggest a conclusion that could be made about the chemical behaviour of **FB 6** from the observations in **(i)**. Explain the reasons for your answer.

.....  
 .....

What conclusion can be made about the chemical behaviour of **FB 6** from the observations in **(ii)**?

.....  
 .....

[6]

[Total: 15]

For  
Examiner's  
Use

I	
II	
III	
IV	
V	
VI	

## Qualitative Analysis Notes

Key: [ ppt. = precipitate ]

## 1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al <sup>3+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> <sup>+</sup> (aq)	no ppt. ammonia produced on heating	
barium, Ba <sup>2+</sup> (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr <sup>3+</sup> (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu <sup>2+</sup> (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe <sup>3+</sup> (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
lead(II), Pb <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
magnesium, Mg <sup>2+</sup> (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. soluble in excess

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

## 2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, $\text{CO}_3^{2-}$	$\text{CO}_2$ liberated by dilute acids
chromate(VI), $\text{CrO}_4^{2-}(\text{aq})$	yellow solution turns orange with $\text{H}^+(\text{aq})$ ; gives yellow ppt. with $\text{Ba}^{2+}(\text{aq})$ ; gives bright yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$ ); gives yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
nitrate, $\text{NO}_3^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil
nitrite, $\text{NO}_2^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil; $\text{NO}$ liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown $\text{NO}_2$ in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ or with $\text{Pb}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	$\text{SO}_2$ liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

## 3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ )
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	"pops" with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint
sulfur dioxide, $\text{SO}_2$	turns acidified aqueous potassium dichromate(VI) from orange to green

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.