



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

MATHEMATICS 9709/06

Paper 6 Probability & Statistics 1 (S1)

October/November 2008

1 hour 15 minutes

Additional Materials: Answer Booklet/Paper

Graph Paper

List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.



1	below.								
	32 35 45 37 38 44 33 39 36 45								
	Find the mean and standard deviation of the lengths of these leaves. [3]								
2	On a production line making toys, the probability of any toy being faulty is 0.08. A random sample of 200 toys is checked. Use a suitable approximation to find the probability that there are at least 15 faulty toys.								
3 (i) The daily minimum temperature in degrees Celsius (°C) in January in Ottawa is a random with distribution N(-15.1, 62.0). Find the probability that a randomly chosen day in Ja Ottawa has a minimum temperature above 0 °C.									
	(ii) In another city the daily minimum temperature in $^{\circ}$ C in January is a random variable with distribution N(μ , 40.0). In this city the probability that a randomly chosen day in January has a minimum temperature above 0 $^{\circ}$ C is 0.8888. Find the value of μ .								
4 A builder is planning to build 12 houses along one side of a road. He will build 2 houses in 2 houses in style <i>B</i> , 3 houses in style <i>C</i> , 4 houses in style <i>D</i> and 1 house in style <i>E</i> .									
	(i) Find the number of possible arrangements of these 12 houses. [2]								
	(ii)								
	Road								
	First group Second group								
The 12 houses will be in two groups of 6 (see diagram). Find the number of possible arrangement if all the houses in styles A and D are in the first group and all the houses in styles B , C and are in the second group.									
	(iii) Four of the 12 houses will be selected for a survey. Exactly one house must be in style <i>B</i> are exactly one house in style <i>C</i> . Find the number of ways in which these four houses can be selected.								
5	The pulse rates, in beats per minute, of a random sample of 15 small animals are shown in the following table.								
	115 120 158 132 125								
	104 142 160 145 104								
	162 117 109 124 134								
	(i) Draw a stem-and-leaf diagram to represent the data. [3]								
	(ii) Find the median and the quartiles. [2]								

© UCLES 2008 9709/06/O/N/08

6	There are three sets of traffic lights on Karinne's journey to work. The independent probabilities that
	Karinne has to stop at the first, second and third set of lights are 0.4, 0.8 and 0.3 respectively.

- (i) Draw a tree diagram to show this information. [2]
- (ii) Find the probability that Karinne has to stop at each of the first two sets of lights but does not have to stop at the third set. [2]
- (iii) Find the probability that Karinne has to stop at exactly two of the three sets of lights. [3]
- (iv) Find the probability that Karinne has to stop at the first set of lights, given that she has to stop at exactly two sets of lights.
- 7 A fair die has one face numbered 1, one face numbered 3, two faces numbered 5 and two faces numbered 6.
 - (i) Find the probability of obtaining at least 7 odd numbers in 8 throws of the die. [4]

The die is thrown twice. Let X be the sum of the two scores. The following table shows the possible values of X.

		Second throw								
		1	3	5	5	6	6			
	1	2	4	6	6	7	7			
	3	4	6	8	8	9	9			
First	5	6	8	10	10	11	11			
throw	5	6	8	10	10	11	11			
	6	7	9	11	11	12	12			
	6	7	9	11	11	12	12			

(ii) Draw up a table showing the probability distribution of X. [3]

(iii) Calculate E(X). [2]

(iv) Find the probability that X is greater than E(X). [2]

© UCLES 2008 9709/06/O/N/08

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.