## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

## MARK SCHEME for the May/June 2010 question paper for the guidance of teachers

## 9701 CHEMISTRY

9701/23

Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.




Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	23


1 (a) enthalpy change when 1 mol of a compound is formed (1) from its elements (1) in their standard states under standard conditions (1) [3]

(b) (i) 
$$N_2H_4(I) + O_2(g) \rightarrow N_2(g) + 2H_2O(g)$$
  
 $\Delta H_f^{\circ}/kJ \text{ mol}^{-1} +50.6 -241.8$   
 $\Delta H^{\circ}_{\text{reaction}} = 2(-241.8) - (+50.6) (1)$   
 $= -534.2 \text{ kJ mol}^{-1} (1)$ 

- (ii)  $E_a$  is too high (1)
- (iii) products are H<sub>2</sub>O and N<sub>2</sub> which are harmless/non toxic or are already present in the atmosphere (1)
- (c) (i) 'dot-and-cross' diagram (1)



(iii) minimum is



allow bond angle around N atom between 109° and 104° (1) [4]

[Total: 12]

[4]

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	23

2 (a) the energy required to remove one electron from each atom (1) in one mole of gaseous atoms (1)

or

the enthalpy change in kJ mol<sup>-1</sup> for (1)

$$M(g) \rightarrow M^{\dagger}(g) + e^{-}(1)$$
 [2]

- (b) (i) first ionisation energy decreases down Group 1 (1) outermost electron is further from nucleus or has greater shielding (1)
  - (ii) outermost electron experiences less attraction
     or formation of M<sup>+</sup> cation becomes easier down Group 1 (1)
     [3]
- (c) (i)  $n(Li) = \frac{0.83}{6.9} = 0.12(1)$ 
  - (ii) 2 mol Li  $\to$  1 mol H<sub>2</sub> 0.12 mol Li  $\to$   $\frac{1 \times 0.12}{2}$  = 0.06 mol H<sub>2</sub> (1) volume of H<sub>2</sub> = 0.06 × 24.0 = 1.44dm<sup>3</sup> (1)
  - (iii) 2 mol Li  $\to$  2 mol LiOH 0.12 mol Li  $\to$  0.12 mol LiOH in 0.50 dm<sup>3</sup> (1) [LiOH] =  $\frac{0.12 \times 1}{0.50}$  = 0.24 mol dm<sup>-3</sup> (1) [5]
- (d) sodium burns with a yellow flameor white solid formedor colour of chlorine disappears (1)

$$2Na + Cl_2 \rightarrow 2NaCl(1)$$
 [2]

[Total: 12]

3	(a) (i)	Ca (1)		
	(ii)	S or C [allow H (H <sub>2</sub> O <sub>2</sub> ) or I	N (NO, NO <sub>2</sub> )] (1)	
	(iii)	He (1)		
	(iv)	Al (1)		
	(v)	Si <b>or</b> Ge (1)		
	(vi)	Al (1)		[6]
	<b>(b)</b> any	two from N or O or F (1)		[1]
	(c) (i)	$Al_2O_3$ or $SiO_2$ (1)		
	(ii)	$SO_2$ or $P_2O_3/P_4O_6$ (1) and and $SO_3$ or $P_2O_5/P_4O_{10}$ (1		
	(iii)	Na₂O (1)		
	(iv)	Al <sub>2</sub> O <sub>3</sub> (1)		[5]
				[Total: 12]
4	(a) rea	action 1 free radical subs	stitution (1)	
	rea	ection 2 elimination (1)		[2]
	(b) (i)	in reaction 4	CH <sub>3</sub> C(OH)(CN)CH <sub>3</sub> (1)	
	(ii)	in reaction 3	I <sup>-</sup> (1)	
	(iii)	in reaction 3 <b>or</b> in reaction 4	CH <sub>3</sub> I CH <sub>3</sub> COCH <sub>3</sub> (1)	[3]
		pecies which has a lone pai which reacts with an electro	ir of electrons on deficient $(\delta+)$ centre in a molecule $(1)$	[1]
		eaction 3 OH <sup>-</sup> (1) eaction 4 CN <sup>-</sup> (1)		[2]
	<b>(e)</b> π b	onding is electron rich (1)		[1] [Total: 9]

Mark Scheme: Teachers' version GCE AS/A LEVEL – May/June 2010

Page 4

Paper 23

Syllabus 9701

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	23

5 (a)

[3]

(b)

	reagent(s)	condition(s)
step 1	Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup> / H <sup>+</sup>	distil off aldehyde
	(1)	(1)
step 2	HCN in presence of CN <sup>-</sup> or KCN + dil H <sub>2</sub> SO <sub>4</sub> (1)	room temperature (1)
step 3	aqueous mineral acid/ /H <sub>2</sub> SO <sub>4</sub> /HC/  not HNO <sub>3</sub> (1)	heat under reflux (1)

in **each** case, the reagent must be correct before the condition mark is awarded [6]

- (c) (i) a protein (1)
  - (ii) 2,4-dinitrophenylhydrazine/Brady's reagent (1) yellow-orange-red ppt. (1)
  - (iii) acidified  $K_2Cr_2O_7$  or Lucas test or  $CH_3CO_2H/H^+$  (1) colour changes or cloudiness or fruity smell from orange to green (1)
  - (iv) LiA/H<sub>4</sub>/NaBH<sub>4</sub> or H<sub>2</sub>/Ni etc. (1) [6]

[Total: 15]