

Cambridge International AS & A Level

CHEMISTRY		9701/05
Paper 5 Planning, Analysis and Evaluation		For examination from 2022
MARK SCHEME		
Maximum Mark: 30		
	Specimen	

This document has 8 pages. Blank pages are indicated.

© UCLES 2019 [Turn over

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
 - the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded positively:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
 - marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind

Science-Specific Marking Principles

3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus
	terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).

correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where

5 <u>List rule' guidance</u> (see examples below)

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided
- Any response marked *ignore* in the mark scheme should not count towards *n*
- Incorrect responses should not be awarded credit but will still count towards *n*
- awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be be treated as a single incorrect response
 - Non-contradictory responses after the first **n** responses may be ignored even if they include incorrect science.

Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, unless the question states 'show your working'

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values

For answers given in standard form, (e.g. $a \times 10^{n}$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme

Mark scheme abbreviations:

separates marking points

separates alternatives within a marking point

gnore mark as if this material was not present

accept (a less than ideal answer which should be marked) indicates mark is conditional on previous marking point COND

or words to that effect (accept other ways of expressing the same idea) OWTTE

alternative wording (where responses vary more than usual)

⋛

actual word given must be used by candidate (grammatical variants accepted) underline

credit a correct statement that follows a previous wrong answer indicates the maximum number of marks that can be awarded max ECF

he word / phrase in brackets is not required, but sets the context

ဖ

© UCLES 2019

Page 4 of 8

~

(discount 3)

3. Correct CON (of 3.)

G (5 responses)

1. Correct 2. Correct

Cambridge International AS & A Level – Mark Scheme **SPECIMEN**

က

ignore ignore

3. Correct Correct CON (of 4.)

H (4 responses)

				7	
oly the list rule	1		<i>^</i>	>	×
 Examples of now to apply the list rule 	State three reasons [3]	4	1. Correct	2. Correct	3. Wrong

F (4 responses)

1. Correct 2. Correct

D (4 responses)		
1. Correct, Correct	\ \ '	
2. Correct	>	က
3. Wrong	ignore	

			7	
2		>	× ,×	ignore
	C (4 responses)	1. Correct	2. Correct, Wrong	3. Correct
Page 5 of 8				

	7		
>	× ,×	ignore	
1. Correct	2. Correct, Wrong	3. Correct	

1. Correct	>	
2. Correct, CON (of 2.)	*, (discount 2)	7
3. Correct	>	

7				က	
*, (discount 2)	>		>	>	>
2. Correct, CON (of 2.) x, (discount 2)	3. Correct	E (4 responses)	1. Correct	2. Correct	3. Correct. Wrong

1. Correct	>	
2. Correct	×	2
3. CON (of 2.) Correct	(discount 2)	ĺ
(4 responses)		

7			
>	×	ر (discount 2)	
1. Correct	2. Correct	3. Correct CON (of 2.)	

Question	Answer	Marks
1(a)(i)	CuCO ₃ and Cu(OH) ₂ both react (with HC <i>l</i>)	_
1(a)(ii)	(Transfer) 12.5(0) cm ³ of (10.0 mol dm ⁻³) HCl ; using a burette / graduated pipette ; add to a 250 cm ³ volumetric flask AND make to mark with distilled water ;	ო
1(a)(iii)	Suitable apparatus for production of ${\rm CO}_2$ without initial gas loss e.g. divided conical flask or solid suspended in small test tube (using cotton) above acid; Suitable means of measuring ${\rm CO}_2$ produced e.g. gas syringe or inverted measuring cylinder above water;	2
1(a)(iv)	Correct labels on axes y-axis: volume (of gas) AND x-axis: time or t;	2
	curved line (from origin) to reach a plateau ; e.g.	
1(a)(v)	mol of $CuCO_3 = 0.494 \div 123.5 = 4.00 \times 10^{-3}$ mol;	7
	volume of HC l = 2 × 4.00 × 10 ⁻³ ÷ 0.500 = 0.0160 dm ³ = 16.0 cm ³ ;	
1(b)(i)	25.55, 23.90, 24.40, 24.15	7
1(b)(ii)	$[(0.05 \times 2) / 23.90] \times 100 = 0.42\%$	-

Question	Answer	Marks
1(b)(iii)	moles of $H_2SO_4 = 0.40 \times \frac{24.15}{1000} = 9.66 \times 10^{-3} \text{ mol}$; mass of $Cu_3(CO_3)_2(OH)_2 = 344.5 \times 9.66 \times 10^{-3} \div 3 = 1.11 \text{ g}$;	ო
	% by mass = $\frac{1.11}{1.50} \times 100\% = 74.0\%$;	
1(b)(iv)	problem 1 titres are not concordant / are too far apart / are $0.5(0)$ cm ³ apart / difference is too large;	4
	improvement 1 repeat until (two) concordant titres have been achieved / two readings within $0.1(0)\mathrm{cm}^3$;	
	problem 2 colour change (of indicator) will be difficult to see ;	
	improvement 2 use an alternative indicator / named indicator ;	

					,	Mosko
guestion					Allswei	Mar NS
2(a)(i)	difference in conc. D	alD	$\left(\frac{\overline{M}}{\overline{M}}\right)$ bol	[X] gol		8
	24.04	120.20	2.08	-0.02		
	24.31	97.24	1.99	-0.16		
	24.40	81.33	1.91	-0.22		
	24.59	70.26	1.85	-0.39		
	24.67	61.68	1.79	-0.48		
	24.73	54.96	1.74	-0.57		
	24.77	49.54	1.69	-0.64		
	24.80	45.09	1.65	-0.70		
	24.83	41.38	1.62	-0.77		
	D data correc log [X] data o	x AND to 2 d orrect AND to	D data correct AND to 2 decimal places; log [X] data correct AND to 2 decimal places;	s ; ılaces ;		
2(a)(ii)	final concentration of blue dye / [X]	ration of blue	dye / [X]			_
2(a)(iii)	as mass of cl greater surfa	harcoal incre ce area leads	as mass of charcoal increases the final concentr greater surface area leads to higher adsorption ;	l concentrati Isorption ;	as mass of charcoal increases the final concentration of the dye decreases ; greater surface area leads to higher adsorption ;	2
2(b)	all nine points plotted correctly; (straight) line of best fit drawn;	s plotted corr of best fit dra	ectly ; awn ;			2
2(c)	point (at -0.2 charcoal too	2, 1.91) iden low / surface	tified AND sta area not high	atement exp ı enough / to	point (at -0.22, 1.91) identified AND statement explaining lack of adsorption, e.g. not enough stirring / mass of activated charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal too low / surface area not high enough / too low / coagulation of charcoal / bulkier particles used / not left long enough	_
2(d)(i)	coordinates read and recorded correctly (mus difference between the y coordinates must be difference between the x coordinates must be	ead and recc tween the y c tween the x c	coordinates read and recorded correctly (mus difference between the y coordinates must be difference between the x coordinates must be		coordinates read and recorded correctly (must not be points from table unless on the line); difference between the y coordinates must be at least half the range of the y values (0.3) AND difference between the x coordinates must be at least half the range of the x values (0.45)	2
	gradient dete	rmined (expe	gradient determined (expected value = 0.625)	0.625);		
2(d)(ii)	intercept on y-axis read and recorded correct	/-axis read aı	nd recorded c	correctly (ex	y (expected value = 2.09–2.10)	_

© UCLES 2019 Page 8 of 8