## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

## MARK SCHEME for the October/November 2009 question paper for the guidance of teachers

## 9701 CHEMISTRY

9701/32

Paper 32 (Advanced Practical Skills), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	ge 2 Mark Scheme: Teachers' version		Paper
	GCE A/AS LEVEL – October/November 2009	9701	32

Question	Sections	Indicative material	Mark	
1 (a)	PDO Recording	Appropriate headings and units Balance readings <u>consistent</u> to 1 dp, to 2 dp or to 3 dp according to precision of balance used Examiner to check subtraction – do not penalise any error here – see section <b>I(d)</b>	1	[2]
(b)				
(c)	MMO Collection  PDO Recording  MMO Quality	<ul> <li>(i) Total volume and temperature recorded for at least 2 different concentrations</li> <li>(ii) Total volume increased by 2.00 cm³ (±0.10 cm³) for at least 3 different concentrations of solution Volumes must be recorded to 2 dp.</li> <li>(iii) Appropriate headings and units</li> <li>(iv) All temperatures recorded to 0.5 °C.</li> <li>Check mass of FB 1 on Supervisor and candidate scripts, particularly if Supervisor and candidate temperatures differ greatly.</li> <li>Round temperatures to nearest 0.5°C if necessary.</li> <li>Where there are repeated temperature readings, take the average correct to</li> <li>1 decimal place (do not round).</li> <li>(v) &amp; (vi) .</li> <li>Compare temperature at 14.00 cm³ with temperature recorded by the Supervisor.</li> <li>Award (v) &amp; (vi) if within 2.5 °C</li> <li>Award (vi) only if temperature difference is &gt; 2.5 °C and ≤ 5 °C</li> <li>(vii) &amp; (viii)</li> <li>Compare temperature at 16.00 cm³ with temperature recorded by the Supervisor.</li> <li>Award (vii) &amp; (viii) if within 2.5 °C</li> <li>Award (viii) only if temperature difference is &gt; 2.5 °C and ≤ 5 °C</li> </ul>	1 1 1 1 2 2	[8]

Acceptable form of units – illustrated for mass.

using the solidus. / g
 using brackets. (g)

fully in words.
 composite.
 mass of ...... in grams
 mass of ..... in g

Where the unit is not incorporated into the heading it must appear after every data entry.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	32

Question	Sections	Indicative material	Mark	
(d)	ACE Interpretation	Examiner to check solubility calculated for each volume of water. Use candidate's mass of <b>FB 1</b> from <b>(a)</b> .	2	
if cand tabulat and,	e of water	Correct rounding required for sig fig displayed. Give 2 marks if <u>all four</u> solubilities are correct. Give 1 of these 2 marks if <b>three</b> only are correct.  Deduct 1 mark from those to be awarded if  1. there is an error in subtraction in (a).  OR  2. the mass recorded in (a) is less than 10 g or		
	PDO Display	more than 30 g.  No negative marks.  3 or 4 significant figures given in each answer given by the candidate	1	[3]
(e)	PDO Layout	Plots solubility on <i>y</i> -axis and temperature on <i>x</i> -axis.  Axes labelled (ignore incorrect or absent units) and points plotted over ½ of each axis - (count large squares containing the points).  Consider	1	
solubili volume the foll	idate plots ity against e of water owing only may be	<ol> <li>Spread of the 4 experimental points alone or</li> <li>Spread of the 4 experimental points and 42.5°C even if it is not possible to plot the point at 42.5°C on the y-axis</li> </ol>		
awarde L5 L6 L3		Scales used are easy for examiner to use and enable all points to be plotted 4 °C/large square is acceptable. Penalise non-linear scales here.	1	
	aximum for ction is 3	Correctly plots all solubility/temperature points.  Plotting to within ½ small square – in the correct square.  Attempts to draw a straight line or curve through plotted	1	
		points (minimum of 3 points)  Plots appropriate curve through a minimum of 4 points.  This is a mark for the quality of the line drawn and should be a <u>close</u> match to the plotted points.  Look for:  (i) good spread of points;	1	
	ACE Interpretation	<ul> <li>(ii) even distribution of points either side of the line (3 on line, 1 off, OK);</li> <li>(iii) smooth curve or regular straight line</li> <li>The solubility of FB 1 at 42.5 °C (± 1/4 square) read from the graph.</li> <li>Correct solubility recorded or correct solubility rounded to nearest gram recorded</li> </ul>	1	[6]
(f)	ACE Conclusions	Give 1 mark for stating there is a higher solubility at higher temperature	1	[1]

Page 4	ge 4 Mark Scheme: Teachers' version		Paper
	GCE A/AS LEVEL – October/November 2009	9701	32

Question	Sections	Indicative material	Mark	
(g)	ACE Conclusions	States that process is endothermic. Uses knowledge of equilibria to explain why it is endothermic. e.g. Refers to absorption of <a href="heat energy">heat energy</a> to dissolve potassium nitrate or States that a rise in temperature shifts equilibrium to the right. (or words to that effect). Beware answers that simply repeat information given in (f).	1	[2]
(h)	ACE Interpretation	Give 1 mark for each acceptable source of inaccuracy.  Evaporation of water;  Supercooling;  Thermal time lag in thermometer;  Difficulty in seeing crystals form;  Time delay in seeing crystals and reading thermometer;  Some solid forms on side of tube – not initial  FB 1;  Precision of thermometer;  not human error	2	[2]
(i)	ACE Improvement PDO Display	Outlines steps necessary to determine solubility, (weigh solution, evaporate, weigh residue)  Shows working for calculating the solubility.  mass of solid (mass of solution - mass of solid)  or  mass of solid mass of water  x 100	1	[2]
Qn 1	Total			[26]

Page 5	Page 5 Mark Scheme: Teachers' version		Paper
	GCE A/AS LEVEL – October/November 2009	9701	32

Question	Sections	Indicative material	Mark	
	FB 2 is Fe <sub>2</sub> (SO	$(a_4)_3(s)$ ; <b>FB 3</b> is ZnO(s)/Na <sub>2</sub> SO <sub>3</sub> (s); <b>FB 4</b> is CaSO <sub>3</sub> (s) [Na <sub>2</sub> SO <sub>3</sub> (s)]	93]	
2 (a)	MMO Collection MMO Decisions	Observes: colourless solution with <b>FB 4</b> , yellow, orange or brown solution with <b>FB 2</b> . Describes <u>one</u> of the following <u>gas tests</u> : (i) Potassium dichromate(VI) used in testing for SO <sub>2</sub> with <b>FB 3</b> or <b>FB 4</b> . (orange to green in colour); (ii) Blue litmus turning red on warming with hydrochloric acid; (iii) Limewater used to test gas – no change in the limewater.	1	[2]
(b)	PDO Recording MMO Collection	<ul> <li>(i) All observations in a single table. There must be no repetition of "headings".</li> <li>(ii) Reports addition of reagents to excess whenever a precipitate is formed on first addition of the reagent.</li> <li>(iii) and (iv)  Give 2 marks for correct observations for all 3 cations with both reagents  Give 1 of these 2 marks for 2 of the three observations correct.  FB 2 – red brown or brown ppt (insol in excess not required but con if either ppt is soluble in excess) with both reagents  FB 3 – white ppt soluble in excess with both reagents  FB 4 – no ppt with either reagent.</li> <li>(v) Give one mark for conclusions that match observations.  minimium observation required  FB 2 – Fe<sup>3+</sup>; red-brown/brown colour of each precipitate  FB 3 – Zn<sup>2+</sup>; white ppt with each reagent and ppt with NH<sub>3</sub> soluble in excess of aqueous ammonia  FB 4 – two of Ba<sup>2+</sup> or Ca<sup>2+</sup> or NH<sub>4</sub><sup>+</sup> from no ppt with either reagent</li> <li>If conclusions fit observations for more than one pair of ions – allow this mark.</li> </ul>	1 2 1	[5]
(c)	MMO Decisions	Selects appropriate reagent to distinguish between <a href="mailto:any">any</a> pair of cations identified in (b).  For Ba <sup>2+</sup> /Ca <sup>2+</sup> – accept H <sub>2</sub> SO <sub>4</sub> or chromate/dichromate;  For Ba <sup>2+</sup> /NH <sub>4</sub> <sup>+</sup> – accept H <sub>2</sub> SO <sub>4</sub> or chromate/dichromate  or  gas turns red litmus blue with NaOH (warming not required);  For Ca <sup>2+</sup> /NH <sub>4</sub> <sup>+</sup> – accept gas turns red litmus blue with NaOH (warming not required)	1	[1]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper	
	GCE A/AS LEVEL – October/November 2009	9701	32	

Question	Sections	Indicative material	Mark	
(d)	MMO Collection  ACE Conclusion	Observes white ppt with BaCl <sub>2</sub> , soluble in HCl with <b>FB 3</b> and <b>FB 4</b> . Observes white; off-white or cream (but not yellow) ppt with BaCl <sub>2</sub> , and <b>FB 2</b> and ppt insoluble on adding acid (ignore any colour change in the precipitate) Conclusion matching observations. Expected ions - sulfate in <b>FB 2</b> , sulfite in <b>FB 3</b> and <b>FB 4</b> . Formulae must be correct.	1 1 1	[3]
(e)	ACE Conclusion	Explains that $BaCl_2$ must be added first to form the precipitate (whose solubility is then tested with acid) or Addition of $BaCl_2$ to acidified solution will give no ppt even if sulfite present. Allow wtte.	1	[1]
(f)	MMO Collection	Observes <b>one</b> of the following when aqueous potassium iodide is added to the solution of <b>FB 2.</b> A darker solution than in <b>(a)</b> if there is no reference to colour change;  Acceptable solution colours are in the range: yellow, orange, red-brown, brown (but not red).  accept red-brown or brown solution formed; accept grey or black precipitate <b>AND</b> blue, blue-black, black, purple colour (solution or solid) when starch is added.  Identifies <b>FB 2</b> as an oxidant/oxidising agent or iodide as a reductant/reducing agent or converse <b>or</b> states that it is a redox reaction.  This mark can be given from either observation.	1	[2]
Qn 2	Total	<u> </u>		[14]