UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2010 question paper for the guidance of teachers

9702 PHYSICS

9702/35

Paper 31 (Advanced Practical Skills 1), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



(a)	(i) Value of d to the nearest 0.01 mm or 0.001 mm with consistent unit. 0.20 < d < 0.60 mm.	[1]
(b)	(iii) Value of <i>x</i> in range 40 cm–60 cm with consistent unit. Value of <i>I</i> with units.	[1]
(c)	Six sets of readings of x and I scores 5 marks, five sets scores 4 marks etc. Incorrect trend then -1 . Minor help from supervisor -1 ; major help from supervisor -2	[5]
	Range $x_{max} > 70$ cm; $x_{min} < 30$ cm	[1]
	Column headings Each column heading must contain a quantity and a unit. There must be some distinguishing mark between the quantity and the unit (solidus is expected but accept, for example, $1/I$ (A ⁻¹). Do not allow $1/I$ (A))	[1]
	Consistency of presentation of <u>raw</u> readings. All values of <i>x</i> must be given to the nearest mm.	[1]
	Significant figures S.F. in $1/I$ must be the same as, or one more than, the least number of significant figures used in raw I .	[1]
	Calculation	[1]

Mark Scheme: Teachers' version

GCE AS/A LEVEL – October/November 2010

Syllabus

9702

Paper

35

Page 2

Correct calculation of 1/I.

1

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper	
	GCE AS/A LEVEL – October/November 2010	9702	35	
(d) (i) Axes Sensible scales must be used. Awkward scales (e.g. 3:10) are not allowed Scales must be chosen so that the plotted points occupy at least half the grid in both x and y directions. Scales must be labelled with the quantity which is being plotted. Ignore un Scale markings should be no more than three large squares apart.			ne graph	[1
All Do Rin	tting of points observations must be plotted on the grid. not accept blobs (points with diameter > 0.5 small squa g and check a suspect plot. ork to an accuracy of half a small square.	re).		[1
Juc The leng	e of best fit dge by the balance of at least 5 points about the candida ere must be an even distribution of points either side gth. es must not be kinked. Do not accept lines thicker than	of the line alon	_	[1
All	ality points in the table (minimum 5) must be plotted for this nts must be within 2 cm (to scale) in <i>x</i> direction of a stra		ored. All	[1
`´ The	adient e hypotenuse of the triangle must be at least half the ler th read-offs must be accurate to half a small square.	igth of the drawi	n line.	[1
Eith Che	ercept ner: eck correct read-off from a point on the line, and subs ad-off must be accurate to half a small square. Allow ed	•		[1

Or:

Check read-off of intercept directly from graph.

(e) Values obtained in (a)(ii) and (d)(iii) substituted correctly into equation: $\frac{M}{N} = \frac{\rho}{AR}$ [1]

Do not allow substitution methods to find M or N

Value for ρ in range: $1 \times 10^{-7} \Omega \,\mathrm{m} - 5 \times 10^{-6} \Omega \,\mathrm{m}$ with consistent unit. [1]

[Total: 20]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2010	9702	35

- 2 (a) (ii) Measurement of x to nearest mm. x < 15.0 cm with consistent unit. [1] -1 for supervisor's help.
 - **(b) (iii)** Measurement of θ (less than 90°) with unit. [1]
 - (iv) Absolute uncertainty in θ in the range 2°-10°.
 If repeated readings have been taken, then the uncertainty can be half the range.
 Correct method of calculation of percentage uncertainty.
 - (v) m = 50 g with consistent unit [1]
 - M = 60 g with consistent unit [1]
 - (vi) Correct calculation of m/M (0.83 or 0.833). No units. [1]
 - (c) Measurement of θ [1]
 - m = 40 g; M = 70 g [1]
 - Quality: $\theta_2 > \theta_1$ [1]
 - (d) (i) Correct calculation of two values of *k*. [1]
 - (ii) Justification of sf in k linked to θ , m and M [1]
 - (iii) Valid conclusion based on the calculated values of *k*. [1] Candidate must test against a stated criterion.

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2010	9702	35

(e) Identifying limitations (4 marks) and suggesting improvements (4 marks)

	(i) Limitations [4]	(ii) Improvements [4]	Do not credit
A	Two readings are not enough (to draw a conclusion.	Take more readings and plot a graph/calculate more <i>k</i> values (and compare).	Few readings. Take more readings and calculate average. Only one reading.
В	Difficult to balance with reason e.g. unstable or effect of fans/draughts/a.c.	Drill hole higher up/switch off fans/a.c./close windows.	Closed room.
С	Difficult to judge when wooden strip horizontal/parallel (to the bench).	Method of ensuring strip is horizontal/parallel to bench e.g. use a spirit level or metre rule(s) to measure height of both ends/sight against window. Allow detailed use of set square.	Strip not straight/parallel/ horizontal. Use set square.
D	Difficult keeping <i>x</i> constant/ weights move.	Method of fixing cotton loop to rule e.g. tape, glue.	
E	Difficult to measure θ because hard to judge vertical/movement of hand.	Use a plumb line/clamped ruler/clamp protractor.	Bigger protractor. Paper behind protractor.
F	Friction at pulley/between nail and wooden strip.	Use lubricant/method of reducing friction.	Friction. Better pulley/ smooth(er) string/thin(ner) string. Friction between string and pulley. Lubrication between string and pulley.
G	Mass (values) not accurate.	Use balance/method of weighing mass.	Weigh masses.

Do not credit 'parallax problems', 'use assistant' or references to sensors, computers or videos.

[Total: 20]