

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**GCE Advanced Level**

## **MARK SCHEME for the October/November 2013 series**

### **9701 CHEMISTRY**

**9701/42**

Paper 4 (A2 Structured Questions), maximum raw mark 100


This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42



8 e<sup>-</sup> around chlorine

[1]

1 H–electron (+) on the Cl<sup>-</sup> ion

[1]

3 covalent (oo) and one dative (oo) around N

[1]

[3]

(b) (i) it would react (with H<sub>2</sub>SO<sub>4</sub>)

[1]

(ii)  $\text{CaO} + \text{H}_2\text{O} \longrightarrow \text{Ca(OH)}_2$

[1]

(iii) CaO absorbs more water or CaO has greater affinity for water

[1]

[3]

(c) (i)  $2\text{Ca(NO}_3)_2 \longrightarrow 2\text{CaO} + 4\text{NO}_2 + \text{O}_2$

[1]

(ii) (Down the group, the nitrates)

become more stable/stability increases

[1]

because the size/radius of ion (M<sup>2+</sup>) increases

[1]

thus causing less polarisation/distortion  
of the anion/NO<sub>3</sub><sup>-</sup>/N-O bond




[1]

[4]

[Total: 10]

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

- 2 (a) (i) Si-Si bonds are weaker (than C-C bonds) [1]
- (ii) metallic (Sn) is weaker than (giant) covalent (Ge) [1]
- [2]



[4]

[Total: 6]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

- 3 (a) (i)  $\text{NH}_3 + \text{HZ} \longrightarrow \text{NH}_4^+ + \text{Z}^-$  [1]  
 $\text{CH}_3\text{OH} + \text{HZ} \longrightarrow \text{CH}_3\text{OH}_2^+ + \text{Z}^-$  [1]
- (ii)  $\text{NH}_3 + \text{B}^- \longrightarrow \text{NH}_2^- + \text{BH}$  [1]  
 $\text{CH}_3\text{OH} + \text{B}^- \longrightarrow \text{CH}_3\text{O}^- + \text{BH}$  [1]
- [4]
- (b) (i) a reaction that can go in either direction [1]
- (ii) **rate** of forward = **rate** of backward reaction [1]  
or forward/back reactions occurring but concentrations of all species do not change [1]
- [2]
- (c) (i) a solution that resists changes in pH [1]  
when small quantities of acid or base/alkali are added [1]
- (ii) in the equilibrium system  $\text{HZ} + \text{H}_2\text{O} \rightleftharpoons \text{Z}^- + \text{H}_3\text{O}^+$  [1]  
addition of acid: reaction moves to the left  
or  $\text{H}^+$  combines with  $\text{Z}^-$  **and** forms  $\text{HZ}$  [1]  
addition of base: the reaction moves to the right  
or  $\text{H}^+$  combines with  $\text{OH}^-$  **and** more  $\text{Z}^-$  formed [1]
- [5 max 4]
- (d) (i)  $[\text{H}^+] = \sqrt{0.5 \times 1.34 \times 10^{-5}} = 2.59 \times 10^{-3} \text{ (mol dm}^{-3}\text{)}$  [1]  
pH = **2.59/2.6** (min 1 d.p) ecf [1]
- (ii)  $\text{CH}_3\text{CH}_2\text{CO}_2\text{H} + \text{NaOH} \longrightarrow \text{CH}_3\text{CH}_2\text{CO}_2\text{Na} + \text{H}_2\text{O}$  [1]
- (iii)  $n(\text{acid})$  in  $100 \text{ cm}^3 = 0.5 \times 100/1000 = 0.05 \text{ mol}$   
 $n(\text{acid})$  remaining =  $0.05 - 0.03 = 0.02 \text{ mol}$   
 $[\text{acid remaining}] = \mathbf{0.2} \text{ (mol dm}^{-3}\text{)}$  [1]  
likewise,  $n(\text{salt}) = 0.03 \text{ mol}$   
 $[\text{salt}] = \mathbf{0.3} \text{ (mol dm}^{-3}\text{)}$  [1]
- (iv)  $\text{pH} = 4.87 + \log(0.3/0.2) = \mathbf{5.04-5.05}$  ecf [1]
- [6]
- (e) **G** is  $\text{CH}_3\text{CH}_2\text{COCl}$   
**H** is  $\text{SOCl}_2$  or  $\text{PCl}_5$   
**J** is  $\text{NaCl}$  [2]  
(or corresponding Br compounds for **G**, **H** and **J**;  $\text{CH}_3\text{CH}_2\text{COBr}$ ,  $\text{SOBr}_2$ ,  $\text{NaBr}$ )

[Total: 18]

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

- 4 (a) (the energy change) when 1 mol of bonds is broken in the gas phase [1]  
[1]

[2]

- (b) (i) (C-X bond energy) decreases/becomes weaker (from F to I) [1]

due to bond becoming longer/not such efficient orbital overlap [1]

- (ii) (as the bond energy of C-X decreases) the halogenalkanes become more reactive (answer must imply that it is from F to I) [1]

[3]


- (c) The C-Cl bond is weaker than the C-F and C-H bonds or C-Cl bond (E = 340) and C-H (E = 410) [1]

so is (easily) broken to form Cl<sup>•</sup>/Cl radicals/Cl atoms [1]

causing the breakdown of O<sub>3</sub> into O<sub>2</sub> [1]

[3]

- (d) Cl-CH<sub>2</sub>CH<sub>2</sub>-CO<sub>2</sub>H [1]  
HO-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-Cl [1]



[1]

[3]

- (e) (i) light/UV/hν or 300°C [1]

- (ii) (free) radical substitution [1]

- (iii)  $\Delta H = E(\text{C-H}) - E(\text{H-Cl}) = 410 - 431 = -21 \text{ kJ mol}^{-1}$  [1]

- (iv)  $\Delta H = E(\text{C-H}) - E(\text{H-I}) = 410 - 299 = +111 \text{ kJ mol}^{-1}$  ecf [1]

- (v) The reaction with iodine is endothermic or  $\Delta H$  is positive or requires energy [1]

- (vi)  $\text{Cl}_2 \longrightarrow 2\text{Cl}^{\bullet}$  [1]



[8]

[Total: 19]

Page 6	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

5 (a) (i) **many** monomers form a polymer [1]


(ii) addition [1]

(iii) C=C/double/ $\pi$  bond is broken **and** new C-C single bonds s are formed  
or double bond breaks and forms single bonds with other monomers [1]

[3]

(b) propenoic acid [1]


[1]



carbon chain **and** CO<sub>2</sub>H  
**at least** one sodium salt [1]  
[1]

(ii) 120° to 109(.5)° [1]  
due to the change from a trigonal/sp<sup>2</sup> carbon to a tetrahedral/sp<sup>3</sup> carbon [1]

[4]



Any four:

hydrogen bond **labelled**

water H-bonded to O through H atom

$\delta^+/\delta^-$  shown on each end of a H-bond

lone pair shown on O<sup>-</sup> or C=O or H<sub>2</sub>O on a **correct H-bond**

Na<sup>+</sup> shown as coordinated to a water molecule [3]

(ii) Solution became paler **and** Cu<sup>(2+)</sup> swapped with Na<sup>(+)</sup>  
or darker in colour **and** polymer absorbs water [1]

[4]

<b>Page 7</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – October/November 2013</b>	<b>9701</b>	<b>42</b>


- (e) (i) alkene(1), amide(1) [2]
- (ii)  $\text{NH}_3$  [1]
- (iii)  $\text{H}_2\text{O}$  [1]
- (iv)  $\text{HCl}$  (aq)/ $\text{H}_3\text{O}^+$  **and** heat/reflux (**not** warm) [1]  
or  $\text{OH}^-$  (aq), heat and acidify [5]

**[Total: 17]**

Page 8	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

### Section B

6 (a) (i) **six/6** (gsv, sgv, gvs, vgs, svg, vsg) [1]



two **displayed** peptide bonds [1]  
correct formula of peptide [1]

(iii) valine (**allow** glycine) [1]


(iv) *any two of:*  
hydrogen bonds **and**  $\text{CO}_2\text{H}$  or  $\text{OH}$  or  $\text{NH}_2$  or  $\text{CONH}$  or  $\text{CO}$  or  $\text{NH}$  or  $\text{CO}_2^-$   
ionic bonds **and**  $\text{NH}_3^+$  or  $\text{CO}_2^-$   
van der Waals' **and**  $-\text{CH}_3$  or  $-\text{H}$  2 × [1]

[6]

(b) (i) same shape/structure as substrate [1]

(inhibitor) competes/blocks/binds/bonds to **active site**  
or substrate cannot bind to **active site** [1]

(ii) binds with enzyme **and** changes shape/3D structure (of enzyme/active site) [1]



[1]


[4]

[Total: 10]



Page 9	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	42

7 (a)



power supply (idea of complete circuit)  
 electrolyte/buffer solution  
 gel/filter paper/absorbent paper  
 (amino acid) sample/mixture [centre of plate]

4 × [1]

[4]

(b) any two from:

size/ $M_r$  (of the amino acid species)  
 charge (on the amino acid species)  
 temperature

2 × [1]

[2]

(c) Ratio of the concentration of a solute in each of two (immiscible) solvents  
 or equilibrium constant representing the distribution of a solute between two solvents  
 or  $PC = [X]_a/[X]_b$  (at a constant temperature)

[1]

[1]

(d) (i)  $K_{pc} = [Z \text{ in ether}]/[Z \text{ in } H_2O]$  – allow reverse ratio  
 $40 = (x/0.05)/((4-x)/0.5)$

[1]

= 3.2 g

ecf [1]

(ii) First extraction

$40 = (x/0.025)/((4-x)/0.5)$

x = 2.67 g

ecf [1]

(iii) Second extraction: 1.33g remain in solution

Second extraction

$40 = (y/0.025)/((1.33-y)/0.5)$

y = 0.887 g

mass extracted = 2.67 + 0.89 = 3.56/3.6 g

ecf [1]

[4]

[Total: 11]

<b>Page 10</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – October/November 2013</b>	<b>9701</b>	<b>42</b>

- 8 (a) (i)** (nitrates are) soluble [1]
- (ii)** Ba<sup>(2+)</sup> **and** Pb<sup>(2+)</sup> [1]
- SO<sub>4</sub><sup>(2-)</sup> [1]
- BaCO<sub>3</sub>/PbCO<sub>3</sub>/CaSO<sub>4</sub> are insoluble [1]
- [4]**
- (b) (i)** fertilisers/animal manure [1]
- (ii)** washing powder/detergents/fertilisers/animal manure [1]
- (iii)** growth/production of algae/weeds/plants  
or eutrophication [1]
- [3]**
- (c) (i)** *any one of:*
- 2SO<sub>2</sub> + O<sub>2</sub> → 2SO<sub>3</sub> **and** SO<sub>3</sub> + H<sub>2</sub>O → H<sub>2</sub>SO<sub>4</sub>
- or SO<sub>2</sub> + NO<sub>2</sub> → SO<sub>3</sub> + NO **and** SO<sub>3</sub> + H<sub>2</sub>O → H<sub>2</sub>SO<sub>4</sub>
- or SO<sub>2</sub> + ½O<sub>2</sub> + H<sub>2</sub>O → H<sub>2</sub>SO<sub>4</sub> [1]
- (ii)** roasting sulfide ores/extraction of metals from sulfide ores [1]
- [2]**
- [Total: 9]**