

Improving Colorectal Cancer Screening in South Asian Americans



A Mixed Methods Study with a CBPR framework

**Ma Somsouk, MD, MAS (UCSF)
Scarlett Lin Gomez, PhD (CPIC)
Susan L. Ivey, MD, MHSA (UCB)**

**UCSF Prevention Research Center
(U48DP004998)**

State of colorectal cancer screening




Ma Somsouk, MD
Division of Gastroenterology

University of California, San Francisco


Why CRC screening works

- Affects men and women of all races, lifetime risk of 5% (1 in 20 people)
- Most colon cancers arise in preexisting colonic polyps
- Has a long natural history
- Identification and early removal of precursor lesions (polyps) prevents colon cancer




USPSTF Recommendations

A Benefit: Life-years gained per 1000 individuals screened




B Benefit: Colorectal cancer deaths averted per 1000 individuals screened




USPSTF Recommendations

C Harms: Complications (gastrointestinal and cardiovascular events) of colorectal cancer screening and follow-up testing per 1000 individuals screened^c



D Burden: Lifetime No. of colonoscopies per 1000 individuals screened



State of CRC Screening

- Screening landscape dominated by colonoscopy & FIT/FOBT (>95%)
- All tests are effective, importance of adherence
 - Fecal immunochemical test (FIT) \$ 22
 - Colonoscopy \$900
 - Less common – sigmoidoscopy, CT colonography, stool DNA+FIT
- Adherence higher in immigrant groups when offered non-invasive screening tests



Why study South Asians?



- Dramatic growth of South Asian populations in California
 - Largest group of South Asians in California is Asian Indian (nearly 600K)
 - Most rapidly increasing population in California
 - Bangladeshi increased by 157% over 10 years
- Colorectal cancer incidence not decreasing among South Asian males, increasing among South Asian females
 - Contrast to declining incidence and mortality trends in other major racial/ethnic groups

State of CRC Screening in South Asians



- CRC screening in South Asians reported to be lower at 41% (as low as 21% in some surveys)
- Less acculturated, lower income, or more recent migrants may have lower use of CRC screening
 - Medical mistrust may also impact screening
- South Asians appear less educated about CRC
- Major limitations in current literature
 - Data based on community-based samples
 - Data do not reflect diversity of South Asian community

Objectives



- Public Health Impact:

- To understand the knowledge, attitudes and behaviors of South Asian immigrants related to CRC screening
- To design more effective and more culturally and linguistically appropriate interventions that lead to greater adherence to CRC screening guidelines

Ultimately to reduce death from CRC

Aim 1: Project Qualitative Aims

Susan L Ivey, MD, MHSA; Winston Tseng, PhD



Conduct a CBPR, mixed methods study to better understand patient attitudes and knowledge, acceptability of CRC screening modalities, barriers and facilitators to screening and relationship to delayed detection, access and use of health services, and health information seeking. Aim 1:

- Convened an Advisory Board of providers, community members, patient
 - Reviewed focus group questions; gathered input on outreach
 - Reviewed analysis plan
- Conducted 10 KI interviews with local community members on CAB to:
 - Understand contextual and system-level factors
 - Assess interventions, strategies, messages; add 4 KI interviews with religious leaders
- Conducted 8 focus groups among South Asian men and women (4 Asian Indian and 4 Bangladeshi groups) to assess:
 - Perception/awareness of CRC risks and screening, CRC screening facilitators, communication barriers and strategies

Improving Colorectal Cancer Screening in South Asian Americans



**Quantitative Results for Analysis of Statewide
Behavioral Risk Factor and Cancer Registry Data**

**Scarlett Lin Gomez, Ph.D.
Salma Shariff-Marco, Ph.D.
Juan Yang, Ph.D., M.S.**

Cancer Prevention Institute of California

Aim 2: Project Quantitative Aim



Applied traditional regression and novel classification techniques to **identify subgroups at highest risks of low adherence to CRC screening and late stage diagnosis**

- Guidance from Advisory Board on key factors
- Identify high-risk populations with low adherence to CRC screening guidelines
 - California Health Interview Survey data
- Assessed patterns of advanced stage CRC
 - California Cancer Registry data

Patterns of CRC screening in South Asians




- To identify high-risk populations with low adherence to current CRC screening guidelines:
 - Used California Health Interview Survey (CHIS) data
 - Focused on health care utilization, insurance coverage, use of other types of preventive services, length of US residence, limited English proficiency, and health literacy

California Health Interview Survey (CHIS)




- A random-dial telephone survey on a wide range of health topics
 - 40,000 adult participants
 - Representative of California's diverse population
- Total N=459 South Asians age 50+ (2001-2009)
 - N=2053 age 18+
 - South Asians are defined as including Asian Indian, Pakistani, Bangladeshi, Nepalese, Sri Lankans, and Other South Asians
- Variables available for assessment
 - Age, gender, race/ethnicity
 - Socioeconomic status, insurance
 - Marital status, immigration (% life in the US), English proficiency
 - Use of preventive services, health status, body mass index, physical activity

Adherence* with CRC screening in South Asians



*Adherent=FOBT annually, flexible sigmoidoscopy every 5 years, double-contrast barium enema every 5 years, colonoscopy every 10 years

Adherence* with CRC screening in South Asians



*Adherent=FOBT annually, flexible sigmoidoscopy every 5 years, double-contrast barium enema every 5 years, colonoscopy every 10 years

Non-adherence to CRC screening guidelines using multivariable model

Characteristic	% non-adherence (N=236)	Association with non-adherence	
	Weighted %	Adjusted OR*	(95% CI)
Percent of life lived in US			
0-20%	65.06	1.00	
21-40%	56.78	1.04	(0.38- 2.82)
41-60%	41.51	0.49	(0.21- 1.16)
61-80%	30.61	0.41	(0.15- 1.15)
81%+	27.67	0.32	(0.06- 1.62)
Unknown (2001 data)	51.73	0.26	(0.01- 7.84)
<i>P-trend³</i>			0.02
Language used at home			
English	37.18	1.00	
Non-English, 1 only	66.50	2.95	(1.05- 8.29)
English + other	48.82	1.74	(0.86- 3.49)
BMI (Asian cut points)			
Underweight/Normal (<23)	56.45	1.00	
Overweight (23-<27.5)	48.40	0.64	(0.33- 1.27)
Obese (27.5+)	38.24	0.39	(0.16- 0.95)
<i>P-trend⁵</i>			0.04
Had flu shot in past 12 months			
Yes	33.01	1.00	
No	62.24	2.75	(1.50- 5.04)
Unknown	55.82	5.35	(0.16- 180.07)

*Model also adjusted for age, gender, insurance, high blood pressure, heart disease, and # of times visited doctor


Recursive Partitioning: an agnostic classification technique for identifying high-risk subgroups

Less likely to be non-adherent

More likely to be non-adherent

N (sample size of this group)
%sample (among total sample, $N/459$)
% non-adherent(among this group, number of
not up-to-date/ N)


Tree 1: Flu shot strongly linked to CRC screening




Weighted distribution of non-adherence by recursive partitioning group

Had flu shot?	Total N	Non-adherence (N=236)			
		N	Weighted %	(95% CI)	
Yes	193	65	33.01	(24.79	- 41.24)
No or Unk	266	158	61.12	(52.40	- 69.83)

Tree 2: Other than flu shot, additional high-risk subgroups were identified, especially Group 1 & Group 4



Tree 2: Other than flu shot, additional high-risk subgroups were identified, especially Group 1 & Group 4



Tree 2: Weighted distribution of non-adherence by partitioning group

Recursive Partitioning Group	Total N	Not up-to-date (N=236)		
		N	Weighted %	(95% CI)
Group 1	48	37	70.05	(50.63 - 89.48)
Group 2	117	73	62.19	(50.61 - 73.76)
Group 3	166	49	30.46	(21.61 - 39.31)
Group 4	17	14	89.88	(74.05 - 100.00)
Group 5	29	19	62.85	(40.58 - 85.11)
Group 6	82	31	35.60	(22.68 - 48.52)

Group 1: Doctor visit >1 year or never

Group 2: Doctor visit ≤ 1 year & Age ≤ 54

Group 3: Doctor visit ≤ 1 year & Age ≥ 55 & Percent life in US ≤ 40% or Unknown

Group 4: Doctor visit ≤ 1 year & Age ≥ 55 & Percent life in US ≥ 41% & Insurance/Age = Uninsured or <65&Public/Medi-Cal

Group 5: Doctor visit ≤ 1 year & Age ≥ 55 & Percent life in US ≥ 41% & Insurance/Age = Private, Public/Medicare or 65+&Public/Medi-Cal & Urbanicity=Suburban

Group 6: Doctor visit ≤ 1 year & Age ≥ 55 & Percent life in US ≥ 41% & Insurance/Age = Private, Public/Medic-Cal or 65+&Public/Medi-Cal & Urbanicity = Urban, Smaller City, Town or Rural

Key findings from CHIS analyses




- 51% of South Asian adults in California, 50 years and older, were adherent to CRC screening for 2001-2009.
- Factors associated with **adherence**:
 - Being obese (BMI 27.5+), relative to BMI<23
- Factors associated with **non-adherence**:
 - Lower percent life lived in the US (2.6x less adherent)
 - Speaking only non-English at home (3x less adherent)
 - Did not have a flu shot in past 12 months (nearly 3x less adherent)
- High-risk subgroups identified by recursive partitioning:
 - No recent doctor's visits
 - Recent doctor's visit, age ≥ 55 , not recent immigrant, uninsured or public insurance

Aim 2: Colorectal cancer stage at diagnosis



- Assessed stage at diagnosis (early v. late) among CRC patients:
 - Used California Cancer Registry data
 - Focused on patient sociodemographic, hospital, and neighborhood characteristics

CRC stage at diagnosis by race/ethnicity, California, 1988-2012



California Cancer Registry (CCR) Variables

- Patient sociodemographic characteristics

- Age at diagnosis
- Sex
- Marital status
- Race/ethnicity
- Nativity
- Health insurance

- Neighborhood (block group)

- Socioeconomic status
- Ethnic enclave (Asian, Hispanic)

- Year of diagnosis

- Registry reporting region

- Reporting hospital type

- NCI designated cancer center
- ACOS Teaching Program
- ACOS Comprehensive Community Cancer Program
- ACOS Community Cancer Program

Multivariable associations with late stage at diagnosis for South Asians in California, 1988-2012



	OR	95% CI	
<u>Age at diagnosis</u> (ref = 60-69)			
50-59	1.62	1.11	2.35
60-69	1.00		
70-79	1.00	0.68	1.45
80+	1.30	0.84	2.01
p-trend	0.78	0.44	1.40
<u>Sex</u> (ref = male)			
Female	1.14	0.87	1.50
<u>Marital status</u> (ref = married)			
Single	2.43	1.40	4.21
Separated/Divorced/Widowed	1.36	0.92	2.03
Unknown	0.70	0.37	1.32
<u>Nativity</u> (ref = US-born)			
Foreign-born	2.42	1.64	3.58
<u>Health insurance</u> (ref = private)			
Not insured, self-pay	0.87	0.39	1.98
Public	1.04	0.73	1.49
Medicare	1.06	0.64	1.75
Unknown	0.83	0.46	1.48

- Younger (age<60) patients 62% more likely to be diagnosed at late stage
- Single patients >2x more likely than married to be diagnosed at late stage
- Foreign-born >2x more likely than US-born to be diagnosed at late stage

OR = odds ratio (adjusted)
CI = confidence interval

Multivariable associations with late stage at diagnosis for South Asians in California, 1988-2012



	OR	95% CI	
<u>Neighborhood SES</u> (ref = Q5 – high SES)			
Quintile 1- low SES	0.86	0.52	1.43
Quintile 2	0.82	0.54	1.25
Quintile 3	1.15	0.79	1.67
Quintile 4	1.08	0.77	1.52
p-trend			0.52
<u>Ethnic Enclave</u> (ref = Q1 – low enclave)			
Quintile 2	1.25	0.66	2.38
Quintile 3	1.39	0.75	2.60
Quintile 4	1.28	0.70	2.31
Quintile 5 - high enclave	0.86	0.48	1.53
Unknown	0.41	0.04	3.95
p-trend			0.06
<u>Reporting hospital type: NCI-Designated Cancer Center</u> (ref = no)			
Yes	1.20	0.73	1.96
<u>Year of Diagnosis (continuous)</u>	0.98	0.96	1.01


- No associations with neighborhood and hospital factors, nor year of diagnosis

OR = odds ratio (adjusted)
CI = confidence interval

Is late stage CRC incidence higher among South Asians?




In-situ and localized



Males



Regional and Distant



— NH White — API — South Asian

Females

— NH White — API — South Asian




Is late stage CRC incidence higher among South Asians?



In-situ and localized

Males

Regional and Distant



Key findings from CCR analysis



- 57% of South Asian adults in California, 50 years and older, were diagnosed with late stage (regional + distant) disease, comparable to other racial/ethnic groups.
- Factors associated with **late stage disease**:
 - Age < 60
 - Single marital status
 - Foreign-born
- High-risk subgroup identified by recursive partitioning:
 - Foreign-born

Aim 3: Health Communication and Outreach



- **Places** for information and outreach
 - Community-based organizations
 - Faith-based organizations
- **Preferred languages**
 - Both **native language and English** are needed
 - Should **reach underserved language groups**
- **Types** of outreach
 - Provide **health education classes** or **group discussions**
 - Host a booth at **community events**
 - Organize **speaker presentations**; participants will attend
 - People go to physicians/community/family/friends for information
- **Key messages** for outreach – consider other media
 - **Verbal messages** perceived as **more effective** than written messages

Next Steps in CBPR process



- Design materials and test usability with community members (In Progress: Summer – fall 2016)
- Create slide deck for physician-led outreach to community organizations (4 planned for fall 2016)
- Discuss with community partners additional outreach strategies (e.g., social media, digital storytelling, “champions”, etc.)

Community Advisory Board



- **Amit Agarwal, M.B.A.**
 - Principal, Deloitte;
Patient representative
- **Raisa Sultana**
 - General Secretary
Bay Area Bangladeshi Association
- **Pragati Grover, M.S.W.**
 - Executive Director
India Community Center
- **Alka Kanaya, M.D.**
 - Professor, UCSF
Department of Medicine
- **Suhaila Khan, M.D., Ph.D.**
 - Director of Research,
National Council of Asian Pacific
Islander Physicians
- **Mary Modayil, Ph.D., M.S.P.H.**
 - Epidemiologist
Alberta Health Services
- **Latha Palaniappan, M.D., M.Sc.**
 - Clinical Professor
Stanford University
- **Rachel J. Mesia, M.P.H.**
 - Manager, Cancer Clinical Trials and
Community Partnership Program
Stanford Cancer institute
- **Aparajita Singh, M.D.**
 - Assistant Clinical Professor, UCSF
Division of Gastroenterology
- **Ronesh Sinha, M.D.**
 - PRANA Initiative
Palo Alto Medical Foundation

Study Team



CANCER PREVENTION INSTITUTE
OF CALIFORNIA
Preventing Cancer. Promoting Life.



- UCSF
 - Ma Somsouk, M.D., M.A.S. (Contact PI)
 - Carly Rachocki, M.P.H.
- UC Berkeley – Health Research for Action
 - Susan L. Ivey, M.D., M.H.S.A. (Co-PI)
 - Winston Tseng, Ph.D.
- Cancer Prevention Institute of California
 - Scarlett Lin Gomez, Ph.D., M.P.H. (Co-PI)
 - Salma Shariff-Marco, Ph.D., M.P.H.
 - Juan Yang, Ph.D., M.P.H.
- Consultant:
 - Arnab Mukherjea, Dr.P.H., M.P.H.

Thanks for listening!



Contact us for additional information on the project

Ma Somsouk, MD, MAS – ma.somsouk@ucsf.edu

Susan L. Ivey, MD, MHSA – sivey@berkeley.edu

Scarlett Lin-Gomez, PhD - scarlett.gomez@cpic.org