

Dust Stars in the Minimal Exponential Measure (MEMe) Model

Justin C. Feng

CEICO, FZU, Institute of Physics of the Czech Academy of Sciences

In collaboration with Reyhan D. Lambaga, Norman Hsia, and Pisin Chen


Based on *Phys. Rev. D* 112, 024038 [*arXiv:2505.11591*]

Aug 5–9, 2025 Black Holes & Cosmology, University of Iceland, Reykjavik, Iceland



Co-funded by
the European Union





The Minimal Exponential Measure (MEMe) model

- Aims: maintain equivalence principle, no new dynamical dofs

$$\mathfrak{g}_{\mu\nu} = e^{(4-A^\mu{}_\mu)/2} A_\mu{}^\alpha A_\nu{}^\beta g_{\alpha\beta}, \quad \mathfrak{g} = \det(\mathfrak{g}..)$$

$$S[g.., A..] = \frac{1}{2\kappa} \int d^4x \left[\sqrt{|g|} (R - 2\tilde{\Lambda}) - \frac{2\kappa}{q} \sqrt{|\mathfrak{g}|} \right] + S_m[\varphi, \mathfrak{g}..]$$

- Field equations ($\kappa := 8\pi G/c^4$, $\tilde{\Lambda} := \Lambda + \frac{2\kappa}{q}$):

$$A_\beta{}^\alpha - \delta_\beta{}^\alpha = q \left[\frac{1}{4} \mathfrak{T} A_\beta{}^\alpha - \mathfrak{T}_{\beta\nu} \bar{\mathfrak{g}}^{\alpha\nu} \right]$$


$$G_{\mu\nu} + \left[\Lambda - \frac{\kappa}{q} (1 - |A|) \right] g_{\mu\nu} = \kappa |A| \bar{A}^\alpha{}_\mu \bar{A}^\beta{}_\nu \mathfrak{T}_{\alpha\beta}$$

- $A_\beta{}^\alpha$ nondynamical, coincides with GR in vacuum ($\mathfrak{T}_{\alpha\beta} = 0$) or vanishing q
- $1/q$ is characteristic energy density

¹Generalized coupling theories: JCF and S Carloni, Phys. Rev. D 101, 064002 (2020)

²JCF, S Mukohyama, S Carloni, Phys.Rev.D 103 (2021) 8, 084055; Phys.Rev.D 105 (2022) 10, 104036

Perfect fluid



EoS for $\hat{p} = 0, q > 0$.


A: $\hat{p} = 2/q$ **B:** $\hat{p} = 8/q$ **C:** $\hat{p} \rightarrow +\infty$

- Can get A_μ^α for: $\mathfrak{T}_{\mu\nu} = (\hat{\rho} + \hat{p})u_\mu u_\nu + \hat{p} g_{\mu\nu}$
 $G_{\mu\nu} = \kappa [(\rho + p)U_\mu U_\nu + p g_{\mu\nu}], \quad |A| \propto (1 - \hat{p}q)^3$
- Effective p, ρ , modified EoS

$$p = \frac{|A|(q\hat{p} - 1) + 1}{q} - \frac{\Lambda}{\kappa}, \quad \rho = |A|(\hat{p} + \hat{\rho}) - p$$
- Dust acquires pressure
- **EoS has maximum $\rho_{\max} = 7/(9q)$ for $q > 0$**
- For $q > 0$, can form compact objects from dust!


- For $q < 0$, find no dust stars for TOV

Dust star TOV solutions for $q > 0$



Mass function


$$[ds^2 = -e^{2\phi(r)} dt^2 + dr^2 / (1 - 2m(r)/r) + r^2 d\Omega^2]$$



Density profile

Nonmonotonic for central $\hat{\rho}_0 > 2/q$

Suppression of low mass BH formation



- Recall EoS for $q > 0$ has $\max \rho_{\max} = 7/(9q)$

- From Buchdahl limit,* $R > \frac{9}{4}M$, get collapse threshold density:

$$\rho_c = \frac{16}{243\pi M^2}$$

Avg. densities $\rho > \rho_c$ collapse to BHs.

- Maximum density in EoS places lower limit on mass that can be formed from collapse:


$$M \gtrsim M_{\min} = \frac{4}{9} \sqrt{\frac{3q}{7\pi}}$$

*H. A. Buchdahl, Phys. Rev. 116, 1027 (1959)

Summary

- MEMe model is simple modification of couplings between metric and matter DoFs
 - Equivalent to GR in vacuum
 - For a perfect fluid: GR with a modified equation of state
- For $q > 0$, can find compact dust star solutions
 - Can have solutions slightly larger than photon radius, potential BH mimickers
- Equation of state for $q > 0$ has a maximum density $\rho_{\text{max}} = 7/(9q)$
 - Maximum density places lower limit on mass that can be formed from collapse


Dust star properties



Scaling behavior for mass and radius of dust stars with a central density of $\hat{\rho}_0 = 1/q$

- For $\hat{\rho}_0 > 2/q$, density peaks at some $r > 0$
 - Expect a Rayleigh–Taylor instability
 - Radially unstable for $\hat{\rho}_0 > 2/q$
- For $\hat{\rho}q \ll 1$, dust EoS: $p \approx 3q\rho^2/8$
- For small $\hat{\rho}_0 \ll 1/q$, radius tied to scale $R \approx \pi\sqrt{3q/2\kappa}$ (since $p \approx 3q\rho^2/8$,)
 - Scaling holds for even for $\hat{\rho}_0 \sim 1/q$
 - Dust star size strongly tied to scale
- Dust stars can be compact; if formed from DM, can mimic low mass black holes.

Radial stability analysis



Lagrangian perturbation of pressure
at surface

- We follow procedure of Chandrasekhar and Chanmugam, with radial pert. Eqs in the comoving gauge

$$\begin{aligned} \frac{d\zeta}{dr} &= - \left(\frac{3}{r} + \frac{dp}{dr} \frac{1}{p + \rho} \right) \zeta - \frac{1}{r} \frac{\Delta p}{\Gamma p}, \\ \frac{d(\Delta p)}{dr} &= \zeta \left\{ \frac{(p + \rho)r\omega^2}{e^{2\phi}(1 - 2m/r)} - 4 \frac{dp}{dr} \right\} \\ &\quad + \zeta \left\{ \left(\frac{dp}{dr} \right)^2 \frac{r}{(p + \rho)} - \frac{8\pi(p + \rho)pr}{(1 - 2m/r)} \right\} \\ &\quad + \Delta p \left\{ \frac{dp}{dr} \frac{1}{p + \rho} - \frac{4\pi(p + \rho)r}{(1 - 2m/r)} \right\}. \end{aligned}$$

- Assuming regularity at $r = 0$, we solve perturbation equations for trial freqs. ω
- Solns. that satisfy $\Delta p = 0$ when $p = 0$ are eigenmodes.

Perfect fluids, high densities

- Perfect fluid: $\mathfrak{T}_{\mu\nu} = (\hat{\rho} + \hat{p})u_\mu u_\nu + \hat{p} \mathfrak{g}_{\mu\nu}$

can solve for A_μ^α :

$$A_\mu^\alpha = Y \delta_\mu^\alpha - 4(1 - Y) U_\mu U^\alpha$$

$$Y := \frac{4(1 - \hat{p}q)}{4 - q(3\hat{p} - \hat{\rho})}$$

- Determinant $|A|$

$$|A| = Y^3(4 - 3Y) = \frac{256(1 - \hat{p}q)^3(q\hat{\rho} + 1)}{[4 - q(3\hat{p} - \hat{\rho})]^4}$$

- If $q < 0$ then there is a crit. density $q\hat{\rho} = -1$,
If $q > 0$ then there is a crit. pressure $\hat{p}q = 1$,

$$G_{\mu\nu} \approx (\kappa/q)g_{\mu\nu}, \quad |\kappa/q| \gg |\Lambda|$$

- Can rewrite gravitational eqn:

$$G_{\mu\nu} = \kappa [(\rho + p)U_\mu U_\nu + p g_{\mu\nu}]$$

Effective fluid with pressure and density:

$$p = \frac{|A|(q\hat{p} - 1) + 1}{q} - \frac{\Lambda}{\kappa}$$

$$\rho = |A|(\hat{p} + \hat{\rho}) - p$$

- Tolman-Oppenheimer-Volkoff (TOV) equation:

$$p'(r) = -[p(r) + \rho(r)] \frac{m(r) + 4\pi r^3 p(r)}{[r - 2m(r)]r}$$

$$m'(r) = 4\pi \rho(r) r^2$$