

# Political Methodology III: Model Based Inference

Justin Grimmer

Professor  
Department of Political Science  
Stanford University

April 16th, 2019

# Most Important Problems

Effects of presidents going public

- $Y_i = 1$  if respondent identifies MIP as topic of president's speech
- $\mathbf{X}_i = (1, \text{Treatment}, \text{Republican})$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\mathbf{X}_i' \underbrace{\boldsymbol{\beta}}_{\text{Infer (Fixed)}})$$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\mathbf{X}_i' \underbrace{\boldsymbol{\beta}}_{\text{Infer (Fixed)}})$$

$$L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) = f(\mathbf{Y}|\boldsymbol{\beta}, \mathbf{X})$$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\mathbf{X}_i' \underbrace{\boldsymbol{\beta}}_{\text{Infer (Fixed)}})$$

$$\begin{aligned} L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= f(\mathbf{Y}|\boldsymbol{\beta}, \mathbf{X}) \\ &= \prod_{i=1}^N \Phi(\mathbf{X}_i' \boldsymbol{\beta})^{Y_i} (1 - \Phi(\mathbf{X}_i' \boldsymbol{\beta}))^{1-Y_i} \end{aligned}$$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\mathbf{X}_i' \underbrace{\boldsymbol{\beta}}_{\text{Infer (Fixed)}})$$

$$\begin{aligned} L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= f(\mathbf{Y}|\boldsymbol{\beta}, \mathbf{X}) \\ &= \prod_{i=1}^N \Phi(\mathbf{X}_i'\boldsymbol{\beta})^{Y_i} (1 - \Phi(\mathbf{X}_i'\boldsymbol{\beta}))^{1-Y_i} \\ \log L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left( Y_i \log \Phi(\mathbf{X}_i'\boldsymbol{\beta}) + (1 - Y_i) \log(1 - \Phi(\mathbf{X}_i'\boldsymbol{\beta})) \right) \end{aligned}$$

# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\mathbf{X}_i' \underbrace{\boldsymbol{\beta}}_{\text{Infer (Fixed)}})$$

$$\begin{aligned} L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= f(\mathbf{Y}|\boldsymbol{\beta}, \mathbf{X}) \\ &= \prod_{i=1}^N \Phi(\mathbf{X}_i'\boldsymbol{\beta})^{Y_i} (1 - \Phi(\mathbf{X}_i'\boldsymbol{\beta}))^{1-Y_i} \\ \log L(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left( Y_i \log \Phi(\mathbf{X}_i'\boldsymbol{\beta}) + (1 - Y_i) \log(1 - \Phi(\mathbf{X}_i'\boldsymbol{\beta})) \right) \\ \boldsymbol{\beta}^* &= \boldsymbol{\beta}^*(\mathbf{X}, \mathbf{Y}) = \text{Maximum Likelihood Estimator} \end{aligned}$$



# Review: Probit

$$\underbrace{Y_i}_{\text{Random}} \sim \text{Bernoulli}(\pi_i)$$
$$\pi_i = \Phi(\underbrace{X_i' \beta}_{\text{Infer (Fixed)}})$$

$$\begin{aligned} L(\beta | \mathbf{X}, \mathbf{Y}) &= f(\mathbf{Y} | \beta, \mathbf{X}) \\ &= \prod_{i=1}^N \Phi(X_i' \beta)^{Y_i} (1 - \Phi(X_i' \beta))^{1-Y_i} \\ \log L(\beta | \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left( Y_i \log \Phi(X_i' \beta) + (1 - Y_i) \log(1 - \Phi(X_i' \beta)) \right) \\ \beta^* &= \beta^*(\mathbf{X}, \mathbf{Y}) = \text{Maximum Likelihood Estimator} \\ \hat{\beta} &= \beta^*(\mathbf{X}, \mathbf{y}) \end{aligned}$$

# Review: Probit

$$\beta^* \rightarrow^D \text{Multivariate Normal} \left( \beta, I_N(\hat{\beta})^{-1} \right)$$

# Review: Probit

$$\beta^* \rightarrow^D \text{Multivariate Normal}(\beta, I_N(\hat{\beta})^{-1})$$

$$I_N(\hat{\beta})^{-1} = \begin{pmatrix} \text{var}(\beta_0) & \text{cov}(\beta_0, \beta_1) & \dots & \text{cov}(\beta_0, \beta_K) \\ \text{cov}(\beta_1, \beta_0) & \text{var}(\beta_1) & \dots & \text{cov}(\beta_1, \beta_K) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(\beta_K, \beta_0) & \text{cov}(\beta_K, \beta_1) & \dots & \text{var}(\beta_K) \end{pmatrix}$$

# Review Probit

# Review Probit

For a probit regression coefficient:

# Review Probit

For a probit regression coefficient:

$$CI_{95} = [\hat{\beta}_k - 1.96 \times \sqrt{\text{var}(\beta_k)}, \hat{\beta}_k + 1.96 \times \sqrt{\text{var}(\beta_k)}]$$

# Review Probit

For a probit regression coefficient:

$$\begin{aligned}\text{CI}_{95} &= [\hat{\beta}_k - 1.96 \times \sqrt{\text{var}(\beta_k)}, \hat{\beta}_k + 1.96 \times \sqrt{\text{var}(\beta_k)}] \\ \text{CI}_{1-\alpha} &= [\hat{\beta}_k - z_{\alpha/2} \times \sqrt{\text{var}(\beta_k)}, \hat{\beta}_k + z_{\alpha/2} \times \sqrt{\text{var}(\beta_k)}]\end{aligned}$$

# Review Probit

For a probit regression coefficient:

$$\begin{aligned}\text{CI}_{95} &= [\hat{\beta}_k - 1.96 \times \sqrt{\text{var}(\beta_k)}, \hat{\beta}_k + 1.96 \times \sqrt{\text{var}(\beta_k)}] \\ \text{CI}_{1-\alpha} &= [\hat{\beta}_k - z_{\alpha/2} \times \sqrt{\text{var}(\beta_k)}, \hat{\beta}_k + z_{\alpha/2} \times \sqrt{\text{var}(\beta_k)}]\end{aligned}$$

Recall: If  $Z \sim \text{Normal}(0, 1)$ ,  $z_{\alpha/2}$  is the number such that

$$P(|Z| \geq z_{\alpha/2}) = \alpha$$



# Review Probit

```
> colnames(mips)
```

```
[1] "mip" "treat" "gop"
```

```
> reg1<- glm(mip~treat + gop, family = binomial(link='probit')  
data = as.data.frame(mips))
```

```
> reg1$coef  
(Intercept) treat gop  
-0.259173644 -0.009598938 -0.198126517
```

```
> sqrt(diag(vcov(reg1)))  
(Intercept) treat gop  
0.03297588 0.05268254 0.05115777
```

# Review Probit

95-percent confidence interval:

```
> ci_treat<- c(reg1$coef[2] - 1.96sqrt(diag(vcov(reg1)))[2],  
reg1$coef[2] + 1.96sqrt(diag(vcov(reg1)))[2])
```

```
> ci_treat
```

```
-0.11285672 0.09365884
```

# Review Probit

Quantities of interest:

# Review Probit

Quantities of interest:

$$E[Y_i | \mathbf{X}_i] = \Phi(\mathbf{X}_i' \boldsymbol{\beta})$$

# Review Probit

Quantities of interest:

$$E[Y_i | \mathbf{X}_i] = \Phi(\mathbf{X}_i' \boldsymbol{\beta})$$

$$E[Y_i | X_{i1} = 1, \mathbf{X}_i] - E[Y_i | \tilde{X}_{i1} = 0, \tilde{\mathbf{X}}_i] = \Phi(\mathbf{X}_i' \boldsymbol{\beta}) - \Phi(\tilde{\mathbf{X}}_i' \boldsymbol{\beta})$$

# Review Probit

Quantities of interest:

$$\begin{aligned}E[Y_i|\mathbf{X}_i] &= \Phi(\mathbf{X}_i'\boldsymbol{\beta}) \\E[Y_i|X_{i1} = 1, \mathbf{X}_i] - E[Y_i|\tilde{X}_{i1} = 0, \tilde{\mathbf{X}}_i] &= \Phi(\mathbf{X}_i'\boldsymbol{\beta}) - \Phi(\tilde{\mathbf{X}}_i'\boldsymbol{\beta}) \\ \frac{\partial E[Y_i|X_{i1} = 1, \mathbf{X}_i]}{\partial X_{i1}} &= \phi(\mathbf{X}_i'\boldsymbol{\beta})\beta_1\end{aligned}$$

```
> X_synth<- c(1, 0, 1)

> y.tilde<- reg1$coef%*%X_synth

> y.prob<- pnorm(y.tilde)

> y.prob

[1,] 0.3237277
```

# Uncertainty for Maximum Likelihood Estimation



# Uncertainty for Maximum Likelihood Estimation

- 1) Coefficients: **Asymptotic result**  $\rightsquigarrow$  YMMV in small samples

# Uncertainty for Maximum Likelihood Estimation

- 1) Coefficients: **Asymptotic result**  $\rightsquigarrow$  YMMV in small samples
- 2) And even if asymptotic distribution holds:

# Uncertainty for Maximum Likelihood Estimation

- 1) Coefficients: **Asymptotic result**  $\rightsquigarrow$  YMMV in small samples
- 2) And even if asymptotic distribution holds:
  - 1) Difficult substantive interpretation of coefficients

# Uncertainty for Maximum Likelihood Estimation

- 1) Coefficients: **Asymptotic result**  $\rightsquigarrow$  YMMV in small samples
- 2) And even if asymptotic distribution holds:
  - 1) Difficult substantive interpretation of coefficients
  - 2) Uncertainty for **functions** of coefficients

# Inference Three Ways

How do obtain uncertainty estimates for Quantities of Interest?

- 1) Bootstrap  $\rightsquigarrow$  (no asymptotics, simulation)
- 2) Delta Method  $\rightsquigarrow$  (asymptotic normality, analytic)
- 3) Simulation from Multivariate Normal  $\rightsquigarrow$  (asymptotic normality, simulation)

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$



# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\beta^m = \beta^*(\tilde{Y}, \tilde{\mathbf{X}})$$

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\begin{aligned}\beta^m &= \beta^*(\tilde{Y}, \tilde{\mathbf{X}}) \\ h(\beta)^m &= h(\beta^*(\tilde{Y}, \tilde{\mathbf{X}}))\end{aligned}$$

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\begin{aligned}\beta^m &= \beta^*(\tilde{Y}, \tilde{\mathbf{X}}) \\ h(\beta)^m &= h(\beta^*(\tilde{Y}, \tilde{\mathbf{X}}))\end{aligned}$$

- 3) Do following:

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\begin{aligned}\beta^m &= \beta^*(\tilde{Y}, \tilde{\mathbf{X}}) \\ h(\beta)^m &= h(\beta^*(\tilde{Y}, \tilde{\mathbf{X}}))\end{aligned}$$

- 3) Do following:

$$E[\beta^*] \approx \sum_{m=1}^M \frac{\beta^m}{M}$$

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\begin{aligned}\beta^m &= \beta^*(\tilde{Y}, \tilde{\mathbf{X}}) \\ h(\beta)^m &= h(\beta^*(\tilde{Y}, \tilde{\mathbf{X}}))\end{aligned}$$

- 3) Do following:

$$\text{Var}(\beta^*) \approx \sum_{m=1}^M \frac{\left( \beta^m - \left( \sum_{m=1}^M \frac{\beta^m}{M} \right) \right)^2}{M}$$

# The Bootstrap (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*(\mathbf{X}, \mathbf{Y})$  for  $N$  observations

For each of  $M$  simulations ( $m = 1, 2, \dots, M$ ):

- 1) Sample **with replacement**  $N$  times:

$$(\tilde{Y}_1, \tilde{\mathbf{X}}_1), (\tilde{Y}_2, \tilde{\mathbf{X}}_2) \quad \dots \quad (\tilde{Y}_N, \tilde{\mathbf{X}}_N) = (\tilde{Y}, \tilde{\mathbf{X}})$$

- 2) Calculate  $\beta^*(\mathbf{X}, \mathbf{Y})$  and/or **any other function** of the estimator (expected value, first difference) on bootstrapped sample

$$\begin{aligned}\beta^m &= \beta^*(\tilde{Y}, \tilde{\mathbf{X}}) \\ h(\beta)^m &= h(\beta^*(\tilde{Y}, \tilde{\mathbf{X}}))\end{aligned}$$

- 3) Do following:

$$E[h(\beta^*)] \approx \sum_{m=1}^M \frac{h(\beta)^m}{M}$$

## Example:

```
store_expected<- c()
for(z in 1:1000){
  subset<- sample(1:nrow(mips), nrow(mips), replace=T)
  use_mips<- mips[subset,]
  temp_reg<- glm(mip~treat + gop,
    family = binomial(link='probit'),
    data = as.data.frame(use_mips))
  store_expected[z]<- pnorm(temp_reg$coef%*%X_synth)
}
> mean(store_expected) ## estimate of expected value
[1] 0.3245165
> sd(store_expected) ## estimate of standard error
[1] 0.0157724
```



# The Bootstrap: Justification

Two key ideas:

- 1) Simulation
- 2) Approximation of cumulative distribution function  $F$  with empirical distribution function  $\hat{F}_n$

# Simulation

- Sample  $M$  draws,  $Y_m \sim F$
- For example  $Y_m \sim \text{Normal}(\mu, \sigma^2)$

# Simulation

- Sample  $M$  draws,  $Y_m \sim F$
- For example  $Y_m \sim \text{Normal}(\mu, \sigma^2)$

$$\frac{1}{M} \sum_{m=1}^M Y_m \xrightarrow{P} E[Y] = \int y f(y) dy = \int y dF(y)$$

# Simulation

- Sample  $M$  draws,  $Y_m \sim F$
- For example  $Y_m \sim \text{Normal}(\mu, \sigma^2)$

$$\frac{1}{M} \sum_{m=1}^M Y_m \xrightarrow{P} E[Y] = \int y f(y) dy = \int y dF(y)$$

$$\frac{1}{M} \sum_{m=1}^M h(Y_m) \xrightarrow{P} E[h(Y)] = \int h(y) f(y) dy$$

# Simulation

- Sample  $M$  draws,  $Y_m \sim F$
- For example  $Y_m \sim \text{Normal}(\mu, \sigma^2)$

$$\frac{1}{M} \sum_{m=1}^M Y_m \xrightarrow{P} E[Y] = \int y f(y) dy = \int y dF(y)$$

$$\frac{1}{M} \sum_{m=1}^M h(Y_m) \xrightarrow{P} E[h(Y)] = \int h(y) f(y) dy$$

$$\frac{1}{M} \sum_{m=1}^M \left( Y_m - \sum_{m=1}^M \frac{Y_m}{M} \right)^2 \xrightarrow{P} \int (y - \mu)^2 f(y) dy$$

# Empirical Distribution Function

# Empirical Distribution Function

Univariate version:  $X_i$  is a scalar

$$\hat{F}_n(x) = \sum_{i=1}^N \frac{I(X_i \leq x)}{N}$$

# Empirical Distribution Function

Univariate version:  $X_i$  is a scalar

$$\hat{F}_n(x) = \sum_{i=1}^N \frac{I(X_i \leq x)}{N}$$

Multivariate Version:  $\mathbf{X}_i$  is a vector

$$\hat{F}_n(\mathbf{x}) = \sum_{i=1}^N \frac{I(\mathbf{X}_i \leq \mathbf{x})}{N}$$



# Empirical Distribution Function

Univariate version:  $X_i$  is a scalar

$$\hat{F}_n(x) = \sum_{i=1}^N \frac{I(X_i \leq x)}{N}$$

Multivariate Version:  $\mathbf{X}_i$  is a vector

$$\hat{F}_n(\mathbf{x}) = \sum_{i=1}^N \frac{I(\mathbf{X}_i \leq \mathbf{x})}{N}$$

Empirical density function: every observation has density height (probability)  $\frac{1}{N}$ .

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

$$\text{Var}_F(Y) = E[(Y - \mu)^2] = \int (y - \mu)^2 dF(y)$$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

$$\text{Var}_F(Y) = E[(Y - \mu)^2] = \int (y - \mu)^2 dF(y)$$

$$h_F(Y) = E[h(Y)] = \int h(y) dF(y)$$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

$$\text{Var}_F(Y) = E[(Y - \mu)^2] = \int (y - \mu)^2 dF(y)$$

$$h_F(Y) = E[h(Y)] = \int h(y) dF(y)$$

$$\text{Mean}_F(Y) \approx \text{Mean}_{\hat{F}}(Y) = \int y d\hat{F}(y)$$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

$$\text{Var}_F(Y) = E[(Y - \mu)^2] = \int (y - \mu)^2 dF(y)$$

$$h_F(Y) = E[h(Y)] = \int h(y) dF(y)$$

$$\text{Mean}_F(Y) \approx \text{Mean}_{\hat{F}}(Y) = \int y d\hat{F}(y)$$

$$\text{Var}_F(Y) \approx \text{Var}_{\hat{F}}(Y) = \int (y - \mu)^2 d\hat{F}(y)$$

# Approximation of Cumulative Distribution Function $F$

Expected value depends on  $F$

$$\text{Mean}_F(Y) = E[Y] = \int y dF(y)$$

$$\text{Var}_F(Y) = E[(Y - \mu)^2] = \int (y - \mu)^2 dF(y)$$

$$h_F(Y) = E[h(Y)] = \int h(y) dF(y)$$

$$\text{Mean}_F(Y) \approx \text{Mean}_{\hat{F}}(Y) = \int y d\hat{F}(y)$$

$$\text{Var}_F(Y) \approx \text{Var}_{\hat{F}}(Y) = \int (y - \mu)^2 d\hat{F}(y)$$

$$h_F(Y) \approx h_{\hat{F}}(Y) = \int h(y) d\hat{F}(y)$$



# How to Take Expected Values Using $\hat{F}_N$ ?

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

Bootstrap is a double approximation:

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

Bootstrap is a double approximation:

$$h_F(Y)$$

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

Bootstrap is a double approximation:

$$h_F(Y) \approx h_{\hat{F}}(Y)$$

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

Bootstrap is a double approximation:

$$h_F(Y) \approx h_{\hat{F}}(Y) \approx \sum_{m=1}^M \frac{h(Y_m)}{M}$$

# How to Take Expected Values Using $\hat{F}_N$ ?

Simulation!

Bootstrap is a double approximation:

$$h_F(Y) \approx h_{\hat{F}}(Y) \approx \sum_{m=1}^M \frac{h(Y_m)}{M}$$

$$h_F(\beta^*) \approx h_{\hat{F}}(\beta^*) \approx \sum_{m=1}^M \frac{h(\beta_m^*)}{M}$$

# Percentile Bootstrap Confidence Intervals

- Suppose we have  $M$  Bootstrap iterations of our maximum likelihood estimator  $\beta^*$ .
- Suppose  $h(\beta^*)$  is a scalar.
- Call  $h(\beta^*)_\alpha$  the value such that  $\alpha$  of values less than or equal to it. (For example  $h(\beta^*)_{0.5}$  is the **median** (value such that 50% lower)
- A  $1 - \alpha$  confidence interval (under some additional assumptions) is

$$CI_{1-\alpha} = (h(\beta^*)_{\alpha/2}, h(\beta^*)_{1-\alpha/2})$$



Recall:

```
store_expected<- c()
for(z in 1:1000){
  subset<- sample(1:nrow(mips), nrow(mips), replace=T)
  use_mips<- mips[subset,]
  temp_reg<- glm(mip~treat + gop,
    family = binomial(link='probit'),
    data = as.data.frame(use_mips))
  store_expected[z]<- pnorm(temp_reg$coef%*%X_synth)
}
>conf_int_95<- quantile(store_expected, c(0.025, 0.975))
> conf_int_95
2.5% 97.5%
0.2953548 0.3546646
```

# Bootstrapping is not a Panacea

# Bootstrapping is not a Panacea

- 1) If you have dependence in your data, you need to sample to preserve it (time series, nested data) (but you can do a block bootstrap)

# Bootstrapping is not a Panacea

- 1) If you have dependence in your data, you need to sample to preserve it (time series, nested data) (but you can do a block bootstrap)

# Bootstrapping is not a Panacea

- 1) If you have dependence in your data, you need to sample to preserve it (time series, nested data) (but you can do a block bootstrap)
- 2) Bootstrap doesn't always work: (but you can fix it, usually)
  - Matching
  - LASSO
  - Maximum (e.g. MLE for uniform)

# Inference Three Ways

- 1) Bootstrap  $\rightsquigarrow$  (no asymptotics, simulation)
- 2) Delta Method  $\rightsquigarrow$  (asymptotic normality, analytic)
- 3) Simulation from Multivariate Normal  $\rightsquigarrow$  (asymptotic normality, simulation)

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):



# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

- 1) Apply function  $h$  to maximum likelihood estimates:

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

- 1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

- 2) To calculate variance-covariance matrix:

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

- 1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

- 2) To calculate variance-covariance matrix:

- a) Calculate  $\nabla h(\beta^*)$  at maximum likelihood estimates:

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

2) To calculate variance-covariance matrix:

a) Calculate  $\nabla h(\beta^*)$  at maximum likelihood estimates:

$$\nabla h(\beta^*) = \left( \frac{\partial h(\beta)}{\partial \beta_1} \Big|_{\beta^*}, \frac{\partial h(\beta)}{\partial \beta_2} \Big|_{\beta^*}, \dots, \frac{\partial h(\beta)}{\partial \beta_K} \Big|_{\beta^*} \right)$$

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \xrightarrow{D} \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

2) To calculate variance-covariance matrix:

a) Calculate  $\nabla h(\beta^*)$  at maximum likelihood estimates:

$$\nabla h(\beta^*) = \left( \frac{\partial h(\beta)}{\partial \beta_1} \Big|_{\beta^*}, \frac{\partial h(\beta)}{\partial \beta_2} \Big|_{\beta^*}, \dots, \frac{\partial h(\beta)}{\partial \beta_K} \Big|_{\beta^*} \right)$$

b) Calculate Variance covariance matrix:



# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \longrightarrow^D \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

- 1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

- 2) To calculate variance-covariance matrix:

- a) Calculate  $\nabla h(\beta^*)$  at maximum likelihood estimates:

$$\nabla h(\beta^*) = \left( \frac{\partial h(\beta)}{\partial \beta_1} \Big|_{\beta^*}, \frac{\partial h(\beta)}{\partial \beta_2} \Big|_{\beta^*}, \dots, \frac{\partial h(\beta)}{\partial \beta_K} \Big|_{\beta^*} \right)$$

- b) Calculate Variance covariance matrix:

$$\nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*)$$

# Delta Method (Procedure, Then Justification)

Suppose we have maximum likelihood estimator  $\beta^*$ .

We are interested in  $h(\beta^*)$ . (Assuming a large sample):

$$h(\beta^*) \longrightarrow^D \text{Multivariate Normal} \left( h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*) \right)$$

Inference for  $h(\beta^*)$

1) Apply function  $h$  to maximum likelihood estimates:

$$h(\beta^*) = \Phi(\mathbf{X}_i' \beta^*)$$

2) To calculate variance-covariance matrix:

a) Calculate  $\nabla h(\beta^*)$  at maximum likelihood estimates:

$$\nabla h(\beta^*) = \left( \frac{\partial h(\beta)}{\partial \beta_1} \Big|_{\beta^*}, \frac{\partial h(\beta)}{\partial \beta_2} \Big|_{\beta^*}, \dots, \frac{\partial h(\beta)}{\partial \beta_K} \Big|_{\beta^*} \right)$$

b) Calculate Variance covariance matrix:

$$\nabla h(\beta^*)' (-E[\text{Hessian}]^{-1}) \nabla h(\beta^*)$$

Example:

$$\frac{\partial E[Y|\mathbf{X}_i, \boldsymbol{\beta}]}{\partial \beta_k} = \phi(\mathbf{X}_i \boldsymbol{\beta}) X_k$$

```
prob_grad<- function(X, coef){  
  base<- dnorm(coef%*%X)  
  out<- base*X  
  return(out)  
}  
  
> mle_grad<- prob_grad(X_synth, reg1$coef)  
  
> mle_grad  
[1] 0.359335 0.000000 0.359335  
  
> ses<- sqrt(t(mle_grad)%*%vcov(reg1)%*%mle_grad)  
  
> ses  
[1,] 0.01618305
```

# Comparison

	Bootstrap	Delta Method
Standard Error	0.016	0.016
Confidence Interval	[0.295, 0.355]	

# Delta Method (Justification)

Recall that

# Delta Method (Justification)

Recall that

$$\boldsymbol{\beta}^* \longrightarrow^D \text{Multivariate Normal } (\boldsymbol{\beta}, I_N(\boldsymbol{\beta}^*)^{-1})$$

# Delta Method (Justification)

Recall that

$$\beta^* \longrightarrow^D \text{Multivariate Normal } (\beta, I_N(\beta^*)^{-1})$$

Multivariate Taylor Series:

# Delta Method (Justification)

Recall that

$$\boldsymbol{\beta}^* \longrightarrow^D \text{Multivariate Normal } (\boldsymbol{\beta}, I_N(\boldsymbol{\beta}^*)^{-1})$$

Multivariate Taylor Series:

$$h(\boldsymbol{\beta}^*) = h(\boldsymbol{\beta}) + \nabla h(\boldsymbol{\beta})'(\boldsymbol{\beta}^* - \boldsymbol{\beta}) + \text{Remainder}$$



# Delta Method (Justification)

Recall that

$$\boldsymbol{\beta}^* \longrightarrow^D \text{Multivariate Normal } (\boldsymbol{\beta}, I_N(\boldsymbol{\beta}^*)^{-1})$$

Multivariate Taylor Series:

$$h(\boldsymbol{\beta}^*) = h(\boldsymbol{\beta}) + \nabla h(\boldsymbol{\beta})'(\boldsymbol{\beta}^* - \boldsymbol{\beta}) + \text{Remainder}$$

# Delta Method (Justification)

Take the limit:

$$\lim_{n \rightarrow \infty} h(\beta^*) = \lim_{n \rightarrow \infty} \left( h(\beta) + \nabla h(\beta)'(\beta^* - \beta) + \text{Remainder} \right)$$

$$\lim_{n \rightarrow \infty} h(\beta^*) \xrightarrow{P} h(\beta)$$

$$\lim_{n \rightarrow \infty} h(\beta^*) \xrightarrow{D} h(\beta) + \lim_{n \rightarrow \infty} \left( \nabla h(\beta)' \underbrace{(\beta^* - \beta)}_{\text{MVN}(0, I_N(\beta)^{-1})} \right)$$

# Delta Method (Justification)

Take the limit:

$$\lim_{n \rightarrow \infty} h(\beta^*) = \lim_{n \rightarrow \infty} \left( h(\beta) + \nabla h(\beta)' (\beta^* - \beta) + \text{Remainder} \right)$$

$$\lim_{n \rightarrow \infty} h(\beta^*) \xrightarrow{P} h(\beta)$$

$$\lim_{n \rightarrow \infty} h(\beta^*) \xrightarrow{D} h(\beta) + \lim_{n \rightarrow \infty} \left( \nabla h(\beta)' \underbrace{(\beta^* - \beta)}_{\text{MVN}(0, I_N(\beta)^{-1})} \right)$$

Then by Slutsky's Theorem:

$$h(\beta^*) \xrightarrow{D} \text{MVN}(h(\beta), \nabla h(\beta^*)' I_N(\beta^*)^{-1} \nabla h(\beta^*))$$

# Delta Method Confidence Intervals

- Suppose  $h(\beta^*)$  is a scalar
- $SE(h(\beta^*)) = \sqrt{(\nabla h(\beta^*))' I_N(\beta^*)^{-1} \nabla h(\beta^*)}$
- Then a  $1 - \alpha$  confidence interval is:

$$CI_{1-\alpha} = [h(\beta^*) - 1.96 \times SE(h(\beta^*)), h(\beta^*) + 1.96 \times SE(h(\beta^*))]$$

# Delta Method: Example

```
> y.prob<- pnorm(y.tilde)
> X_synth<- c(1, 0, 1)
> y.tilde<- reg1$coef%*%X_synth
> y.prob<- pnorm(y.tilde)
> y.prob
[1,] 0.3237277

> ses<- sqrt(t(mle_grad)%*%vcov(reg1)%*%mle_grad)
> ses
[1,] 0.01618305
> delta_95<- c(y.prob - 1.96*ses, y.prob + 1.96*ses)
> delta_95
[1] 0.2920089 0.3554464
```

# Comparison

	Bootstrap	Delta Method
Standard Error	0.016	0.016
Confidence Interval	[0.295, 0.355]	[0.292, 0.355]

# Inference Three Ways

- 1) Bootstrap  $\rightsquigarrow$  (no asymptotics, simulation)
- 2) Delta Method  $\rightsquigarrow$  (asymptotic normality, analytic)
- 3) Simulation from Multivariate Normal  $\rightsquigarrow$  (asymptotic normality, simulation)

# Simulation from Multivariate Normal (Procedure)

$$\boldsymbol{\beta}^* \longrightarrow^D \text{MVN}(\boldsymbol{\beta}, I(\boldsymbol{\beta})^{-1})$$



# Simulation from Multivariate Normal (Procedure)

$$\boldsymbol{\beta}^* \longrightarrow^D \text{MVN}(\boldsymbol{\beta}, I(\boldsymbol{\beta})^{-1})$$

Inference about  $h(\boldsymbol{\beta}^*)$ :

# Simulation from Multivariate Normal (Procedure)

$$\boldsymbol{\beta}^* \longrightarrow^D \text{MVN}(\boldsymbol{\beta}, I(\boldsymbol{\beta})^{-1})$$

Inference about  $h(\boldsymbol{\beta}^*)$ :

For each simulation  $m$  of  $M$  simulations:

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\beta^m \sim \text{MVN}(\beta^*, I_N(\beta)^{-1})$$

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\begin{aligned}\beta^m &\sim \text{MVN}(\beta^*, I_N(\beta)^{-1}) \\ h(\beta)^m &= h(\beta^m)\end{aligned}$$

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\begin{aligned}\beta^m &\sim \text{MVN}(\beta^*, I_N(\beta)^{-1}) \\ h(\beta)^m &= h(\beta^m)\end{aligned}$$

Use collection of simulated values  $(h(\beta)^1, h(\beta)^2, \dots, h(\beta)^M)$  to approximate distribution of  $h(\beta^*)$ .

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\begin{aligned}\beta^m &\sim \text{MVN}(\beta^*, I_N(\beta)^{-1}) \\ h(\beta)^m &= h(\beta^m)\end{aligned}$$

Use collection of simulated values  $(h(\beta)^1, h(\beta)^2, \dots, h(\beta)^M)$  to approximate distribution of  $h(\beta^*)$ .

$$E[h(\beta^*)] = \sum_{m=1}^M \frac{h(\beta)^m}{M}$$

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\begin{aligned}\beta^m &\sim \text{MVN}(\beta^*, I_N(\beta)^{-1}) \\ h(\beta)^m &= h(\beta^m)\end{aligned}$$

Use collection of simulated values  $(h(\beta)^1, h(\beta)^2, \dots, h(\beta)^M)$  to approximate distribution of  $h(\beta^*)$ .

$$\text{var}[h(\beta^*)] = \sum_{m=1}^M \frac{\left(h(\beta)^m - \sum_{m=1}^M \frac{h(\beta)^m}{M}\right)^2}{M}$$

# Simulation from Multivariate Normal (Procedure)

$$\beta^* \rightarrow^D \text{MVN}(\beta, I(\beta)^{-1})$$

Inference about  $h(\beta^*)$ :

For each simulation  $m$  of  $M$  simulations:

$$\begin{aligned}\beta^m &\sim \text{MVN}(\beta^*, I_N(\beta)^{-1}) \\ h(\beta)^m &= h(\beta^m)\end{aligned}$$

Use collection of simulated values  $(h(\beta)^1, h(\beta)^2, \dots, h(\beta)^M)$  to approximate distribution of  $h(\beta^*)$ .

$$f[h(\beta^*)] = \sum_{m=1}^M \frac{f(h(\beta)^m)}{M}$$



# Simulation: Example

```
library(MASS)
draw_coef<- mvrnorm(1000, mu = reg1$coef, Sigma = vcov(reg1))

> dim(draw_coef)
[1] 1000 3

dist_exp<- pnorm(draw_coef%*%X_synth)
> length(dist_exp)
[1] 1000

sd(dist_exp)
[1] 0.01572543
```

# Comparison

	Bootstrap	Delta Method	MVN Simulation
Standard Error	0.016	0.016	0.016
Confidence Interval	[0.295, 0.355]	[0.292, 0.355]	

# Simulation from Multivariate Normal (Justification)

Simulation is a (stronger assumption) double approximation

$$\begin{aligned} E_f[h(\boldsymbol{\beta})^*] &= \int h(\boldsymbol{\beta})^* f(\boldsymbol{\beta}^*) d\boldsymbol{\beta}^* \\ &\approx \int h(\boldsymbol{\beta}^*) \underbrace{\tilde{f}(\boldsymbol{\beta}^*)}_{\text{MVN}} d\boldsymbol{\beta}^* \approx \sum_{m=1}^M \frac{h(\boldsymbol{\beta})^m}{M} \end{aligned}$$

# Percentile Simulation Confidence Intervals

- Suppose we have  $M$  draws
- Suppose  $h(\beta^*)$  is a scalar
- Then a  $1 - \alpha$  confidence interval is:

$$CI_{1-\alpha} = [h(\beta^*)_{\alpha/2}, h(\beta^*)_{1-\alpha/2}]$$

- In practice: order the values, select quantile values at  $\alpha/2$  and  $1 - \alpha/2$ .

# Simulation Confidence Intervals

```
sim_95<- quantile(dist_exp, c(0.025, 0.975))  
> sim_95  
2.5% 97.5%  
0.2953568 0.3559635
```

# Comparison

	Bootstrap	Delta Method	MVN Simulation
Standard Error	0.016	0.016	0.016
Confidence Interval	[0.295, 0.355]	[0.292, 0.355]	[0.295, 0.356]

# So Which Should I Use?

## Lots of data

- Then all three methods will be close
- Delta Method + MVN Simulation  $\rightsquigarrow$  faster (but what else are you doing?)

## Not lots of data

- Then only bootstrap will perform well
- And will be comparable in speed to Delta + MVN

# Thursday

## Model fit for Binary DVs + Ordered Probit