

Political Methodology III: Model Based Inference

Justin Grimmer

Professor
Department of Political Science
Stanford University

May 21st, 2019

Model Based Inference

- 1) Likelihood inference
- 2) Machine Learning
 - a) Unsupervised Latent Features
 - b) Classification/Prediction/Regression

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2$$

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2$$
$$\hat{\boldsymbol{\beta}} = \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\}$$

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \\ \hat{\boldsymbol{\beta}} &= \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\} \\ &= \left(\mathbf{X}' \mathbf{X} \right)^{-1} \mathbf{X}' \mathbf{Y} \end{aligned}$$

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \\ \hat{\boldsymbol{\beta}} &= \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\} \\ &= \left(\mathbf{X}' \mathbf{X} \right)^{-1} \mathbf{X}' \mathbf{Y} \end{aligned}$$

Problem:

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \\ \hat{\boldsymbol{\beta}} &= \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\} \\ &= \left(\mathbf{X}' \mathbf{X} \right)^{-1} \mathbf{X}' \mathbf{Y} \end{aligned}$$

Problem:

- J will likely be large (perhaps $J > N$)

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \\ \hat{\boldsymbol{\beta}} &= \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\} \\ &= \left(\mathbf{X}' \mathbf{X} \right)^{-1} \mathbf{X}' \mathbf{Y} \end{aligned}$$

Problem:

- J will likely be large (perhaps $J > N$)
- There many correlated variables

Regression models

Suppose we have N documents, with each document i having label $y_i \in \{-1, 1\} \rightsquigarrow \{\text{not, credit claiming}\}$

We represent each document i is $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{iJ})$.

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \\ \hat{\boldsymbol{\beta}} &= \arg \min_{\boldsymbol{\beta}} \left\{ \sum_{i=1}^N \left(y_i - \boldsymbol{\beta}' \mathbf{x}_i \right)^2 \right\} \\ &= \left(\mathbf{X}' \mathbf{X} \right)^{-1} \mathbf{X}' \mathbf{Y} \end{aligned}$$

Problem:

- J will likely be large (perhaps $J > N$)
- There many correlated variables

Predictions will be **variable**

Mean Square Error

Suppose θ is some value of the true parameter

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$E[(\hat{\theta} - \theta)^2]$$

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$E[(\hat{\theta} - \theta)^2] = E[\hat{\theta}^2] - 2\theta E[\hat{\theta}] + \theta^2$$

Mean Square Error

Suppose θ is some value of the true parameter
Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$\begin{aligned} E[(\hat{\theta} - \theta)^2] &= E[\hat{\theta}^2] - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + E[\hat{\theta}]^2 - 2\theta E[\hat{\theta}] + \theta^2 \end{aligned}$$

Mean Square Error

Suppose θ is some value of the true parameter

Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$\begin{aligned} E[(\hat{\theta} - \theta)^2] &= E[\hat{\theta}^2] - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + E[\hat{\theta}]^2 - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + (E[\hat{\theta}] - \theta)^2 \end{aligned}$$

Mean Square Error

Suppose θ is some value of the true parameter

Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$\begin{aligned} E[(\hat{\theta} - \theta)^2] &= E[\hat{\theta}^2] - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + E[\hat{\theta}]^2 - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + (E[\hat{\theta}] - \theta)^2 \\ &= \text{Var}(\hat{\theta}) + \text{Bias}^2 \end{aligned}$$

Mean Square Error

Suppose θ is some value of the true parameter

Bias:

$$\text{Bias} = E[\hat{\theta} - \theta]$$

We may care about average distance from truth

$$\begin{aligned} E[(\hat{\theta} - \theta)^2] &= E[\hat{\theta}^2] - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + E[\hat{\theta}]^2 - 2\theta E[\hat{\theta}] + \theta^2 \\ &= E[\hat{\theta}^2] - E[\hat{\theta}]^2 + (E[\hat{\theta}] - \theta)^2 \\ &= \text{Var}(\hat{\theta}) + \text{Bias}^2 \end{aligned}$$

To reduce MSE, we are willing to induce bias to decrease variance \rightsquigarrow
methods that **shrink** coefficients toward zero

Ridge Regression

Penalty for model complexity

Ridge Regression

Penalty for model complexity

$$f(\beta, \mathbf{X}, \mathbf{Y})$$

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2$$

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \underbrace{\lambda \sum_{j=1}^J \beta_j^2}_{\text{Penalty}}$$

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \underbrace{\lambda \sum_{j=1}^J \beta_j^2}_{\text{Penalty}}$$

where:

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \underbrace{\lambda \sum_{j=1}^J \beta_j^2}_{\text{Penalty}}$$

where:

- $\beta_0 \rightsquigarrow$ intercept

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \underbrace{\lambda \sum_{j=1}^J \beta_j^2}_{\text{Penalty}}$$

where:

- $\beta_0 \rightsquigarrow$ intercept
- $\lambda \rightsquigarrow$ penalty parameter

Ridge Regression

Penalty for model complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \underbrace{\lambda \sum_{j=1}^J \beta_j^2}_{\text{Penalty}}$$

where:

- $\beta_0 \rightsquigarrow$ intercept
- $\lambda \rightsquigarrow$ penalty parameter
- Standardized \mathbf{X} (coefficients on same scale)

Ridge Regression \rightsquigarrow Optimization

$$\beta^{\text{Ridge}} = \arg \min_{\beta} \{f(\beta, \mathbf{X}, \mathbf{Y})\}$$

Ridge Regression \rightsquigarrow Optimization

$$\begin{aligned}\beta^{\text{Ridge}} &= \arg \min_{\beta} \{f(\beta, \mathbf{X}, \mathbf{Y})\} \\ &= \arg \min_{\beta} \left\{ \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \beta_j^2 \right\}\end{aligned}$$

Ridge Regression \rightsquigarrow Optimization

$$\begin{aligned}\beta^{\text{Ridge}} &= \arg \min_{\beta} \{f(\beta, \mathbf{X}, \mathbf{Y})\} \\ &= \arg \min_{\beta} \left\{ \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \beta_j^2 \right\} \\ &= \arg \min_{\beta} \left\{ (\mathbf{Y} - \mathbf{X}'\beta)'(\mathbf{Y} - \mathbf{X}'\beta) + \lambda \beta' \beta \right\}\end{aligned}$$

Demmean the data and set $\beta_0 = \bar{y} = \sum_{i=1}^N \frac{y_i}{N}$

Ridge Regression \rightsquigarrow Optimization

$$\begin{aligned}\beta^{\text{Ridge}} &= \arg \min_{\beta} \{f(\beta, \mathbf{X}, \mathbf{Y})\} \\ &= \arg \min_{\beta} \left\{ \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \beta_j^2 \right\} \\ &= \arg \min_{\beta} \left\{ (\mathbf{Y} - \mathbf{X}'\beta)'(\mathbf{Y} - \mathbf{X}'\beta) + \lambda \beta' \beta \right\} \\ &= \left(\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}'\mathbf{Y}\end{aligned}$$

Demean the data and set $\beta_0 = \bar{y} = \sum_{i=1}^N \frac{y_i}{N}$

Ridge Regression \rightsquigarrow Optimization

$$\begin{aligned}\beta^{\text{Ridge}} &= \arg \min_{\beta} \{f(\beta, \mathbf{X}, \mathbf{Y})\} \\ &= \arg \min_{\beta} \left\{ \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \beta_j^2 \right\} \\ &= \arg \min_{\beta} \left\{ (\mathbf{Y} - \mathbf{X}'\beta)'(\mathbf{Y} - \mathbf{X}'\beta) + \lambda \beta' \beta \right\} \\ &= \left(\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}'\mathbf{Y}\end{aligned}$$

Demmean the data and set $\beta_0 = \bar{y} = \sum_{i=1}^N \frac{y_i}{N}$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y}$$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\begin{aligned}\hat{\boldsymbol{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} \\ &= \mathbf{X}'\mathbf{Y}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\begin{aligned}\hat{\boldsymbol{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} \\ &= \mathbf{X}'\mathbf{Y} \\ \boldsymbol{\beta}^{\text{ridge}} &= (\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\begin{aligned}\hat{\boldsymbol{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} \\ &= \mathbf{X}'\mathbf{Y} \\ \boldsymbol{\beta}^{\text{ridge}} &= (\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y} \\ &= (\mathbf{I}_J + \lambda \mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\begin{aligned}\hat{\boldsymbol{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} \\ &= \mathbf{X}'\mathbf{Y} \\ \boldsymbol{\beta}^{\text{ridge}} &= (\mathbf{X}'\mathbf{X} + \lambda\mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y} \\ &= (\mathbf{I}_J + \lambda\mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y} \\ &= (\mathbf{I}_J + \lambda\mathbf{I}_J)^{-1} \hat{\boldsymbol{\beta}}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (1)

Suppose $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$.

$$\begin{aligned}\hat{\boldsymbol{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} \\ &= \mathbf{X}'\mathbf{Y} \\ \boldsymbol{\beta}^{\text{ridge}} &= (\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y} \\ &= (\mathbf{I}_J + \lambda \mathbf{I}_J)^{-1} \mathbf{X}'\mathbf{Y} \\ &= (\mathbf{I}_J + \lambda \mathbf{I}_J)^{-1} \hat{\boldsymbol{\beta}} \\ \beta_j^{\text{Ridge}} &= \frac{\hat{\beta}_j}{1 + \lambda}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\beta_j \sim \text{Normal}(0, \tau^2)$$

$$y_i \sim \text{Normal}(\beta_0 + \mathbf{x}_i' \boldsymbol{\beta}, \sigma^2)$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\beta_j \sim \text{Normal}(0, \tau^2)$$

$$y_i \sim \text{Normal}(\beta_0 + \mathbf{x}_i' \boldsymbol{\beta}, \sigma^2)$$

$$p(\boldsymbol{\beta} | \mathbf{X}, \mathbf{Y}) \propto \prod_{j=1}^J p(\beta_j) \prod_{i=1}^N p(y_i | \mathbf{x}_i, \boldsymbol{\beta})$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\beta_j \sim \text{Normal}(0, \tau^2)$$

$$y_i \sim \text{Normal}(\beta_0 + \mathbf{x}_i' \boldsymbol{\beta}, \sigma^2)$$

$$\begin{aligned} p(\boldsymbol{\beta} | \mathbf{X}, \mathbf{Y}) &\propto \prod_{j=1}^J p(\beta_j) \prod_{i=1}^N p(y_i | \mathbf{x}_i, \boldsymbol{\beta}) \\ &\propto \prod_{j=1}^J \frac{1}{\sqrt{2\pi\tau}} \exp\left(-\frac{\beta_j^2}{2\tau^2}\right) \prod_{i=1}^N \frac{1}{\sqrt{2\pi\sigma}} \exp\left(-\frac{(y_i - \beta_0 - \mathbf{x}_i' \boldsymbol{\beta})^2}{2\sigma^2}\right) \end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) = -\sum_{j=1}^J \frac{\beta_j^2}{2\tau^2} - \sum_{i=1}^N \frac{(y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2}{2\sigma^2}$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\begin{aligned}\log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= -\sum_{j=1}^J \frac{\beta_j^2}{2\tau^2} - \sum_{i=1}^N \frac{(y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2}{2\sigma^2} \\ -2\sigma^2 \log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N (y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2 + \sum_{j=1}^J \frac{\sigma^2}{\tau^2} \beta_j^2\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (2)

$$\begin{aligned}\log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= -\sum_{j=1}^J \frac{\beta_j^2}{2\tau^2} - \sum_{i=1}^N \frac{(y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2}{2\sigma^2} \\ -2\sigma^2 \log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N (y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2 + \sum_{j=1}^J \frac{\sigma^2}{\tau^2} \beta_j^2\end{aligned}$$

where:

Ridge Regression \rightsquigarrow Intuition (2)

$$\begin{aligned}\log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= -\sum_{j=1}^J \frac{\beta_j^2}{2\tau^2} - \sum_{i=1}^N \frac{(y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2}{2\sigma^2} \\ -2\sigma^2 \log p(\boldsymbol{\beta}|\mathbf{X}, \mathbf{Y}) &= \sum_{i=1}^N (y_i - \beta_0 - \mathbf{x}'\boldsymbol{\beta})^2 + \sum_{j=1}^J \frac{\sigma^2}{\tau^2} \beta_j^2\end{aligned}$$

where:

$$- \lambda = \frac{\sigma^2}{\tau^2}$$

Ridge Regression \rightsquigarrow Intuition (3)

Definition

Suppose \mathbf{X} is an $N \times J$ matrix. Then \mathbf{X} can be written as:

$$\mathbf{X} = \underbrace{\mathbf{U}}_{N \times N} \underbrace{\mathbf{S}}_{N \times J} \underbrace{\mathbf{V}'}_{J \times J}$$

Where:

$$\begin{aligned}\mathbf{U}'\mathbf{U} &= \mathbf{I}_N \\ \mathbf{V}'\mathbf{V} &= \mathbf{V}\mathbf{V}' = \mathbf{I}_J\end{aligned}$$

\mathbf{S} contains $\min(N, J)$ singular values, $\sqrt{\lambda_j} \geq 0$ down the diagonal and then 0's for the remaining entries

Ridge Regression \rightsquigarrow Intuition (3)

Recall: PCA:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \underbrace{\mathbf{W}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{W}'}_{\text{eigenvectors}}$$

Ridge Regression \rightsquigarrow Intuition (3)

Recall: PCA:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \underbrace{\mathbf{W}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{W}'}_{\text{eigenvectors}}$$

Using SVD:

Ridge Regression \rightsquigarrow Intuition (3)

Recall: PCA:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \underbrace{\mathbf{W}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{W}'}_{\text{eigenvectors}}$$

Using SVD:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \mathbf{V} \mathbf{S}' \underbrace{(\mathbf{U}' \mathbf{U})}_{\mathbf{I}_J} \mathbf{S} \mathbf{V}'$$

Ridge Regression \rightsquigarrow Intuition (3)

Recall: PCA:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \underbrace{\mathbf{W}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{W}'}_{\text{eigenvectors}}$$

Using SVD:

$$\begin{aligned} \frac{1}{N} \mathbf{X}' \mathbf{X} &= \mathbf{V} \mathbf{S}' \underbrace{(\mathbf{U}' \mathbf{U})}_{\mathbf{I}_J} \mathbf{S} \mathbf{V}' \\ &= \frac{1}{N} \mathbf{V} \mathbf{S}' \mathbf{S} \mathbf{V}' \end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (3)

Recall: PCA:

$$\frac{1}{N} \mathbf{X}' \mathbf{X} = \underbrace{\mathbf{W}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{W}'}_{\text{eigenvectors}}$$

Using SVD:

$$\begin{aligned} \frac{1}{N} \mathbf{X}' \mathbf{X} &= \mathbf{V} \mathbf{S}' \underbrace{(\mathbf{U}' \mathbf{U})}_{\mathbf{I}_J} \mathbf{S} \mathbf{V}' \\ &= \frac{1}{N} \mathbf{V} \mathbf{S}' \mathbf{S} \mathbf{V}' \\ &= \underbrace{\mathbf{V}}_{\text{eigenvectors}} \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_J \end{pmatrix} \underbrace{\mathbf{V}'}_{\text{eigenvectors}} \end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (3)

We can write the predicted values for a regular regression as

$$\begin{aligned}\hat{Y} &= \mathbf{X}\hat{\beta} \\ &= \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}'\mathbf{Y} = \sum_{j=1}^J u_j u_j' \mathbf{Y}\end{aligned}$$

Ridge Regression \rightsquigarrow Intuition (3)

We can write the predicted values for a regular regression as

$$\begin{aligned}\hat{Y} &= \mathbf{X}\hat{\boldsymbol{\beta}} \\ &= \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}'\mathbf{Y} = \sum_{j=1}^J u_j u_j' \mathbf{Y}\end{aligned}$$

We can write $\boldsymbol{\beta}^{\text{ridge}}$ as

Ridge Regression \rightsquigarrow Intuition (3)

We can write the predicted values for a regular regression as

$$\begin{aligned}\hat{Y} &= \mathbf{X}\hat{\beta} \\ &= \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}'\mathbf{Y} = \sum_{j=1}^J u_j u_j' \mathbf{Y}\end{aligned}$$

We can write β^{ridge} as

$$\hat{Y}^{\text{ridge}} = \mathbf{X}(\mathbf{X}'\mathbf{X} + \lambda\mathbf{I}_J)^{-1}\mathbf{X}'\mathbf{Y}$$

Ridge Regression \rightsquigarrow Intuition (3)

We can write the predicted values for a regular regression as

$$\begin{aligned}\hat{Y} &= \mathbf{X}\hat{\beta} \\ &= \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}'\mathbf{Y} = \sum_{j=1}^J u_j u_j' \mathbf{Y}\end{aligned}$$

We can write β^{ridge} as

$$\begin{aligned}\hat{Y}^{\text{ridge}} &= \mathbf{X}(\mathbf{X}'\mathbf{X} + \lambda\mathbf{I}_J)^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\tilde{\mathbf{S}}\mathbf{U}'\mathbf{Y}\end{aligned}$$

Where

$$\tilde{\mathbf{S}} = \left[\mathbf{S}(\mathbf{S}'\mathbf{S} + \lambda\mathbf{I}_J)^{-1}\mathbf{S} \right]$$

Ridge Regression \rightsquigarrow Intuition (3)

We can write the predicted values for a regular regression as

$$\begin{aligned}\hat{Y} &= \mathbf{X}\hat{\beta} \\ &= \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}'\mathbf{Y} = \sum_{j=1}^J \mathbf{u}_j \mathbf{u}_j' \mathbf{Y}\end{aligned}$$

We can write β^{ridge} as

$$\begin{aligned}\hat{Y}^{\text{ridge}} &= \mathbf{X}(\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J)^{-1}\mathbf{X}'\mathbf{Y} \\ &= \mathbf{U}\tilde{\mathbf{S}}\mathbf{U}'\mathbf{Y}\end{aligned}$$

Where

$$\tilde{\mathbf{S}} = \left[\mathbf{S}(\mathbf{S}'\mathbf{S} + \lambda \mathbf{I}_J)^{-1}\mathbf{S} \right]$$

Which we can write as:

$$\hat{Y}^{\text{ridge}} = \sum_{j=1}^J \mathbf{u}_j \frac{\lambda_j}{\lambda_j + \lambda} \mathbf{u}_j' \mathbf{Y}$$

Degrees of Freedom for Ridge

We will say that the degrees of freedom for Ridge regression with penalty λ is

$$\text{dof}(\lambda) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \lambda}$$

Lasso Regression Objective Function

Different Penalty for Model Complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \underbrace{|\beta_j|}_{\text{Penalty}}$$

Lasso Regression Objective Function

Different Penalty for Model Complexity

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \underbrace{|\beta_j|}_{\text{Penalty}}$$

Lasso Regression Optimization

Definition

Coordinate Descent Algorithms:

Consider $g : \mathbb{R}^J \rightarrow \mathbb{R}$. Our goal is to find $\mathbf{x}^ \in \mathbb{R}^J$ such that $g(\mathbf{x}^*) \leq g(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}$.*

To find \mathbf{x}^ :*

Until convergence: for each iteration t and each coordinate j

$$x_j^{t+1} = \arg \min_{x_j \in \mathbb{R}} g(x_1^{t+1}, x_2^{t+1}, \dots, x_{j-1}^{t+1}, x_j, x_{j+1}^t, \dots, x_J^t)$$

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$
- **Case 2:** If $\beta_j > (<) 0 \rightsquigarrow$ differentiable \rightsquigarrow differentiate and solve for β_j

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$
- **Case 2:** If $\beta_j > (<)0 \rightsquigarrow$ differentiable \rightsquigarrow differentiate and solve for β_j

Define $\tilde{y}_i^j = \beta_0 + \sum_{l \neq j} x_{il} \beta_l$

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$
- **Case 2:** If $\beta_j > (<) 0 \rightsquigarrow$ differentiable \rightsquigarrow differentiate and solve for β_j

Define $\tilde{y}_i^j = \beta_0 + \sum_{l \neq j} x_{il} \beta_l$

$$r^j \equiv \frac{1}{N} \sum_{i=1}^N x_{ij} (y_i - \tilde{y}_i^j)$$

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$
- **Case 2:** If $\beta_j > (<) 0 \rightsquigarrow$ differentiable \rightsquigarrow differentiate and solve for β_j

Define $\tilde{y}_i^j = \beta_0 + \sum_{l \neq j} x_{il} \beta_l$

$$r^j \equiv \frac{1}{N} \sum_{i=1}^N x_{ij} (y_i - \tilde{y}_i^j)$$

Update step for β_j is

Lasso Regression Optimization: Coordinate Descent

$$\tilde{f}(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{2N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J |\beta_j|$$

- **Case 1:** If $\beta_j = 0 \rightsquigarrow$ not differentiable. But $\beta_j = 0$
- **Case 2:** If $\beta_j > (<)0 \rightsquigarrow$ differentiable \rightsquigarrow differentiate and solve for β_j

Define $\tilde{y}_i^j = \beta_0 + \sum_{l \neq j} x_{il} \beta_l$

$$r^j \equiv \frac{1}{N} \sum_{i=1}^N x_{ij} (y_i - \tilde{y}_i^j)$$

Update step for β_j is

$$\beta_j \leftarrow \text{sign}(r^j) \max(|r^j| - \lambda, 0)$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j|$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j| \\ &= -2\mathbf{X}'\mathbf{Y}\boldsymbol{\beta} + \boldsymbol{\beta}'\boldsymbol{\beta} + \lambda \sum_{j=1}^J |\beta_j| \end{aligned}$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j| \\ &= -2\mathbf{X}'\mathbf{Y}\boldsymbol{\beta} + \boldsymbol{\beta}'\boldsymbol{\beta} + \lambda \sum_{j=1}^J |\beta_j| \end{aligned}$$

The coefficient is

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j| \\ &= -2\mathbf{X}'\mathbf{Y}\boldsymbol{\beta} + \boldsymbol{\beta}'\boldsymbol{\beta} + \lambda \sum_{j=1}^J |\beta_j| \end{aligned}$$

The coefficient is

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$\begin{aligned}f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j| \\&= -2\mathbf{X}'\mathbf{Y}\boldsymbol{\beta} + \boldsymbol{\beta}'\boldsymbol{\beta} + \lambda \sum_{j=1}^J |\beta_j|\end{aligned}$$

The coefficient is

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

- $\text{sign}(\cdot) \rightsquigarrow 1$ or -1

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Suppose again $\mathbf{X}'\mathbf{X} = \mathbf{I}_J$

$$\begin{aligned} f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) &= (\mathbf{Y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{Y} - \mathbf{X}\boldsymbol{\beta}) + \lambda \sum_{j=1}^J |\beta_j| \\ &= -2\mathbf{X}'\mathbf{Y}\boldsymbol{\beta} + \boldsymbol{\beta}'\boldsymbol{\beta} + \lambda \sum_{j=1}^J |\beta_j| \end{aligned}$$

The coefficient is

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

- $\text{sign}(\cdot) \rightsquigarrow 1$ or -1
- $\left(|\hat{\beta}_j| - \lambda \right)_+ = \max(|\hat{\beta}_j| - \lambda, 0)$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

With hard assignment, selecting M biggest components

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

With hard assignment, selecting M biggest components

$$\beta_j^{\text{subset}} = \hat{\beta}_j \cdot I \left(|\hat{\beta}_j| \geq |\hat{\beta}_{(M)}| \right)$$

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

With hard assignment, selecting M biggest components

$$\beta_j^{\text{subset}} = \hat{\beta}_j \cdot I\left(|\hat{\beta}_j| \geq |\hat{\beta}_{(M)}|\right)$$

Intuition 2: Prior on coefficients \rightsquigarrow Laplace “The Bayesian LASSO”

Lasso Regression \rightsquigarrow Intuition 1, Soft Thresholding

Compare soft assignment

$$\beta_j^{\text{LASSO}} = \text{sign}(\hat{\beta}_j) \left(|\hat{\beta}_j| - \lambda \right)_+$$

With hard assignment, selecting M biggest components


$$\beta_j^{\text{subset}} = \hat{\beta}_j \cdot I\left(|\hat{\beta}_j| \geq |\hat{\beta}_{(M)}|\right)$$

Intuition 2: Prior on coefficients \rightsquigarrow Laplace “The Bayesian LASSO”

Why does LASSO induce sparsity?


Comparing Ridge and LASSO

Ridge Regression



Comparing Ridge and LASSO

LASSO Regression



Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

$$\sum_{j=1}^2 \beta_j^2 = \frac{1}{2} + \frac{1}{2} = 1$$

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

$$\sum_{j=1}^2 \beta_j^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\sum_{j=1}^2 \tilde{\beta}_j^2 = 1 + 0 = 1$$

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

$$\sum_{j=1}^2 \beta_j^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\sum_{j=1}^2 \tilde{\beta}_j^2 = 1 + 0 = 1$$

Under LASSO

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

$$\sum_{j=1}^2 \beta_j^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\sum_{j=1}^2 \tilde{\beta}_j^2 = 1 + 0 = 1$$

Under LASSO

$$\sum_{j=1}^2 |\beta_j| = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

Comparing Ridge and LASSO

Contrast $\beta = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $\tilde{\beta} = (1, 0)$

Under ridge:

$$\sum_{j=1}^2 \beta_j^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\sum_{j=1}^2 \tilde{\beta}_j^2 = 1 + 0 = 1$$

Under LASSO

$$\sum_{j=1}^2 |\beta_j| = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

$$\sum_{j=1}^2 |\tilde{\beta}_j| = 1 + 0 = 1$$

Ridge and LASSO: The Elastic-Net

Combining the two criteria \rightsquigarrow Elastic-Net

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \left(\frac{1}{2} (1 - \alpha) \beta_j^2 + \alpha |\beta_j| \right)$$

Ridge and LASSO: The Elastic-Net

Combining the two criteria \rightsquigarrow Elastic-Net

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \left(\frac{1}{2} (1 - \alpha) \beta_j^2 + \alpha |\beta_j| \right)$$

The new update step (for coordinate descent:)

Ridge and LASSO: The Elastic-Net

Combining the two criteria \rightsquigarrow Elastic-Net

$$f(\boldsymbol{\beta}, \mathbf{X}, \mathbf{Y}) = \frac{1}{N} \sum_{i=1}^N \left(y_i - \beta_0 - \sum_{j=1}^J \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^J \left(\frac{1}{2} (1 - \alpha) \beta_j^2 + \alpha |\beta_j| \right)$$

The new update step (for coordinate descent:)

$$\beta_j \leftarrow \frac{\text{sign}(r^j) \max(|r^j| - \lambda \alpha, 0)}{1 + \lambda(1 - \alpha)}$$

Selecting λ

How do we determine λ ? \rightsquigarrow Cross validation (Recall code from Monday)

Selecting λ

How do we determine λ ? \rightsquigarrow Cross validation (Recall code from Monday)
Applying models gives score (probability) of document belong to class \rightsquigarrow
threshold to classify

Selecting λ

How do we determine λ ? \rightsquigarrow Cross validation (Recall code from Monday)
Applying models gives score (probability) of document belong to class \rightsquigarrow
threshold to classify

Credit Claiming (Grimmer, Westwood, and Messing 2014)

```
library(glmnet)
set.seed(8675309) ##setting seed
folds<- sample(1:10, nrow(dtm), replace=T) ##assigning to fold
out_of_samp<- c() ##collecting the predictions
```

Credit Claiming (Grimmer, Westwood, and Messing 2014)

```
for(z in 1:10){  
  train<- which(folds!=z) ##the observations we will use to train the model  
  
  test<- which(folds==z) ##the observations we will use to test the model  
  part1<- cv.glmnet(x = dtm[train,], y = credit[train], alpha = 1, family =  
  binomial) ##fitting the LASSO model on the data.  
  ## alpha = 1 -> LASSO  
  ## alpha = 0 -> RIDGE  
  ## 0<alpha<1 -> Elastic-Net  
  out_of_samp[test]<- predict(part1, newx= dtm[test,], s = part1$lambda.min,  
  type = "class") ##predicting the labels  
  print(z) ##printing the labels  
}  
  
conf_table<- table(out_of_samp, credit) ##calculating the confusion table  
> round(sum(diag(conf_table))/len(credit), 3)  
[1] 0.844
```

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

$$\beta^{\text{Ridge}} = \left(\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}'\mathbf{Y}$$

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

$$\begin{aligned}\beta^{\text{Ridge}} &= \left(\mathbf{X}'\mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}'\mathbf{Y} \\ \hat{\mathbf{Y}} &= \mathbf{X}(\beta)^{\text{Ridge}}\end{aligned}$$

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

$$\begin{aligned}\beta^{\text{Ridge}} &= \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y} \\ \hat{\mathbf{Y}} &= \mathbf{X} (\beta)^{\text{Ridge}} \\ &= \underbrace{\mathbf{X} \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y}}_{\text{Hat Matrix}}\end{aligned}$$

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

$$\begin{aligned}\beta^{\text{Ridge}} &= \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y} \\ \hat{\mathbf{Y}} &= \mathbf{X} (\beta)^{\text{Ridge}} \\ &= \underbrace{\mathbf{X} \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y}}_{\text{Hat Matrix}} \\ \hat{\mathbf{Y}} &= \underbrace{\mathbf{H}}_{\text{Smoother Matrix}} \mathbf{Y}\end{aligned}$$

Generalized Cross Validation and Ridge Regression

In some special cases there are analytic solutions:

$$\begin{aligned}\beta^{\text{Ridge}} &= \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y} \\ \hat{\mathbf{Y}} &= \mathbf{X} (\beta)^{\text{Ridge}} \\ &= \underbrace{\mathbf{X} \left(\mathbf{X}' \mathbf{X} + \lambda \mathbf{I}_J \right)^{-1} \mathbf{X}' \mathbf{Y}}_{\text{Hat Matrix}} \\ \hat{\mathbf{Y}} &= \underbrace{\mathbf{H}}_{\text{Smoother Matrix}} \mathbf{Y}\end{aligned}$$

Generalized Cross Validation and Ridge Regression

Why do we care?

Generalized Cross Validation and Ridge Regression

Why do we care?

Leave one out cross validation

Generalized Cross Validation and Ridge Regression

Why do we care?

Leave one out cross validation

$$\text{Cross Validation(1)} = \frac{1}{N} \sum_{i=1}^N (Y_i - f(\mathbf{X}_{-i}, \mathbf{Y}_{-i}, \lambda, \hat{\beta}))^2$$

Generalized Cross Validation and Ridge Regression

Why do we care?

Leave one out cross validation

$$\begin{aligned}\text{Cross Validation(1)} &= \frac{1}{N} \sum_{i=1}^N (Y_i - f(\mathbf{X}_{-i}, \mathbf{Y}_{-i}, \lambda, \hat{\boldsymbol{\beta}}))^2 \\ &= \frac{1}{N} \sum_{i=1}^N \left(\frac{Y_i - f(\mathbf{X}, \mathbf{Y}, \lambda, \hat{\boldsymbol{\beta}})}{1 - H_{ii}} \right)^2\end{aligned}$$

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters}$ (class regression = number of independent variables + 1)

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters}$ (class regression = number of independent variables + 1)
- For Ridge regression:

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters (class regression = number of independent variables + 1)}$
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters}$ (class regression = number of independent variables + 1)
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\Sigma = \mathbf{X}'\mathbf{X}$

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters}$ (class regression = number of independent variables + 1)
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\mathbf{\Sigma} = \mathbf{X}'\mathbf{X}$ (!!!!!!)

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters (class regression = number of independent variables + 1)}$
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\Sigma = \mathbf{X}'\mathbf{X}$ (!!!!)

Define generalized cross validation:

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters (class regression = number of independent variables + 1)}$
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\mathbf{\Sigma} = \mathbf{X}'\mathbf{X}$ (!!!!)

Define generalized cross validation:

$$\text{GCV} = \frac{1}{N} \sum_{i=1}^N \left(\frac{Y_i - \hat{Y}_i}{1 - \frac{\text{Tr}(\mathbf{H})}{N}} \right)^2$$

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters (class regression = number of independent variables + 1)}$
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\Sigma = \mathbf{X}'\mathbf{X}$ (!!!!)

Define generalized cross validation:

$$\text{GCV} = \frac{1}{N} \sum_{i=1}^N \left(\frac{Y_i - \hat{Y}_i}{1 - \frac{\text{Tr}(\mathbf{H})}{N}} \right)^2$$

Applicable in any setting where we can write **Smoother** matrix

Generalized Cross Validation and Ridge Regression

Calculating \mathbf{H} can be computationally expensive

- $\text{Trace}(\mathbf{H}) \equiv \text{Tr}(\mathbf{H}) = \sum_{i=1}^N H_{ii}$
- $\text{Tr}(\mathbf{H}) = \text{Effective number of parameters (class regression = number of independent variables + 1)}$
- For Ridge regression:

$$\text{Tr}(\mathbf{H}) = \sum_{j=1}^J \frac{\lambda_j}{\lambda_j + \underbrace{\lambda}_{\text{Penalty}}}$$

where λ_j is the j^{th} Eigenvalue from $\Sigma = \mathbf{X}'\mathbf{X}$ (!!!!)

Define generalized cross validation:

$$\text{GCV} = \frac{1}{N} \sum_{i=1}^N \left(\frac{Y_i - \hat{Y}_i}{1 - \frac{\text{Tr}(\mathbf{H})}{N}} \right)^2$$

Applicable in any setting where we can write **Smoother** matrix