

Classical Mechanics 1, Autumn 2021 CMI
Problem set 9
- Govind S. Krishnaswami

Soham Chatterjee

Roll: BMC202175

1. The frame S' that coincides with frame S at $t = t' = 0$ and moves uniformly at speed v along the positive x axis of S . Hence

$$x' = \gamma(x - vt) \quad \text{and} \quad t' = \gamma\left(t - \frac{vx}{c^2}\right)$$

Hence

$$x'_2 - x'_1 = \gamma((x_2 - x_1) - v(t_2 - t_1)) \quad \text{and} \quad t'_2 - t'_1 = \gamma\left((t_2 - t_1) - \frac{v(x_2 - x_1)}{c^2}\right)$$

Let's denote $\Delta_x = x_2 - x_1$ and $\Delta_t = t_2 - t_1$. Hence given that $c^2\Delta_t^2 - \Delta_x^2 = \Delta$ Now

$$\begin{aligned} c^2(t'_2 - t'_1)^2 - (x'_2 - x'_1)^2 &= c^2\gamma^2\left(\Delta_t - \frac{v\Delta_x}{c^2}\right)^2 - \gamma^2(\Delta_x - v\Delta_t)^2 \\ &= \gamma^2\left[\left(c\Delta_t - \frac{v\Delta_x}{c}\right)^2 - (\Delta_x - v\Delta_t)^2\right] \\ &= \gamma^2\left[\left(c^2\Delta_t^2 + \frac{v^2\Delta_x^2}{c^2} - 2v\Delta_t\Delta_x\right) - (\Delta_x^2 + v^2\Delta_t^2 - 2v\Delta_x\Delta_t)\right] \\ &= \gamma^2\left[\Delta_t^2(c^2 - v^2) - \Delta_x^2\left(\frac{v^2}{c^2} - 1\right)\right] \\ &= c^2\Delta_t^2 - \Delta_x^2 = \Delta \end{aligned}$$

Hence the square of space-time separation is unchanged under Lorentz Transformation.

2. (a)

$$\begin{aligned} E^2 - \mathbf{p}^2c^2 &= \gamma^2m^2c^4 - \gamma^2m^2\mathbf{v}^2c^2 \\ &= \gamma^2m^2c^2(c^2 - \mathbf{v}^2) \\ &= m^2c^4 = (mc^2)^2 \end{aligned}$$

- (b)

$$\gamma = \left(1 - \frac{\mathbf{v}^2}{c^2}\right)^{\frac{1}{2}} \approx 1 + \frac{1}{2} \frac{\mathbf{v}^2}{c^2}$$

Here we are ignoring the higher order terms of $\frac{\mathbf{v}^2}{c^2}$ for non-relativistic limit. Hence

$$\begin{aligned} E &= \gamma mc^2 \approx mc^2 \left(1 + \frac{\mathbf{v}^2}{2c^2}\right) = mc^2 + \frac{m\mathbf{v}^2}{2} \\ \mathbf{p} &= \gamma m\mathbf{v} \approx m\mathbf{v} \left(1 + \frac{\mathbf{v}^2}{2c^2}\right) = m\mathbf{v} + \frac{m\mathbf{v}^3}{2c^2} \end{aligned}$$

(c) According to the dispersion relation $E^2 - \mathbf{p}^2 c^2 = (mc^2)^2 \implies E^2 = (mc^2)^2 + \mathbf{p}^2 c^2$. Hence

$$\begin{aligned}
 E &= ((mc^2)^2 + \mathbf{p}^2 c^2)^{\frac{1}{2}} \\
 &= mc^2 \left(1 + \frac{\mathbf{p}^2 c^2}{m^2 c^4} \right)^{\frac{1}{2}} \\
 &= mc^2 \left(1 + \frac{\mathbf{p}^2}{m^2 c^2} \right)^{\frac{1}{2}} \\
 &\approx mc^2 \left(1 + \frac{1}{2} \frac{\mathbf{p}^2}{m^2 c^2} \right) \\
 &= mc^2 + \frac{\mathbf{p}^2}{2m}
 \end{aligned}$$

3. (a) In the graph we have taken E in the x -axis, p_1 in the y -axis and p_2 in the z -axis.




Figure: Surface of $E - p_1 - p_2$ in \mathbb{R}^3

Here the x -axis intersected the surfaces in two points. It intersected in positive side at $(mc^2, 0, 0)$ and in negative side at $(-mc^2, 0, 0)$

- (b) Name of the surface can be "Relativistic Dispersion Surface"
- (c) Since $E = \gamma mc^2$ we can say $E > 0$. Hence the surface which is in the negative side of the x -axis is not physically relevant. Therefore the surface in the positive direction of x -axis is physically relevant.

4. (a) I select the chapters
 - Collisions or scattering and conservation laws
 - Special theory of relativity
- (b) Let we have two identical rods l_1, l_2 of length l and mass m (The rods have uniform mass distribution. Hence the centre of the rods are centre of mass). Suppose l_1 is in rest along the x axis whose left end is at the $(\frac{l}{2}, 0)$. l_2 parallel to x -axis and it is approaching towards the origin from the positive side of y -axis with velocity v whose centre is on the y -axis such that it collides with l_1 at the end point.
 - After the collision the rods will start rotating about their centres. As l_2 's right end collided with l_1 's left end both l_1 and l_2 will start rotating counter-clockwise if we observe from the positive direction of z -axis. See here before the collision their was no angular momentum . But after the collision they have angular momentum which does not cancel out each other instead add up. Hence angular momentum conservation rule didn't follow here. But as per the law in absence of torque angular momentum is supposed to be conserved. What exactly wrong here ?
 - If we know m, l, v then the whole situation after the collision is predetermined. How to calculate the final outcome in general condition ?
- (c) Suppose S is an inertial frame. S' is a frame which moving at the speed of light c relative to S . Then γ becomes infinite. Hence with respect to S any time interval in S becomes 0 in S' . Hence everything in S happens instantaneously in S' . In that case any time interval in S' makes no sense. Then how can we apply Lorentz Transformation here where in one frame time interval does not make sense and in another frame time interval makes sense. If time interval does not make sense in S' how can we define time in S' ?