

Analytic Two-Allele Model

Liu et al.

February 7, 2019

1 Introduction

To understand the roles that mutation probability and growth rate during treatment play in determining clinical prevalence of specific resistance mutations, we developed a model for two potential resistance variants.

Consider a population of pathogenic cells which expand prior to treatment and collapse during treatment. During the lifetime of this population of cells, the initial sensitive population has the potential to seed resistant subclones which have a positive net growth rate in the presence of drug. The population dynamics of a resistant clone depend on allele-specific parameters, in particular its mutational probability (i.e. the likelihood that the clone will spawn) and its degree of resistance (i.e. the growth rate during treatment).

We are interested in identifying the conditions for which mutation bias plays a significant role in determining prevalence at the scale of patient populations. To address this, we consider the simplest case of two distinct mutations that confer drug resistance (Allele A and Allele B). To simplify the analysis, we do not consider the potential for compound mutations.

1.1 Resistance and Mutational Probability Profiles

It is interesting to consider the case where there exists a trade-off between resistance and mutational probability. We assign Allele A a high degree of resistance and low mutational probability; we assign the opposite profile for Allele B. These profiles are summarized in **Supplementary Table 1**.

Supplementary Table 1

	Allele A	Allele B
Resistance	high	low
Mutational Probability	low	high

We then ask the question: what will the dominant allele be upon relapse? In other words, which is the first clone to reach clinical detection? In identifying conditions where Allele B (the allele with a greater mutational probability, but a lower degree of resistance) is more likely to be the dominant allele, we find

parameter sets where we expect mutational probability to play a significant role in shaping distributions of specific amino acid mutations in clinical populations.

2 General Outcomes

In identifying which will be the dominant allele upon relapse in a two-allele system, consider the four possible outcomes:

Supplementary Table 2

		Allele A present?	
		No	Yes
Allele B present?	No	Outcome 1	Outcome 2
	Yes	Outcome 3	Outcome 4

Here, we say an allele is present if its mutational event occurs during the lifetime of the sensitive population and the newly spawned clone escapes stochastic eradication (i.e. it eventually reaches clinical detection). Thus, the results of each outcome are summarized as follows.

Outcome 1: Neither allele is present.

Outcome 2: Allele A is fixed.

Outcome 3: Allele B is fixed.

Outcome 4: Both alleles are present.

Outcome 4 can be further divided into two suboutcomes:

Outcome 4.A: Allele A reaches clinical detection first.

Outcome 4.B: Allele B reaches clinical detection first.

Towards addressing our initial question, we see that:

$$\begin{aligned}
P[\text{Allele A is dominant upon relapse}] &= \\
&P[\text{Outcome 2}] + P[\text{Outcome 4.A}|\text{Outcome 4}] \cdot P[\text{Outcome 4}] \\
P[\text{Allele B is dominant upon relapse}] &= \\
&P[\text{Outcome 3}] + P[\text{Outcome 4.B}|\text{Outcome 4}] \cdot P[\text{Outcome 4}]
\end{aligned} \tag{1}$$

In the following analyses, we will derive these probabilities.

3 Probability of Mutation

Let us begin by considering the probabilities of each outcome. Because the presence of each allele is independent of the other, it follows that:

$$\begin{aligned}
P[\mathbf{Outcome\ 1}] &= P[\text{Allele A is absent}] \cdot P[\text{Allele B is absent}] \\
P[\mathbf{Outcome\ 2}] &= P[\text{Allele A is present}] \cdot P[\text{Allele B is absent}] \\
P[\mathbf{Outcome\ 3}] &= P[\text{Allele A is absent}] \cdot P[\text{Allele B is present}] \\
P[\mathbf{Outcome\ 4}] &= P[\text{Allele A is present}] \cdot P[\text{Allele B is present}]
\end{aligned} \tag{2}$$

To find the probability that an allele is absent, let $N_i(t)$ be the number of mutational events generating the i th allele that occur on the time interval $(0, t]$. The discrete random variable $N_i(t)$ can be modeled as a nonhomogenous Poisson process. Thus, the probability that k mutations occur on the interval $(0, t]$ is given by:

$$P[N_i(t) = k] = \frac{[\int_0^t \lambda_i(y) dy]^k}{k!} \exp\left(-\int_0^t \lambda(y) dy\right) \tag{3}$$

where the mutational hazard rate $\lambda_i(t)$ is the probability of a mutation to the i th allele occurring on the infinitesimal interval $(t, t + dt)$. This hazard rate is a function of the size of the sensitive population $x_S(t)$ from which the resistant variant arises, and is given by:

$$\lambda_i(t) = x_S(t) b_S \mu \rho_i \frac{r_i}{b_i} \tag{4}$$

Here, b_S is the division rate of the sensitive population; μ is the probability that a resistance mutation occurs per division event of the sensitive population; ρ_i is the probability of mutation to the i th allele given that a resistance mutation occurs. The parameter ρ_i is used to describe differences in allele-specific mutational probabilities. Note that $\sum_i \rho_i = 1$. We multiply by the allele-specific survival probability r_i/b_i to consider only mutation events where the newly spawned clone escapes stochastic eradication.

The size of the sensitive population can be modeled assuming exponential growth prior to treatment and exponential decay during treatment. If treatment starts at time T , then the number of sensitive cells $x_S(t)$ is:

$$x_S(t) = \begin{cases} \frac{b_S}{r_S} e^{r_S t} & 0 \leq t < T \\ M e^{r'_S (t-T)} & T \leq t \end{cases} \tag{5}$$

Here, r_S is the net growth rate of sensitive cells prior to treatment, r'_S is their net growth rate during treatment ($r'_S < 0$) and M is the size of the sensitive

population at the start of treatment (i.e. the detection size). For our analysis, we assume that drug treatment increases the death rate and leaves the birth rate unchanged, i.e. $b'_S = b_S$.

The time to detection T can be related to M by solving the equation $\frac{b_S}{r_S} e^{r_S T} = M$, yielding:

$$T = \frac{1}{r_S} \ln \left(\frac{Mr_S}{b_S} \right) \quad (6)$$

Substituting equations (4) and (5) into (3) and choosing $k = 0$ events, we arrive at an expression $S_i(t)$ for the probability that a mutation to the i th allele has not occurred by time t .

$$S_i(t) = P[N_i(t) = 0] = \begin{cases} \exp \left(-\frac{b_S^2 r_i \mu \rho_i}{r_S b_i} \int_0^t e^{r_S y} dy \right) & 0 \leq t < T \\ \exp \left(-\frac{b_S^2 r_i \mu \rho_i}{r_S b_i} \int_0^T e^{r_S y} dy - \frac{Mb_S r_i \mu \rho_i}{b'_i} e^{-r'_S T} \int_T^t e^{r'_S y} dy \right) & T \leq t \end{cases} \quad (7)$$


Let $\alpha_i = \frac{b_S^2 r_i \mu \rho_i}{r_S b_i}$ and $\beta_i = \frac{Mb_S r_i \mu \rho_i}{b'_i} e^{-r'_S T}$. Solving the integrals, we can simplify equation (7) to:

$$S_i(t) = \begin{cases} \exp \left(-\frac{\alpha_i}{r_S} (e^{r_S t} - 1) \right) & 0 \leq t < T \\ \exp \left(-\frac{\alpha_i}{r_S} (e^{r_S T} - 1) - \frac{\beta_i}{r'_S} (e^{r'_S t} - e^{r'_S T}) \right) & T \leq t \end{cases} \quad (8)$$

If we define $\eta_i = e^{\alpha_i/r_S}$ and $\nu_i = \eta_i \exp[\frac{-\alpha_i}{r_S} e^{r_S T} + \frac{\beta_i}{r'_S} e^{r'_S T}]$, $S_i(t)$ can be further simplified:

$$S_i(t) = \begin{cases} \eta_i \exp \left(\frac{-\alpha_i}{r_S} e^{r_S t} \right) & 0 \leq t < T \\ \nu_i \exp \left(\frac{-\beta_i}{r'_S} e^{r'_S t} \right) & T \leq t \end{cases} \quad (9)$$

The complement of $S_i(t)$, $F_i(t) = 1 - S_i(t)$, gives the cumulative mutational risk, i.e. the probability that at least one mutation *has* occurred by time t . An example trajectory of $F_i(t)$ is shown in **Supplementary Figure 1**.



Supplementary Figure 1: $F_i(t)$ for $b_S = b_i = 0.14$ /day, $r_S = 0.01$ /day, $r'_S = -0.03$, $r_i = 0.0075$ /day, $\mu = 10^{-8}$, $\rho_i = 0.9$, and $M = 10^9$.

Recalling that $r'_S < 0$, the probability that at least one mutation to the i th allele occurs during the lifetime of the tumor is:

$$F_i(\infty) = 1 - S_i(\infty) = 1 - \nu_i \quad (10)$$

This gives us expressions for the probabilities of the four outcomes outlined in **Supplementary Table 1**:

$$\begin{aligned} P[\text{Outcome 1}] &= S_A(\infty) \cdot S_B(\infty) = \nu_A \cdot \nu_B \\ P[\text{Outcome 2}] &= F_A(\infty) \cdot S_B(\infty) = (1 - \nu_A) \cdot \nu_B \\ P[\text{Outcome 3}] &= S_A(\infty) \cdot F_B(\infty) = \nu_A \cdot (1 - \nu_B) \\ P[\text{Outcome 4}] &= F_A(\infty) \cdot F_B(\infty) = (1 - \nu_A) \cdot (1 - \nu_B) \end{aligned} \quad (11)$$

4 Outcome 4: Competition

We now turn our attention to the **Outcome 4**, where both alleles are present. Since we are interested in which allele will be dominant at relapse, we ask the question: which clone will reach clinical detection first?

We see that

$$\begin{aligned} P[\text{Outcome 4.A}] &= P[T_{det,A} < T_{det,B}] \\ P[\text{Outcome 4.B}] &= P[T_{det,B} < T_{det,A}] \end{aligned} \quad (12)$$


where $T_{det,i}$ is the time to detection of the i th allele (i.e. the time for the i th allele to reach detection size M). We focus on finding an expression for the probability

of **Outcome 4.B**. Since, given **Outcome 4**, **Outcome 4.A** and **Outcome 4.B** are mutually exclusive and collectively exhaustive, the probability of **Outcome 4.A** can be found given $P[\mathbf{Outcome\ 4.A}|\mathbf{Outcome\ 4}] = 1 - P[\mathbf{Outcome\ 4.B}|\mathbf{Outcome\ 4}]$.

If $T_{mut,i}$ is the time to the first mutational event of the i th allele (given there is one sensitive cell at $t = 0$) and $T_{og,i}$ is the time to outgrowth for the newly spawned clone of the i th allele, then

$$T_{det,i} = T_{mut,i} + T_{og,i} \quad (13)$$

For Alleles A and B, we might expect a timeline that resembles:



We expect Allele A to grow out faster (i.e. $T_{og,A} < T_{og,B}$) because it is more resistant and thus has a faster growth rate in the presence of drug. However, this does not necessarily mean that it will reach clinical detection first. We see that if an Allele B clone spawns early enough, it will be the dominant allele upon relapse. This is true if $T_{og,B} - T_{og,A} < T_{mut,A} - T_{mut,B}$. Thus,

$$P[\mathbf{Outcome\ 4.B}] = P[T_{det,B} < T_{det,A}] = P[\Delta T_{og} < T_{mut,A} - T_{mut,B}] \quad (14)$$

where $\Delta T_{og} = T_{og,B} - T_{og,A}$.

Because we consider the variation in $T_{og,i}$ to be very small compared to $T_{mut,i}$, we treat ΔT_{og} as a deterministic variable. Its value can be calculated using resistance allele-specific variations of equation (6):

$$\Delta T_{og} = T_{og,B} - T_{og,A} = \frac{1}{r'_B} \ln\left(\frac{Mr'_B}{b_B}\right) - \frac{1}{r'_A} \ln\left(\frac{Mr'_A}{b_A}\right) \quad (15)$$

From here on, we abbreviate times to mutation $T_{mut,A}$ and $T_{mut,B}$ to T_A and T_B , respectively. We consider these to be random variables. To obtain a solution for the right-hand expression in equation (14), we find the probability

that the difference that these two random variables exceeds a threshold value, ΔT_{og} . The probability that two random variables will take on values on a two-dimensional interval is the integral of their joint probability density function over that interval.

$$P[\Delta T_{og} < T_A - T_B] = \iint_D f_{A,B}^*(t_A, t_B) dA \quad (16)$$

Here, $f_{A,B}^*(t_A, t_B)$ is the conditional joint probability density function for T_A and T_B , and D is the region in the first quadrant (where $f_{A,B}$ takes on nonzero values) bounded above by the line $T_B = T_A - \Delta T_{og}$. We consider the *conditional* joint probability density function because we are concerned with **Outcome 4**, where both alleles are present.

Because T_A and T_B are independent random variables, their joint conditional probability density function is given by the product of their individual PDFs:

$$f_{A,B}^*(t_A, t_B) = f_A^*(t_A) \cdot f_B^*(t_B) \quad (17)$$

The probability density function for T_i , $f_i(t_i)$, gives the probability that a mutational event occurs on the interval $(t_i, t_i + dt_i)$. It is given by the derivative of the cumulative density function $f_i(t_i) = \frac{d}{dt} F_i(t_i)$. Therefore,


$$f_i(t_i) = \frac{d}{dt} F_i(t_i) = \begin{cases} \eta_i \alpha_i e^{r_S t_i} \exp\left[\frac{-\alpha_i}{r_S} e^{r_S t_i}\right] & 0 \leq t < T \\ \nu_i \beta_i e^{r'_S t_i} \exp\left[\frac{-\beta_i}{r'_S} e^{r'_S t_i}\right] & T \leq t \end{cases} \quad (18)$$

To determine the *conditional* probability density function $f_i^*(t_i)$, we divide $f_i(t_i)$ by the probability that the i th allele *will* spawn. From equation (10), we have:

$$f_i^*(t_i) = \frac{f_i(t_i)}{F_i(\infty)} = \begin{cases} \frac{\eta_i \alpha_i}{1 - \nu_i} e^{r_S t_i} \exp\left[\frac{-\alpha_i}{r_S} e^{r_S t_i}\right] & 0 \leq t < T \\ \frac{\nu_i \beta_i}{1 - \nu_i} e^{r'_S t_i} \exp\left[\frac{-\beta_i}{r'_S} e^{r'_S t_i}\right] & T \leq t \end{cases} \quad (19)$$

which is the conditional probability density function of T_i given that at least one mutational event to the i th allele eventually takes place.

An example plot of $f_i^*(t_i)$ is shown in **Supplementary Figure 2**.



Supplementary Figure 2: Instantaneous mutational risk, conditioned on the mutational event eventually occurring. The same parameters were used as in **Supplementary Figure 1**.


From equations (17) and (19), we arrive at the formal conditional joint probability density function:

$$f_{A,B}^*(t_A, t_B) = \begin{cases} C_1 \cdot e^{r_S t_A} \exp\left(\frac{-\alpha_A}{r_S} e^{r_S t_A}\right) \cdot e^{r_S t_B} \exp\left(\frac{-\alpha_B}{r_S} e^{r_S t_B}\right) & 0 \leq t_A, t_B < T \\ C_2 \cdot e^{r_S t_A} \exp\left(\frac{-\alpha_A}{r_S} e^{r_S t_A}\right) \cdot e^{r'_S t_B} \exp\left(\frac{-\beta_B}{r'_S} e^{r'_S t_B}\right) & 0 \leq t_A < T \leq t_B \\ C_3 \cdot e^{r'_S t_A} \exp\left(\frac{-\beta_A}{r'_S} e^{r'_S t_A}\right) \cdot e^{r_S t_B} \exp\left(\frac{-\alpha_B}{r_S} e^{r_S t_B}\right) & 0 \leq t_B < T \leq t_A \\ C_4 \cdot e^{r'_S t_A} \exp\left(\frac{-\beta_A}{r'_S} e^{r'_S t_A}\right) \cdot e^{r'_S t_B} \exp\left(\frac{-\beta_B}{r'_S} e^{r'_S t_B}\right) & T \leq t_A, t_B \end{cases} \quad (20)$$

The simplifying constants C_j are defined as follows:

$$\begin{aligned} C_1 &= \frac{\alpha_A \eta_A}{1 - \nu_A} \frac{\alpha_B \eta_B}{1 - \nu_B} \\ C_2 &= \frac{\alpha_A \eta_A}{1 - \nu_A} \frac{\beta_B \nu_B}{1 - \nu_B} \\ C_3 &= \frac{\beta_A \nu_A}{1 - \nu_A} \frac{\alpha_B \eta_B}{1 - \nu_B} \\ C_4 &= \frac{\beta_A \nu_A}{1 - \nu_A} \frac{\beta_B \nu_B}{1 - \nu_B} \end{aligned} \quad (21)$$

An example joint density function $f_{A,B}^*(t_A, t_B)$ is shown in **Supplementary Figure 3**.




Supplementary Figure 3: Joint pdf etc.

Note that $f_{A,B}^*(t_A, t_B)$ is a piecewise function with four subdomains. Thus, when evaluating the integral in equation (16), it is necessary to separate it into appropriate subintegrals. The choice of subintegral and their associated limits of integration depend on the value of ΔT_{og} . From here on, we abbreviate ΔT_{og} to ΔT . Given our assumption that Allele A is more resistant than Allele B, $\Delta T = T_{og,B} - T_{og,A}$ must be positive. Consider the two cases:

Case 1: $0 \leq \Delta T < T$

Case 2: $T \leq \Delta T$

It seems most likely that detection time T will be larger than ΔT , so we focus our attention on **Case 1**.



Supplementary Figure 4: t_A - t_B plane etc

The region of integration is divided into four subregions in the t_A - t_B plane:

$$\iint_D f^* dA = \iint_{D_1} f^* dA + \iint_{D_2} f^* dA + \iint_{D_3} f^* dA + \iint_{D_4} f^* dA \quad (22)$$

We now take the integral for each individual subregion. For D_1 we have:

$$\begin{aligned}
\iint_{D_1} f^* dA &= C_1 \int_{\Delta T}^T e^{r_s t_A} \exp\left(-\frac{\alpha_A}{r_s} e^{r_s t_A}\right) \int_0^{t_A - \Delta T} e^{r_s t_B} \exp\left(-\frac{\alpha_B}{r_s} e^{r_s t_B}\right) dt_B dt_A \\
&= -\frac{C_1}{\alpha_B} \int_{\Delta T}^T e^{r_s t_A} \exp\left(-\frac{\alpha_A}{r_s} e^{r_s t_A}\right) \left[\exp\left(-\frac{\alpha_B}{r_s} e^{-r_s \Delta T} e^{r_s t_A}\right) - \exp\left(-\frac{\alpha_B}{r_s}\right) \right] dt_A \\
&= -\frac{C_1}{\alpha_B} \int_{\Delta T}^T e^{r_s t_A} \exp\left(\left(-\frac{1}{r_s}(\alpha_A + \alpha_B e^{-r_s \Delta T}) e^{r_s t_A}\right)\right) dt_A + \\
&\quad \frac{C_1}{\alpha_B} e^{-\alpha_B/r_s} \int_{\Delta T}^T e^{r_s t_A} \exp\left(-\frac{\alpha_A}{r_s} e^{r_s t_A}\right) dt_A \\
&= \frac{C_1}{\alpha_B(\alpha_A + \alpha_B e^{-r_s \Delta T})} \left[\exp\left(-\frac{e^{r_s T}}{r_s}(\alpha_A + \alpha_B e^{-r_s \Delta T})\right) - \exp\left(-\frac{e^{r_s \Delta T}}{r_s}(\alpha_A + \alpha_B e^{-r_s \Delta T})\right) \right] - \\
&\quad \frac{C_1}{\alpha_A \alpha_B} e^{-\alpha_B/r_s} \left[\exp\left(-\frac{\alpha_A}{r_s} e^{r_s T}\right) - \exp\left(-\frac{\alpha_A}{r_s} e^{r_s \Delta T}\right) \right]
\end{aligned} \tag{23}$$

For D_2 :

$$\begin{aligned}
\iint_{D_2} f^* dA &= C_3 \int_T^{T+\Delta T} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) \int_0^{t_A - \Delta T} e^{r_s t_B} \exp\left(-\frac{\alpha_B}{r_s} e^{r_s t_B}\right) dt_B dt_A \\
&= -\frac{C_3}{\alpha_B} \int_T^{T+\Delta T} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) \cdot \left[\exp\left(-\frac{\alpha_B}{r_s} e^{-r_s \Delta T} e^{r_s t_A}\right) - \exp\left(-\frac{\alpha_B}{r_s}\right) \right] dt_A \\
&= -\frac{C_3}{\alpha_B} \int_T^{T+\Delta T} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) \exp\left(-\frac{\alpha_B}{r_s} e^{-r_s \Delta T} e^{r_s t_A}\right) dt_A + \\
&\quad \frac{C_3}{\alpha_B} e^{-\alpha_B/r_s} \int_T^{T+\Delta T} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) dt_A
\end{aligned} \tag{24}$$

The second integral in the last form of equation (24) has a straightforward solution:

$$\begin{aligned}
\frac{C_3}{\alpha_B} e^{-\alpha_B/r_s} \int_T^{T+\Delta T} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) dt_A &= \\
&= -\frac{C_3}{\beta_A \alpha_B} e^{-\alpha_B/r_s} \left[\exp\left(-\frac{\beta_A}{r'_s} e^{r'_s(T+\Delta T)}\right) - \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s T}\right) \right]
\end{aligned} \tag{25}$$

However, the first integral in the last form of equation (24) requires more analysis. Let $u = e^{r'_s t_A}$, $du = r'_s e^{r'_s t_A} dt_A$, $u_1 = e^{r'_s T}$, and $u_2 = e^{r'_s(T+\Delta T)}$.

Now, using the power series for exponential functions:

$$\begin{aligned}
& -\frac{C_3}{\alpha_B} \int_T^{T+\Delta T} e^{r'_S t_A} \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S t_A}\right) \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} e^{r_S t_A}\right) dt_A \\
& = -\frac{C_3}{\alpha_B r'_S} \int_{u_1}^{u_2} \exp\left(-\frac{\beta_A}{r'_S} u\right) \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} u^{r_S/r'_S}\right) du \\
& = -\frac{C_3}{\alpha_B r'_S} \int_{u_1}^{u_2} \sum_{k=0}^{\infty} \frac{(-u\beta_A/r'_S)^k}{k!} \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} u^{r_S/r'_S}\right) du \\
& = -\frac{C_3}{\alpha_B r'_S} \sum_{k=0}^{\infty} \frac{(-\beta_A/r'_S)^k}{k!} \int_{u_1}^{u_2} \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} u^{r_S/r'_S}\right) u^k du
\end{aligned} \tag{26}$$

Let $v = \frac{\alpha_B}{r_S} e^{-r_S \Delta T} u^{r_S/r'_S}$, $dv = \frac{\alpha_B}{r'_S} e^{-r_S \Delta T} u^{(r_S/r'_S)-1} du$, $v_1 = \frac{\alpha_B}{r_S} e^{r_S(T-\Delta T)}$ and $v_2 = \frac{\alpha_B}{r_S} e^{r_S T}$. Solving for u and du , we have $u = \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T} v\right)^{r'_S/r_S}$ and $du = \frac{r'_S}{r_S} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{r'_S/r_S} v^{(r'_S/r_S)-1} dv$. Now,

$$\begin{aligned}
& -\frac{C_3}{\alpha_B} \int_T^{T+\Delta T} e^{r'_S t_A} \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S t_A}\right) \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} e^{r_S t_A}\right) dt_A \\
& = -\frac{C_3}{\alpha_B r'_S} \sum_{k=0}^{\infty} \frac{(-\beta_A/r'_S)^k}{k!} \int_{v_1}^{v_2} e^{-v} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T} v\right)^{kr'_S/r_S} \cdot \frac{r'_S}{r_S} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{r'_S/r_S} v^{r'_S/r_S-1} dv \\
& = -\frac{C_3}{\alpha_B r'_S} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{r'_S/r_S} \sum_{k=0}^{\infty} \frac{(-\beta_A/r'_S)^k}{k!} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{kr'_S/r_S} \int_{v_1}^{v_2} e^{-v} v^{(kr'_S/r_S+r'_S/r_S-1)} dv
\end{aligned} \tag{27}$$

Finally, using the integral definition of the lower incomplete gamma function $\gamma(a, x) = \int_0^x t^{a-1} e^{-t} dt$, we arrive at an asymptotic solution for this integral:

$$\begin{aligned}
& -\frac{C_3}{\alpha_B} \int_T^{T+\Delta T} e^{r'_S t_A} \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S t_A}\right) \exp\left(-\frac{\alpha_B}{r_S} e^{-r_S \Delta T} e^{r_S t_A}\right) dt_A \\
& = K \sum_{k=0}^{\infty} \left\{ \frac{(-\beta_A/r'_S)^k}{k!} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{kr'_S/r_S} \right. \\
& \quad \left. \left[\gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S} e^{r_S T}\right) - \gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S} e^{r_S(T-\Delta T)}\right) \right] \right\}
\end{aligned} \tag{28}$$

where $K = -\frac{C_3}{\alpha_B r'_S} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T}\right)^{r'_S/r_S}$. Given that the error bound on the series in equation (28) is the $(k+1)$ th term of the series, we can evaluate the integral to a desired degree of precision.

For D_3 :

$$\begin{aligned}
\iint_{D_3} f^* dA &= C_3 \int_{T+\Delta T}^{\infty} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) dt_A \int_0^T e^{r_s t_B} \exp\left(-\frac{\alpha_B}{r_s} e^{r_s t_B}\right) dt_B \\
&= \frac{C_3}{\beta_A \alpha_B} \left[1 - \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s (T+\Delta T)}\right)\right] \cdot \left[\exp\left(-\frac{\alpha_B}{r_s} e^{r_s T}\right) - \exp\left(-\frac{\alpha_B}{r_s}\right)\right]
\end{aligned} \tag{29}$$

For D_4 :

$$\begin{aligned}
\iint_{D_4} f^* dA &= C_4 \int_{T+\Delta T}^{\infty} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) \int_T^{t_A-\Delta T} e^{r'_s t_B} \exp\left(-\frac{\beta_B}{r'_s} e^{r'_s t_B}\right) dt_B dt_A \\
&= -\frac{C_4}{\beta_B} \int_{T+\Delta T}^{\infty} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) [\exp\left(-\frac{\beta_B}{r'_s} e^{-r'_s \Delta T} e^{r'_s t_A}\right) - \exp\left(-\frac{\beta_B}{r'_s} e^{r'_s T}\right)] dt_A \\
&= -\frac{C_4}{\beta_B} \int_{T+\Delta T}^{\infty} e^{r'_s t_A} \exp\left(-\frac{e^{r'_s t_A}}{r'_s} (\beta_A + \beta_B e^{-r'_s \Delta T})\right) dt_A + \\
&\quad \frac{C_4}{\beta_B} \exp\left(-\frac{\beta_B}{r'_s} e^{r'_s T}\right) \int_{T+\Delta T}^{\infty} e^{r'_s t_A} \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s t_A}\right) dt_A \\
&= \frac{C_4}{\beta_B (\beta_A + \beta_B e^{-r'_s \Delta T})} \left[1 - \exp\left(-\frac{e^{r'_s (T+\Delta T)}}{r'_s} (\beta_A - \beta_B e^{-r'_s \Delta T})\right)\right] - \\
&\quad \frac{C_4}{\beta_A \beta_B} \exp\left(-\frac{\beta_B}{r'_s} e^{r'_s T}\right) \left[1 - \exp\left(-\frac{\beta_A}{r'_s} e^{r'_s (T+\Delta T)}\right)\right]
\end{aligned} \tag{30}$$

Thus, the full expression for equation (22) is:

$$\begin{aligned}
\iint_D f^* dA &= \frac{C_1}{\alpha_B(\alpha_A + \alpha_B e^{-r_S \Delta T})} \left[\exp\left(-\frac{e^{r_S T}}{r_S}(\alpha_A + \alpha_B e^{-r_S \Delta T})\right) - \right. \\
&\quad \left. \exp\left(-\frac{e^{r_S \Delta T}}{r_S}(\alpha_A + \alpha_B e^{-r_S \Delta T})\right) \right] - \\
&\quad \frac{C_1}{\alpha_A \alpha_B} e^{-\alpha_B/r_S} \left[\exp\left(-\frac{\alpha_A}{r_S} e^{r_S T}\right) - \exp\left(-\frac{\alpha_A}{r_S} e^{r_S \Delta T}\right) \right] + \\
&\quad K \sum_{k=0}^{\infty} \left\{ \frac{(-\beta_A/r'_S)^k}{k!} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T} \right)^{kr'_S/r_S} \right. \\
&\quad \left. \left[\gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S} e^{r_S T}\right) - \gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S} e^{r_S(T-\Delta T)}\right) \right] \right\} - \\
&\quad \frac{C_3}{\beta_A \alpha_B} e^{-\alpha_B/r_S} \left[\exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S T}\right) \right] + \\
&\quad \frac{C_3}{\beta_A \alpha_B} \left[1 - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) \right] \cdot \left[\exp\left(-\frac{\alpha_B}{r_S} e^{r_S T}\right) - \exp\left(-\frac{\alpha_B}{r_S}\right) \right] + \\
&\quad \frac{C_4}{\beta_B(\beta_A + \beta_B e^{-r_S \Delta T})} \left[1 - \exp\left(-\frac{e^{r'_S(T+\Delta T)}}{r'_S}(\beta_A - \beta_B e^{-r'_S \Delta T})\right) \right] - \\
&\quad \frac{C_4}{\beta_A \beta_B} \exp\left(-\frac{\beta_B}{r'_S} e^{r'_S T}\right) \left[1 - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) \right]
\end{aligned} \tag{31}$$

Similarly, it can be shown that $\iint_D f^* dA$ for **Case 2** ($T \leq \Delta T$) is:

$$\begin{aligned}
\iint_D f^* dA &= K \sum_{k=0}^{\infty} \left\{ \frac{(-\beta_A/r'_S)^k}{k!} \left(\frac{r_S}{\alpha_B} e^{r_S \Delta T} \right)^{kr'_S/r_S} \right. \\
&\quad \left. \left[\gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S} e^{r_S T}\right) - \gamma\left(\frac{r'_S}{r_S}(k+1), \frac{\alpha_B}{r_S}\right) \right] \right\} - \\
&\quad \frac{C_3}{\beta_A \alpha_B} e^{-\alpha_B/r_S} \left[\exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S \Delta T}\right) \right] + \\
&\quad \frac{C_3}{\beta_A \alpha_B} \left[1 - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) \right] \cdot \left[\exp\left(-\frac{\alpha_B}{r_S} e^{r_S T}\right) - \exp\left(-\frac{\alpha_B}{r_S}\right) \right] + \\
&\quad \frac{C_4}{\beta_B(\beta_A + \beta_B e^{-r'_S \Delta T})} \left[1 - \exp\left(-\frac{e^{r'_S(T+\Delta T)}}{r'_S}(\beta_A + \beta_B e^{-r'_S \Delta T})\right) \right] - \\
&\quad \frac{C_4}{\beta_A \beta_B} \exp\left(-\frac{\beta_B}{r'_S} e^{r'_S T}\right) \left[1 - \exp\left(-\frac{\beta_A}{r'_S} e^{r'_S(T+\Delta T)}\right) \right]
\end{aligned} \tag{32}$$

5 Conclusion

In summary, the probability that either allele is dominant upon relapse is the sum of the probability of its fixation and the probability of it outcompeting the other allele (Equation (1)).

The probability of each allele's fixation is given by equations (10) and (11):

$$\begin{aligned} P[\text{Allele A is fixed}] &= P[\mathbf{Outcome\ 2}] = (1 - \nu_A) \cdot \nu_B \\ P[\text{Allele B is fixed}] &= P[\mathbf{Outcome\ 3}] = \nu_A \cdot (1 - \nu_B) \end{aligned} \quad (33)$$

The probability of either allele outcompeting the other allele is given by equations (11), (13), and (15):

$$\begin{aligned} P[\text{Allele A outcompetes}] &= P[\mathbf{Outcome\ 4}] \cdot P[\mathbf{Outcome\ 4.A} | \mathbf{Outcome\ 4}] \\ &= (1 - \nu_A) \cdot (1 - \nu_B) \cdot \left(1 - \iint_D f_{A,B}^*(t_A, t_B) dA\right) \\ P[\text{Allele B outcompetes}] &= P[\mathbf{Outcome\ 4}] \cdot P[\mathbf{Outcome\ 4.B} | \mathbf{Outcome\ 4}] \\ &= (1 - \nu_A) \cdot (1 - \nu_B) \cdot \iint_D f_{A,B}^*(t_A, t_B) dA \end{aligned} \quad (34)$$

where $\iint_D f_{A,B}^*(t_A, t_B) dA$ is given by equation (29) for **Case 1** ($0 \leq \Delta T < T$) and equation (30) for **Case 2** ($T \leq \Delta T$).

Thus, the probability that either allele is dominant upon relapse is given by substituting equations (33) and (34) into equation (1):

$$\begin{aligned} P[\text{Allele A is dominant upon relapse}] &= P[\mathbf{Outcome\ 2}] + P[\mathbf{Outcome\ 4}] \cdot P[\mathbf{Outcome\ 4.A} | \mathbf{Outcome\ 4}] \\ &= (1 - \nu_A) \cdot \nu_B + (1 - \nu_A) \cdot (1 - \nu_B) \cdot \left(1 - \iint_D f_{A,B}^*(t_A, t_B) dA\right) \\ P[\text{Allele B is dominant upon relapse}] &= P[\mathbf{Outcome\ 3}] + P[\mathbf{Outcome\ 4}] \cdot P[\mathbf{Outcome\ 4.B} | \mathbf{Outcome\ 4}] \\ &= (1 - \nu_B) \cdot \nu_A + (1 - \nu_A) \cdot (1 - \nu_B) \cdot \iint_D f_{A,B}^*(t_A, t_B) dA. \end{aligned} \quad (35)$$