

## Eigenvalue Problems

**Problem 1.** Find the positive eigenvalues and eigenfunctions of

$$\begin{cases} X^{(4)} = \lambda X & 0 < x < L \\ X(0) = X(L) = X''(0) = X''(L) = 0 \end{cases}$$

**Solution 1.** We want to find non-trivial solutions to the eigenvalue problem i.e.  $X(x) \neq 0$ .

*Step 1:* We are interested in positive eigenvalues, so we can set  $\lambda = \beta^4 > 0$ , where  $\beta > 0$ . We first find the general solution to the ODE

$$X^{(4)} = \beta^4 X \quad 0 < x < L.$$

This is a fourth order constant coefficient ODE with roots  $r = \pm\beta, \pm\beta i$ , which corresponds to the solution

$$X(x) = A \cos(\beta x) + B \sin(\beta x) + C \cosh(\beta x) + D \sinh(\beta x).$$

*Step 2:* We now solve for the values of  $\beta$  that satisfy the boundary conditions. Plugging the solution into the initial conditions gives

$$\begin{aligned} A + C &= 0 \\ A \cos(\beta L) + B \sin(\beta L) + C \cosh(\beta L) + D \sinh(\beta L) &= 0 \\ -\beta^2 A + \beta^2 C &= 0 \\ -A\beta^2 \cos(\beta L) - B\beta^2 \sin(\beta L) + C\beta^2 \cosh(\beta L) + D\beta^2 \sinh(\beta L) &= 0. \end{aligned}$$

Since  $\beta > 0$ , the first and third equation implies  $A + C = 0$  and  $-A + C = 0$  which can only happen when  $A = C = 0$ . We now have to solve the system

$$\begin{aligned} B \sin(\beta L) + D \sinh(\beta L) &= 0 \\ -B\beta^2 \sin(\beta L) + D\beta^2 \sinh(\beta L) &= 0. \end{aligned}$$

We can write this system of equations in matrix form

$$\begin{bmatrix} \sin(\beta L) & \sinh(\beta L) \\ -\beta^2 \sin(\beta L) & \beta^2 \sinh(\beta L) \end{bmatrix} \begin{bmatrix} B \\ D \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

which has a non-trivial solution when

$$\det \left( \begin{bmatrix} \sin(\beta L) & \sinh(\beta L) \\ -\beta^2 \sin(\beta L) & \beta^2 \sinh(\beta L) \end{bmatrix} \right) = 0 \implies 2\beta^2 \sinh(\beta L) \sin(\beta L) = 0.$$

Since  $\beta > 0$  and  $\sinh(\beta L) > 0$ , this simplifies to  $\sin(\beta L) = 0$ , which occurs precisely when

$$\beta_n = \frac{n\pi}{L}, \quad n \geq 1.$$

Therefore, the corresponding eigenvalues are  $\lambda_n = \frac{n^4 \pi^4}{L^4}$ . Furthermore, notice that from the equation

$$B \sin(\beta_n L) + D \sinh(\beta_n L) = 0,$$

we must have  $D = 0$  since  $\sinh(\beta_n L) > 0$  and  $\sin(\beta_n L) = 0$ . Finally, for  $n \geq 1$ , the corresponding eigenfunction (taking  $B = 1$ ) for the eigenvalue  $\lambda_n = \left(\frac{n\pi}{L}\right)^4$  is

$$X_n(x) = \sin\left(\frac{n\pi x}{L}\right).$$

## Separation of Variables

**Problem 2.** Consider wave equation with the Neumann boundary condition on the left and weird b.c. on the right:

$$\begin{aligned} u_{tt} - c^2 u_{xx} &= 0 & 0 < x < l, \\ u_x(0, t) &= 0, \\ (u_x + i\alpha u_t)(l, t) &= 0 \end{aligned}$$

with  $\alpha \in \mathbb{R}$ .

1. Separate variables
2. Find the “weird” eigenvalue problem for the ODE;
3. Solve the problem;
4. Find the simple solution  $u(x, t) = X(x)T(t)$ .

**Solution 2.** We will use the method of separation of variables to find a complex solution to this PDE.

*Step 1 — Separation of Variables:* We look for a separated solution  $u(x, t) = T(t)X(x)$  to our PDE. Plugging this into our PDE gives

$$T''(t)X(x) - c^2 T(t)X''(x) = 0 \implies \frac{T''(t)}{c^2 T(t)} = \frac{X''(x)}{X(x)} = \lambda.$$

This gives the following ODEs

$$X''(x) - \lambda X(x) = 0 \text{ and } T''(t) - c^2 \lambda T(t) = 0,$$

with boundary conditions

$$X'(0)T(t) = 0, \quad X'(l)T(t) + i\alpha X(l)T'(t) = 0.$$

Instead of trying to find an eigenvalue problem for  $X(x)$  like in the usual cases, we first solve the second order ODEs, which gives

$$X_\lambda(x) = Ae^{\sqrt{\lambda}x} + Be^{-\sqrt{\lambda}x}, \quad T_\lambda(t) = Ce^{c\sqrt{\lambda}t} + De^{-c\sqrt{\lambda}t} \quad \text{when } \lambda \neq 0.$$

and

$$X_0(x) = A + Bx, \quad T_0(t) = C + Dt \quad \text{when } \lambda = 0.$$

*Step 2 — Solve for  $\lambda$ :* We use the boundary conditions to solve for  $\lambda$ , and its corresponding eigenfunctions  $X_\lambda$  and  $T_\lambda$ .

**Zero Eigenvalue:** When  $\lambda = 0$ , then the first initial condition implies that

$$B = 0,$$

so  $X_0(x) = A$ . The second initial condition, then implies that

$$i\alpha DA = 0 \implies D = 0,$$

otherwise we are left with a trivial solution. Therefore, we must have  $T_0(t) = C$ , which means that the constant function

$$u(x, t) = X_0(x)T_0(t) \equiv E$$

where  $E \in \mathbb{C}$  is the solution corresponding to the zero eigenvalue.

**Non-Zero Eigenvalues:** We now find  $\lambda \neq 0$  that satisfies the boundary conditions

$$T_\lambda(t)X'_\lambda(0) = 0, \quad (1)$$

$$X'_\lambda(l)T_\lambda(t) + i\alpha X_\lambda(l)T'_\lambda(t) = 0. \quad (2)$$

1. First Boundary Condition: For all  $t$ , we must have

$$(A\sqrt{\lambda} - B\sqrt{\lambda})(Ce^{c\sqrt{\lambda}t} + De^{-c\sqrt{\lambda}t}) = 0.$$

Since  $\sqrt{\lambda} \neq 0$  and the equation must hold for all  $t$ , we need  $A = B \neq 0$ . In other words, we need  $X_\lambda(x)$  to be an even function.

2. Second Boundary Condition: For all  $t$ , we must have

$$A(\sqrt{\lambda}e^{\sqrt{\lambda}l} - \sqrt{\lambda}e^{-\sqrt{\lambda}l})(Ce^{c\sqrt{\lambda}t} + De^{-c\sqrt{\lambda}t}) + i\alpha cA(e^{\sqrt{\lambda}l} + e^{-\sqrt{\lambda}l})(C\sqrt{\lambda}e^{c\sqrt{\lambda}t} - D\sqrt{\lambda}e^{-c\sqrt{\lambda}t}) = 0$$

and rearranging and dividing out by  $A\sqrt{\lambda} \neq 0$  gives

$$(e^{\sqrt{\lambda}l} - e^{-\sqrt{\lambda}l})(Ce^{c\sqrt{\lambda}t} + De^{-c\sqrt{\lambda}t}) + i\alpha c(e^{\sqrt{\lambda}l} + e^{-\sqrt{\lambda}l})(Ce^{c\sqrt{\lambda}t} - De^{-c\sqrt{\lambda}t}) = 0$$

We can now use the fact  $e^{c\sqrt{\lambda}t}$  and  $e^{-c\sqrt{\lambda}t}$  are linearly independent to equate coefficients to solve for  $\lambda$  such that the above equation holds for all  $t$ . If  $D = 0$  and  $C \neq 0$ , we can divide by  $e^{c\sqrt{\lambda}t}$  to conclude that

$$(e^{\sqrt{\lambda}l} - e^{-\sqrt{\lambda}l}) + (e^{\sqrt{\lambda}l} + e^{-\sqrt{\lambda}l})i\alpha c = 0 \implies \tanh(\sqrt{\lambda}l) = -i\alpha c \implies \sqrt{\lambda} = \frac{i(\pi n - \tan^{-1}(\alpha c))}{l}$$

and if  $C = 0$  and  $D \neq 0$  we can divide by  $e^{-c\sqrt{\lambda}t}$  to conclude that

$$(e^{\sqrt{\lambda}l} - e^{-\sqrt{\lambda}l}) - (e^{\sqrt{\lambda}l} + e^{-\sqrt{\lambda}l})i\alpha c = 0 \implies \tanh(\sqrt{\lambda}l) = i\alpha c \implies \sqrt{\lambda} = \frac{i(\pi n + \tan^{-1}(\alpha c))}{l}.$$

We used the identity  $\tanh(x) = -i \tan(ix)$  in our simplification above.

We get two families of solutions depending on the eigenvalue,

$$X_{\lambda_n}(x)T_{\lambda_n}(t) = (e^{\sqrt{\lambda_n}x} + e^{-\sqrt{\lambda_n}x})e^{c\sqrt{\lambda_n}t} \quad \text{where } \sqrt{\lambda_n} = \frac{i(\pi n - \tan^{-1}(\alpha c))}{l}$$

and

$$X_{\gamma_n}(x)T_{\gamma_n}(t) = (e^{\sqrt{\gamma_n}x} + e^{-\sqrt{\gamma_n}x})e^{-c\sqrt{\gamma_n}t} \quad \text{where } \sqrt{\gamma_n} = \frac{i(\pi n + \tan^{-1}(\alpha c))}{l}.$$

*Step 3 — General Simple Solution:* Therefore, taking the linear combination of our solutions, we have

$$u(x, t) = E + \sum_{i=-\infty}^{\infty} C_n(e^{\sqrt{\lambda_n}x} + e^{-\sqrt{\lambda_n}x})e^{c\sqrt{\lambda_n}t} + \sum_{i=-\infty}^{\infty} D_n(e^{\sqrt{\gamma_n}x} + e^{-\sqrt{\gamma_n}x})e^{-c\sqrt{\gamma_n}t}$$

where  $C_n, D_n, E \in \mathbb{C}$  and

$$\sqrt{\lambda_n} = \frac{i(\pi n - \tan^{-1}(\alpha c))}{l}, \quad \sqrt{\gamma_n} = \frac{i(\pi n + \tan^{-1}(\alpha c))}{l}.$$