Counting Hunger, Missing Deaths: What Food Insecurity Phases Tell Us About Malnutrition and Mortality

Claude Opus 5.5* Charles Kenny† Justin Sandefur‡

October 2026

Working draft — not for citation

Abstract

The humanitarian system counts need in phases of acute food insecurity. The Integrated Food Security Phase Classification (IPC) puts each area on a five-point scale, and the number of people in its third phase or worse is the headline figure behind appeals, famine warnings and the U.S.-funded Famine Early Warning Systems Network (FEWS NET), which forecasts it. We compare these phases with outcomes measured independently in surveys. Phases track household food consumption and coping, loosely: 47 percent of households in areas mapped in Crisis had poor or borderline diets, against 28 percent in areas mapped Minimal. They are a weak guide to child malnutrition and a poor guide to death. In Somalia's scheduled surveys, acute malnutrition averaged 13, 13 and 15 percent in zones forecast to be Stressed, in Crisis and in Emergency; across other countries, crude death rates do not rise with FEWS NET's phase at all. The IPC's reference bands imply gradients several times steeper than those we measure across countries, and roughly ten times steeper than those within Somali zones over time. This fits an old finding: most people who die in food crises die of disease, as health, water and sanitation break down, not of a shortfall in food consumption alone. A response targeted on food insecurity targets a small part of the problem. Scheduled nutrition and mortality surveys do better, and so do cheap signals of drought and conflict: in Somalia they predict malnutrition and death rates out of sample where the phases predict nothing. A zone's last survey predicts its next one far better than the forecast phase does, and if treatment for malnourished children were placed by survey results, Somalia's survey rounds would avert deaths at about $1,200–2,200 each. But surveys avoid the places that most need them: household surveys are postponed in years of conflict, and 65 percent of those run in wartime document leaving out insecure areas. The case for reform concerns the IPC and the way phase counts are used, more than FEWS NET, which forecasts the IPC reasonably well.

*This paper is fully AI generated. Claude Opus 5.5 assembled the data, wrote the code, ran the analysis and drafted the text. Human direction, shared with the companion paper (Claude Opus 5.5 et al., 2026), consisted of 98 prompts totaling about 5,200 words between 30 September and 14 October 2026. Replication code is available at `https://github.com/justinsandefur/fewsnet-evaluation`.

†Center for Global Development.

‡Coefficient Giving.

JEL codes: I15, O19, Q18, F35. **Keywords:** famine, food insecurity, acute malnutrition, mortality, humanitarian aid, surveys.

1 Introduction

In 2022 Somalia was the world's most-watched hunger crisis. Between 5.6 and 6.7 million people were classified in Crisis or worse on the IPC scale, two districts were projected to reach Famine, and agencies raised money on that basis. An estimated 43,000 people died in excess of normal that year, about half of them children under five, and most of them not in the small areas at risk of Famine but in the much larger populations classified in Crisis and Emergency (London School of Hygiene & Tropical Medicine et al., 2023). That is a terrible toll. It is also, by the IPC's own reference bands, two to five times smaller than the toll the phase counts implied. Something in the chain from phase to death did not hold.

This paper asks what acute food insecurity phases measure. The IPC and the Cadre Harmonisé, its West African counterpart, classify areas on a five-phase scale — Minimal, Stressed, Crisis, Emergency and Famine — that is meant to summarize whether households can eat enough and what they give up to do so (IPC Global Partners, 2021). FEWS NET produces the same classification and forecasts it months ahead. The number of people in Phase 3 or worse has become the humanitarian system's common currency: it heads the Global Report on Food Crises (FSIN and Global Network Against Food Crises, 2026), it sizes appeals, and it is how the world decides whether a crisis is getting worse. The IPC also attaches to each phase a band of child acute malnutrition and of crude death rates, so a phase count is easily read as a count of children wasting and people dying.

We test that reading against outcomes measured independently, in four steps. First, the like-for-like test: do phases match household food consumption and coping measured in surveys the IPC did not run? Second and third, do they match acute malnutrition among children and death rates measured in nutrition and mortality surveys? Here the cleanest evidence comes from Somalia, where the Food Security and Nutrition Analysis Unit (FSNAU) surveys rural zones on a fixed calendar twice a year, whatever conditions are, so that results are not driven by where agencies chose to look. Fourth, we ask whether those surveys would be worth paying for in their own right: do they predict malnutrition and death, and would a response that followed them save lives?

The answers are consistent. Phases track household food insecurity, loosely, and look too mild in areas mapped as Minimal or Stressed. They are a weak guide to child malnutrition: within a Somali zone, a phase worse than usual goes with about 0.8 more points of acute malnutrition, where the IPC's bands imply 5 to 10. They are a poor guide to mortality: across 246 recent surveys elsewhere, crude death rates are flat across FEWS NET's phases. The IPC's own malnutrition scale agrees with its food insecurity scale for only 35 percent of areas. Surveys, by contrast, carry the information that phases lack. A zone's previous survey explains 0.42 of the variance in its next survey's acute malnutrition, out of sample, and FEWS NET's forecast explains none. A planner placing treatment for malnourished children by survey results rather than forecasts would avert deaths at about \$1,200–2,200 each.

We see this paper as a continuation of an argument made long ago. Alex de Waal, studying Darfur in 1984–85, showed that most people who die in famines die of disease — diarrhoea, measles, malaria and respiratory infection — spread by displacement, crowding, bad water and the collapse of

health services, with hunger one cause among several (de Waal, 1989b). Refugee epidemiologists and economic historians found the same (Toole and Waldman, 1990; Mokyr and Ó Gráda, 2002; Salama et al., 2004). If they are right, phases of food insecurity cannot be a good guide to malnutrition and death, and the bands that connect them cannot hold. Our first contribution is to show, with the data the system itself relies on, that they do not. Our second is practical: to take the old argument seriously would mean measuring the final outcomes — malnutrition and death — regularly, and directing attention by them. We estimate what that would be worth. That too has been proposed before, under the name of nutritional surveillance (Mason et al., 1984; Bilukha et al., 2012), and Section 10 is explicit about what is old here and what is not.

Two limits frame the argument. First, malnutrition and mortality are not what the food insecurity scale classifies; the IPC says so, and in 2016 it built a separate scale for acute malnutrition (IPC Nutrition Working Group, 2016). Our point is not that the scale fails at its own task but that the system uses it for another. Second, the surveys that measure outcomes best do not go everywhere. Using metadata for every DHS, MICS and World Bank living standards survey since 1990, we find that household surveys are postponed in years of conflict and economic collapse, and that when they do go to war-affected countries, most leave out insecure areas, and almost none say anything about camps. A survey-based system would need to be built for exactly the places current surveys avoid.

The implications concern the IPC more than FEWS NET. A companion paper finds that FEWS NET's forecasts carry information about later IPC classifications that other sources lack, and that humanitarian money follows them (Claude Opus 5.5 et al., 2026). FEWS NET predicts the IPC; the weakness lies in what the IPC's phases capture and in how phase counts are used to stand for malnutrition and death. Section 11 argues for reporting measured malnutrition and mortality alongside phases, for scheduled surveys and surveillance in high-risk areas, and for allocating life-saving treatment by measured need.

The paper proceeds as follows. Section 2 describes the IPC and how its phases are used. Section 3 describes the data. Sections 4, 5 and 6 compare phases with food insecurity, malnutrition and death. Section 7 asks why the link is weak. Section 8 asks what surveys add and what they would be worth, and Section 9 where they do not go. Section 10 places the argument in older debates and Section 11 concludes.

2 What a phase is, and how it is used

The IPC began in Somalia in 2004, built by FSNAU on an earlier proposal to grade famines by intensity and magnitude (Howe and Devereux, 2004), and became a global partnership of UN agencies, NGOs and governments. Its acute food insecurity scale classifies an area in the highest phase that describes at least 20 percent of its households. Analysts meet, review the available evidence — household food consumption and coping, livelihood change, nutrition, mortality, and contributing factors such as rainfall, prices and conflict — and reach a consensus phase. The phase

is a judgment that converges evidence, not a formula applied to it (IPC Global Partners, 2021).

The IPC's reference table attaches indicative values to each phase. For households, borderline food consumption and a reduced coping strategies index of 19 or more correspond to Crisis, and poor consumption to Emergency or worse. For outcomes, it attaches bands of acute malnutrition among children under five (below 5 percent in Phase 1, 5–10 in Phase 2, 10–15 in Crisis, 15–30 in Emergency and above 30 in Famine) and of crude death rates (below 0.5 deaths per 10,000 people per day in Phases 1 and 2, 0.5–1 in Crisis, 1–2 in Emergency and above 2 in Famine) (IPC Global Partners, 2019). Famine requires all three thresholds: 20 percent of households facing extreme food gaps, acute malnutrition of 30 percent and a death rate of 2. Below Famine, malnutrition and mortality are supporting evidence, often missing.

The bands were not derived from food insecurity data. They descend from benchmarks for refugee camps in the 1980s and 1990s — a doubling of a baseline crude death rate of about 0.44, and camp-based relations between wasting and mortality (Young and Jaspars, 2009). When the IPC built its separate acute malnutrition scale in 2016, it noted that food insecurity and acute malnutrition often diverge, in both directions (IPC Nutrition Working Group, 2016; IPC Global Partnership, 2024).

Phase counts are used for much more than classification. The number of people in Phase 3 or worse is the headline of the Global Report on Food Crises and of most humanitarian appeals that involve food (FSIN and Global Network Against Food Crises, 2026). FEWS NET, funded by the United States since 1985, forecasts the same phases four to eight months ahead in about 30 countries, and the companion paper shows that humanitarian funding follows its forecasts (Claude Opus 5.5 et al., 2026). The bands invite a further step, which evaluations and advocates take: converting people in Crisis or worse into expected deaths. That conversion is what this paper tests.

3 Data

We match outcomes measured in surveys to the phase FEWS NET or the IPC gave the same area at the same time. None of the outcome data were produced by the IPC or FEWS NET, though their analysts may have seen them.

- **Household food security.** WFP's household food security surveys and its monthly phone monitoring, compiled by the European Commission's Joint Research Centre (Machefer et al., 2025) and from WFP's monitoring files, by region: 478 areas in 29 countries, 2011–2024. We use the share of households with poor or borderline food consumption and the share coping severely (a reduced coping strategies index of 19 or more). For 2021–2023 the Cadre Harmonisé also publishes the household indicators its own analysts weighed, for 2,805 area-analyses in 18 countries.
- **Somalia's scheduled surveys.** FSNAU, managed by FAO, surveys rural livelihood zones, towns and displaced settlements after each of the two rainy seasons. We assembled all 1,248 results it published from late 2010 to 2026 and kept the rural surveys it ran itself, matched

to FEWS NET's mapping units: 329 zone-seasons in 29 zones over 28 seasons.

- **Other nutrition and mortality surveys.** The SMART+ dashboard run by Action Against Hunger and UNICEF (246 survey results, 2022–2026, 167 in FEWS NET's maps), and national survey rounds in Burkina Faso and Mauritania from FEWS NET's data warehouse (389 results, 2003–2018).
- **The IPC's own two scales.** 1,427 areas classified on both the food insecurity and the acute malnutrition scales, or with published malnutrition prevalence we can classify, in nine countries, 2021–2026.
- **Excess deaths.** Published estimates of excess mortality in four documented crises, from small-area models fitted to mortality surveys (Checchi and Robinson, 2013; Warsame et al., 2023; London School of Hygiene & Tropical Medicine et al., 2023; Checchi et al., 2023).
- **Where surveys go.** Metadata for every DHS, MICS and World Bank LSMS survey in 143 low- and middle-income countries, 1990–2023, and the sampling appendices of 170 DHS and MICS reports from 2000 on (Section 9).

Which surveys are used matters. Ad hoc surveys are sent where agencies suspect trouble, so they overstate the link between phases and outcomes: they are fielded where both are bad. Scheduled surveys, fielded on a calendar whatever conditions are, are the cleaner test. We rely on them where we can and treat ad hoc surveys as an upper bound. Appendix A gives the estimating equations.

4 Phases and household food insecurity

The like-for-like test is whether phases match the household measures they are designed to summarize. They do, loosely (Figures 1 and 2). The share of households with poor or borderline food consumption averages 28, 35, 47 and 46 percent in areas FEWS NET mapped in Phases 1, 2, 3 and 4 or worse; the share coping severely averages 33, 35, 49 and 45 percent. The correlations with the phase are 0.41 and 0.35 across all areas, 0.27 and 0.25 within countries, and 0.15 and 0.25 within an area over time. The spread within each phase is wide, and there is no step up from Crisis to Emergency.

Where the phases and the indicators disagree, the phases are milder. Under the IPC's 20 percent rule, 71 percent of areas mapped in Phases 1 or 2 had too many households with poor or borderline consumption for their phase, and 78 percent had too many coping severely; only 5 and 4 percent of areas mapped in Crisis or worse had too few. The indicators are consistent with the classification in Crisis areas and much worse than it in areas called Minimal or Stressed. This fits Lentz et al. (2025), who find that official estimates undercount acute hunger, and Lentz et al. (2024), who find the IPC's analyses only partly consistent with their own evidence. It does not prove the phases wrong: survey indicators are noisy, and analysts weigh other evidence. But it means that, if anything, phase counts understate household food insecurity.




Figure 1: Household food consumption by FEWS NET phase. Each point is one area in one survey round (blue) or one area-year of WFP's monthly phone monitoring (orange): the share of households with poor or borderline food consumption, against the phase FEWS NET's map gave the area at the time (mean over the FEWS NET units in the area, rounded; points jittered). Data: WFP, compiled in the Joint Research Centre's harmonized food insecurity dataset and WFP's monitoring files, by region, 2011–2024. Grey boxes: the range consistent with each phase under the IPC's rule that an area is in a phase when at least 20 percent of its households are in that phase or worse, with borderline consumption counted as Crisis and poor consumption as Emergency or worse; below 20 percent for Phases 1 and 2, at least 20 percent for Crisis or worse (this indicator cannot separate Crisis from Emergency). Diamonds: means with 95 percent intervals.

The analysts' own inputs tell the same story. Even the indicators Cadre Harmonisé analysts recorded for an area correlate only loosely with the phase those analysts assigned: 0.33 for poor or borderline food consumption, 0.34 for moderate or severe household hunger, 0.15 for crisis livelihood coping and 0.04 for severe consumption coping. A phase is a consensus that weighs several kinds of evidence and adjusts for context. Its link to any single measured outcome is correspondingly loose. That is defensible for a summary of food insecurity. It is a problem when the phase is used as a proxy for outcomes it does not directly weigh.




Figure 2: Household coping by FEWS NET phase. As Figure 1, for the share of households using severe consumption coping strategies (reduced coping strategies index of 19 or more, which the IPC's reference table places at Crisis or worse). Grey boxes: the range consistent with each phase under the IPC's 20 percent rule.

5 Phases and child malnutrition

Somalia's scheduled surveys give the cleanest test. Mean acute malnutrition among children under five was 13 percent in zones FEWS NET forecast to be Stressed or better, 13 percent in zones forecast to be in Crisis, and 15 percent in zones forecast to be in Emergency or worse (Figure 3, blue points). The IPC bands for those phases are 5–10, 10–15 and 15–30 percent. The correlation between the forecast and the measured rate is 0.10. The correlation with FEWS NET's map published just after the survey, which could draw on its results, is 0.09.

Comparing zones with each other could hide a gradient over time, since some zones always have high malnutrition. So we also compare each zone with itself. When FEWS NET forecast a zone to be one phase worse than usual, its acute malnutrition was 0.8 points higher (standard error 0.3), or 0.5 points with season effects removed (Appendix Figure 9). The bands imply 5 points for a step from Stressed to Crisis and 10 for a step from Crisis to Emergency. The estimate may be biased down, because a survey zone's phase averages several FEWS NET units and the worst-hit zones are under-surveyed; but it is consistent with Reusken et al. (2025), who find a link between the official Phase 3+ share and later malnutrition in Somali districts that is real but far weaker than the bands.

Other countries show a steeper gradient, still well below the bands. In SMART+ surveys, mean




Figure 3: Measured acute malnutrition by FEWS NET phase. Each point is one survey: global acute malnutrition (weight-for-height) among children under five, against the phase FEWS NET's map gave the area at the time of the survey (rounded mean over the FEWS NET units in the survey area; points jittered). Grey boxes: the IPC reference band for each phase. Blue: Somalia's scheduled FSNAU rounds, rural zones, 2011–2024. Orange: other recurring national or sub-national rounds (Nigeria's UNICEF rounds, 2023–26; Burkina Faso and Mauritania national rounds, 2003–18). Green: ad hoc surveys in the SMART+ dashboard, 2022–26. Diamonds: mean of all surveys in the phase, with 95 percent intervals.

acute malnutrition was 9, 10, 11 and 18 percent across Phases 1, 2, 3 and 4 or worse, and one phase goes with 1.8 points (standard error 0.5) with country-by-year effects; the jump at Phase 4 comes mostly from a dozen ad hoc surveys in Ethiopia. Nigeria's recurring rounds give 2.7 points per Cadre Harmonisé phase, and the Burkina Faso and Mauritania rounds 3.3 per FEWS NET phase. The IPC's two scales tell the same story from the inside: where an area has both a food insecurity phase and an acute malnutrition phase for the same period, they agree only 35 percent of the time, and areas the food insecurity scale calls Crisis are spread almost evenly across all four malnutrition phases (Appendix Figure 11).

6 Phases and death

For mortality the link is weaker still. In the SMART+ surveys that measured it, crude death rates were 0.32, 0.45, 0.40 and 0.31 deaths per 10,000 people per day in areas FEWS NET mapped in Phases 1, 2, 3 and 4 or worse: flat, with a slope of 0.00 (standard error 0.04) (Figure 4). Against the official IPC or Cadre Harmonisé phase they rise by 0.17 per phase (0.05). The IPC bands step




Figure 4: Measured crude death rates by FEWS NET phase. As Figure 3, for crude death rates (deaths per 10,000 people per day), in surveys that measured mortality. Grey boxes: the IPC reference band for each phase.

by 0.5 to 1 per phase. Within Somali zones, a phase worse than usual goes with 0.12 more deaths per 10,000 per day (0.03), where the bands imply 0.45–0.75.

The crises whose death tolls have been estimated point the same way (Figure 5). Dividing excess deaths by the number of people in Crisis or worse gives an excess death rate per person in Phase 3 or worse of roughly 0.15–0.2 per 10,000 per day in Somalia's 2017–18 and 2022 droughts, about 0.9 in northeast Nigeria in 2016–19, and 1.1–1.9 in the 2010–12 Somalia famine, which met a late response (Warsame et al., 2023; London School of Hygiene & Tropical Medicine et al., 2023; Checchi et al., 2023; Checchi and Robinson, 2013). The two droughts in which aid arrived at scale sit well below the IPC's Crisis band; the two cases above it are a famine with a late response and a conflict. Measured crude death rates in areas reviewed for famine since 2017 have mostly been between 0.3 and 1.9, below the famine threshold of 2 (IPC Famine Review Committee, 2022, 2024b).

Death rates in crises vary enormously, and the phase captures little of that variation. Arithmetic with the bands would have predicted roughly two to five times the excess deaths estimated for Somalia in 2022. A death toll inferred from a phase count is not a measurement.




Figure 5: Excess deaths in documented crises, per person in Crisis or worse. Estimated excess deaths divided by the number of people classified in IPC or Cadre Harmonisé Phase 3 or worse and the length of the crisis; ranges reflect ranges in the death toll and in the Phase 3+ population. Grey band: the IPC Crisis band for crude death rates (0.5–1 per 10,000 per day) less a normal baseline of 0.2–0.3. Sources: Warsame et al. (2023); London School of Hygiene & Tropical Medicine et al. (2023); Checchi et al. (2023); Checchi and Robinson (2013).

7 Why the link is weak

The simplest explanation is that phases measure food insecurity well enough, and that food insecurity is only one cause of malnutrition and death in crises. That is an old finding, not ours.

What people die of. Since Alex de Waal's study of the 1984–85 Darfur famine, the consensus in humanitarian public health has been that most excess deaths in famines are not deaths from starvation. In Darfur the crude death rate roughly trebled, mostly through a seasonal spike in child deaths from diarrhoea, measles and malaria, and access to water predicted who died better than wealth did (de Waal, 1989b,a). Studies of refugee and displaced populations in the 1980s and 1990s found the same four killers — measles, diarrhoea, respiratory infection and malaria — made deadlier by malnutrition (Toole and Waldman, 1990, 1997). Economic historians find the same profile in Ireland, India and Russia before antibiotics (Mokyr and Ó Gráda, 2002). In Gode, Ethiopia, in 2000, measles alone or combined with malnutrition caused about a fifth of under-five deaths, and a measles campaign was mounted only after the bulk of food aid had arrived (Salama et al., 2001). In Darfur in 2003–08, more than 80 percent of an estimated 300,000 excess deaths were not violent, mostly diarrhoea among the displaced (Degomme and Guha-Sapir, 2010). Across 1,759 death rate estimates from crises between 1998 and 2012, non-violent excess deaths outnumber violent ones even in exceptionally brutal conflicts (Heudtlass et al., 2016).

What predicts malnutrition and death. The best statistical models of crisis mortality, fitted for Somalia, South Sudan and northeast Nigeria, lean on conflict, measles, malaria, displacement and admissions for severe malnutrition; none uses the IPC phase as a predictor (Checchi et al., 2022, 2023). Models of acute malnutrition in Somalia and South Sudan found livelihood, measles,

vegetation, terms of trade and water prices most useful, and still performed poorly (Checchi et al., 2022). Wasting in Somalia peaks around May–June and November and falls during the lean season, so it does not follow the food insecurity calendar (Luc et al., 2024); wasting peaks across Africa track heat and disease seasons more than the hunger gap (Venkat et al., 2023). In parts of Chad and South Sudan, acute malnutrition has stayed above the 15 percent emergency threshold for decades, including when the IPC classified the area as Minimal (Young and Marshak, 2017). A meta-analysis of household studies finds that food insecurity predicts stunting and underweight but not wasting (Moradi et al., 2019).

Why the bands are too steep. The bands came from camps, where populations were crowded, newly displaced and often hit by epidemics, and where death rates rose steeply with wasting (Young and Jaspars, 2009). The IPC applies them to whole rural livelihood zones whose baselines and disease environments differ. In south-central Somalia in 2010–12, death rates in areas classified Minimal or Stressed were already 0.5–0.8 per 10,000 per day, inside the Crisis band (Checchi and Robinson, 2013). The same absolute bands make the opposite error where baselines are low: in Gaza, a death rate meeting the Famine threshold would have required a more than twentyfold rise, against six- to sevenfold in rural Somalia (de Zwarte et al., 2026). Both problems come from fixing absolute death rates to phases defined by food.

Where food does matter. None of this says that food does not matter. In sieges and in populations with good water and health services, starvation itself can dominate, as in the European famines of the 1940s (Mokyr and Ó Gráda, 2002), and plausibly in Gaza. At the extreme, as in Somalia in 2011, food deprivation and disease compound each other, and the presence of food aid lowered death rates (Checchi and Robinson, 2013). Preventive food supplements and cash can reduce wasting at the margin (Isanaka et al., 2009; Langendorf et al., 2014). Our claim is narrower: across the large populations in Crisis and Emergency where most crisis deaths occur, the amount of food insecurity explains little of the variation in who becomes malnourished or dies.

Where the money goes. Humanitarian funding follows the food-insecurity frame. Across 2015–2024, the food security cluster received 27 percent of all humanitarian funding reported to OCHA's Financial Tracking Service, health 10 percent, nutrition 4 percent, and water, sanitation and hygiene 4 percent; about 30 percent was not assigned to a sector. Some of the food money saves lives, and some of it buys nutrition and health benefits. But if most deaths come from disease, a system that sizes and prioritizes crises by counts of food-insecure people is aiming at a small part of the problem.

8 What surveys add

If phases say little about malnutrition and death, what does? The simplest predictor of a zone's acute malnutrition is its last survey. In Somalia, a zone's result six months earlier correlates 0.66

with this round's (Appendix Figure 10). Out of sample, leaving out one season at a time, the previous survey explains 0.42 of the variance in acute malnutrition and FEWS NET's forecast explains none (R^2 of -0.10); adding the forecast to the previous survey does not help (0.41). Given one zone that reaches the critical level of 15 percent and one that does not, the previous survey ranks them correctly 0.75 of the time; the forecast does no better than a coin toss (0.47). The same holds for under-five death rates: an out-of-sample R^2 of 0.20 for the previous survey and -0.07 for the forecast. Models that try to predict malnutrition without recent surveys, from rainfall, prices, conflict and disease, have performed poorly (Checchi et al., 2022).

FEWS NET barely uses this information. A Somali zone whose last survey showed 10 more points of acute malnutrition was forecast 0.09 of a phase worse (standard error 0.05). That is not a failure of FEWS NET's analysts: their scale is about food insecurity, and survey results about malnutrition do not translate into it. Where surveys do move FEWS NET's maps, they move them in one direction only. In Somalia, Ethiopia and Sudan, 263 survey results are quoted in FEWS NET's reports; when one was carried out between a forecast and the next map, that map was 8.8 percentage points (standard error 3.1) more likely to be worse than forecast, from a base of 10 percent, and no more likely to be better. Surveys showing acute malnutrition below 15 percent moved the maps up about as much as alarming ones (14.8 against 11.9 points). Agencies send survey teams where they suspect trouble, and the decision to survey is itself the signal. That is useful, but it is not the same as measuring.

Could cheaper signals do the job? Surveys are costly and do not go everywhere (Section 9). Drought, conflict and prices are observed everywhere, from satellites, conflict-event databases and markets. Do they predict malnutrition and death in ways the phases miss? We test this in Somalia, on 458 survey domains and seasons from 2015 to 2026 — rural zones, towns and displaced settlements — where we observe the official IPC share of the population in Phase 3 or worse, FEWS NET's map and forecast, and the triggers: rainfall anomalies over 6 and 12 months, conflict fatalities over 6 months, and cereal and goat prices, all measured up to the survey month.

The triggers beat the phases (Figure 6). Predicting each season's surveys from the others, the triggers explain 0.21 of the variance in acute malnutrition and 0.19 in under-five death rates; the phases explain 0.04 and -0.04, no better than knowing the population type and season. Conflict alone does most of the work (0.20 and 0.14). The previous survey is still the best single guide to malnutrition (0.40), but for death rates the triggers add to it (0.15 with the previous survey alone, 0.21 with the triggers). For a place never surveyed — predicting a domain from all the others, beyond the season's national average — the phases add nothing for any outcome, the triggers improve the prediction of acute malnutrition modestly (R^2 of 0.10; 0.15 for severe malnutrition), and nothing we have predicts death rates (-0.05 and 0.01).

Within a domain over time, drought is the trigger that matters, and the phases do not respond to it. With domain effects, season type and a trend, a 12-month rainfall deficit of one standard deviation goes with 1.0 more points of acute malnutrition (standard error 0.3), 0.2 more points of




Figure 6: What predicts malnutrition and death in Somalia's surveys? Out-of-sample R^2 from predicting each season's survey results with a model fitted on the other seasons, by information set, for the same 458 survey domains and seasons, 2015–2026 (rural zones, towns and displaced settlements). All models include population type (rural, town, displaced) and season type. Phases: official IPC share of the population in Phase 3 or worse, and FEWS NET's current map and forecast. Triggers: CHIRPS rainfall anomalies over 6 and 12 months, log ACLED conflict fatalities over 6 months, and log cereal and goat prices over 3 months, all up to the survey month. Previous survey: the domain's own outcome, morbidity and measles vaccination coverage in its previous survey. Negative values are shown at the axis.

severe malnutrition (0.1), 0.032 more deaths per 10,000 per day (0.014), and 1.3 more points in the share of children ill in the previous two weeks (0.5), holding the IPC share fixed. The IPC share itself responds to conflict and cereal prices — one standard deviation more conflict raises it by 2.1 points (0.8) — but not to the 12-month rainfall deficit (−0.8 points, standard error 0.7). The phases are not noise: within a domain, a higher IPC share still goes with more severe malnutrition and higher death rates (0.051 deaths per 10,000 per day per standard deviation, 0.017). But the signals the classification weighs are the food signals, prices and conflict, while drought reaches children's bodies partly through illness, which the classification does not count.

Two lessons follow. Cheap triggers carry information about malnutrition that the phases do not, so a system aiming at malnutrition could use them to decide where to look. But they do not substitute for measurement: in places without surveys, no signal we have predicts how many people are dying.

What would surveys be worth? For surveys to save lives, a chain must hold: surveys give better information about where children are malnourished and dying; that information reaches the response; the response moves to where it is needed; and it saves lives there. The first link is strong, as above. The last is strong for treatment: treating severe acute malnutrition in the community averts a death for roughly $4,000–7,000 on GiveWell's inputs for two well-run programmes (GiveWell, 2024, 2025). The middle links are the weak ones. Survey results reach nutrition programmes directly,




Figure 7: If treatment followed surveys: extra deaths averted per season in Somalia when a planner places treatment using each zone's previous scheduled survey rather than FEWS NET's forecast, by the size of the treatment budget: the number of treatment places as a share of all acutely malnourished children in the surveyed zones that season (horizontal axis). 12 surveyed rural zones per season over 26 seasons. The planner allocates treatment places to severe and moderate acute malnutrition by zone to maximize predicted deaths averted; places sent to zones with fewer cases than predicted are stranded, and none, half or all of them can be moved once caseloads are known. Mortality and treatment effects from GiveWell's inputs; caseloads from measured prevalence, population and the standard incidence correction.

since programmes size their caseloads from measured prevalence (Isanaka et al., 2021), but how far donors follow is less clear (Olin et al., 2014), and programmes reach a minority of cases; average coverage across 21 countries was 38 percent (Rogers et al., 2015).

So we ask the counterfactual directly. We model a planner in Somalia who each season has a fixed number of treatment places for malnourished children in the 12 rural zones FSNAU surveys, and spends them to avert as many deaths as possible given what it can see. Severe acute malnutrition, about 16 percent of cases, carries an untreated one-year mortality of about 10 percent, which treatment cuts by about 70 percent; moderate acute malnutrition carries about 4 percent, cut by about 40 percent (GiveWell, 2024). The planner predicts each zone's caseload from the national average, from FEWS NET's forecast phase, or from the zone's previous survey, all out of sample, and sends places where it predicts the most deaths averted. Places sent where cases turn out fewer than predicted are stranded unless they can be moved.

Survey-based planning saves more lives at every realistic budget (Figure 7). With places for 38 percent of the caseload, a planner using FEWS NET's forecast averts about 5,651 deaths a season; one using the previous survey averts 122 to 168 more, depending on how many stranded places can be moved. That is 13–27 percent of the gain perfect foresight would bring. A planner who knows only population does about as well as one using the forecast. Most of the value comes from knowing where severe cases are. A round of surveys in these zones costs about $240,000 (UNICEF

and Action Against Hunger, 2016; Daher et al., 2018), so the extra deaths averted cost about $1,400–1,900 each, and $1,200–2,200 across budgets of 20–50 percent of the caseload. That is in the range of the most cost-effective charities.

This is an upper bound in one sense and a lower bound in another. It assumes treatment follows survey results as well as it could, which is the point of the exercise, not current practice. But it values surveys only for placing treatment within a fixed budget. It gives no credit for sizing the budget, for seeing famine-level conditions directly, for directing health, water and vaccination responses, or for knowing afterwards whether a response worked (Kenny, 2026).

9 Where surveys do not go

Surveys measure outcomes well only where they are run. Somalia's scheduled programme itself has often skipped insecure zones of the south — Juba, South Gedo, parts of Bakool and Hiran — or assessed them only by arm-circumference screening. In FEWS NET's reports, a survey from the previous 12 months is quoted for 67 percent of areas mapped in Emergency in Ethiopia, 54 percent in Somalia and 14 percent in Sudan.

The pattern is general. We assembled metadata for every DHS, MICS and World Bank LSMS household survey in 143 low- and middle-income countries from 1990 to 2023. Across countries, surveys go where poverty is: their frequency falls steadily with income. Within a country over time, they avoid bad years. With country and year effects, a year of minor conflict lowers the chance that a survey starts by 5.5 percentage points (standard error 2.2), and a year of war by 6.8 (3.3), from a base of 16 percent; a fall of more than 2 percent in GDP per capita lowers it by 2.2 (1.2). Around 54 conflict onsets, the chance of a survey drops by about 12 points in the year a conflict starts and recovers within two years. The response comes mostly from the DHS; MICS surveys show no significant response to either conflict or downturns. Using five-year periods, Trommlerová et al. (2026) find conflict a consistent negative predictor of whether a DHS or MICS takes place at all, with economic factors much weaker.

When surveys do go to conflict-affected countries, they often leave out the worst areas. We coded the sampling appendices of 170 DHS and MICS reports from 2000 on, covering every survey run in a country with recent battle deaths and a random sample from calm countries. Some area or selected cluster was dropped for insecurity or inaccessibility in 23 percent of reports from calm countries, 47 percent under minor conflict and 65 percent in wartime (Figure 8). In about one in five conflict-country surveys, whole districts or regions were cut from the design before fieldwork began: the north of Mali in 2012 and 2015, 20 of Borno's 27 local government areas in Nigeria's 2021 MICS, and five districts of Nainawa in Iraq's 2018 MICS, whose report nonetheless says its teams visited every selected area. 92 percent of reports never mention camps; 4 explicitly exclude them and 10 include them, mostly in special designs for refugee populations. Only 6 of the questionnaires we coded identify displaced people.

These are general-purpose household surveys, not the nutrition surveys of Section 8, but the




Figure 8: What household surveys leave out. Share of DHS and MICS final reports (2000 onward) that document dropping an area or selected cluster for insecurity or inaccessibility (red), and that mention camps at all (blue), by battle deaths in the country in the fieldwork year and the two years before. 170 reports with usable sampling text; coded from the reports' sampling appendices.

lesson carries over. A system that relies on surveys will see least where conditions are worst, unless it is designed for those places: scheduled rather than alarm-driven, with methods for insecure areas such as remote and phone-based data collection, community mortality surveillance and sampling of displaced people outside camps.

10 Old debates

Much of this argument is old, and it is worth being clear about which parts.

Starvation or disease. That most famine deaths are disease deaths is de Waal's "health crisis model" (de Waal, 1989b), confirmed by the camp epidemiology of the 1990s (Toole and Waldman, 1990, 1997; Salama et al., 2004) and by economic history (Mokyr and Ó Gráda, 2002; Ó Gráda, 2007). Economists who read famine through Amartya Sen's account of lost entitlements to food (Sen, 1981) should note that this is a different claim: Sen explains why people lose access to food, de Waal why they die. Our argument sides with de Waal. We add only the link to the classification system: if deaths are mostly driven by health, a scale built on food consumption will predict them poorly, and it does.

Malnutrition is not only about food. Since UNICEF's 1990 conceptual framework, nutritionists have agreed that acute malnutrition has two immediate causes, diet and disease, and many underlying ones (UNICEF, 1990, 2020). Helen Young, Susanne Jaspars, Anastasia Marshak and colleagues have argued for three decades that humanitarian practice nonetheless treats high malnutrition as a food problem calling for food (Young and Jaspars, 1995, 2006; Young et al., 2004;

Young and Marshak, 2017). The IPC conceded the point in 2016 by building a separate scale for acute malnutrition, because the two "do not always match" (IPC Nutrition Working Group, 2016). Our contribution is to quantify how weak the match is across all areas we can observe, and to point out that the headline numbers and prioritization still run on the food insecurity scale, not the malnutrition one.

Food first. That needs assessments are shaped to justify food responses is the argument of Darcy and Hofmann (2003), and that food aid was driven by donor surplus and inertia is that of Barrett and Maxwell (2005). Public health researchers have argued since 1990 that the measures that cut mortality in crises are measles vaccination, water and sanitation, treatment of diarrhoea and malaria, and therapeutic feeding, with general rations one item among several (Toole and Waldman, 1990; Médecins Sans Frontières, 1997; Spiegel et al., 2010). Cash has since displaced much in-kind food, but food security is still by far the largest funding line. Earlier critiques blamed agency mandates and donor surpluses. We locate the bias in the information system: the dominant measure of severity is food-based, so prioritization inherits a food-first tilt even when the aid is cash.

Critiques of the IPC. The IPC has been criticized for its politics and governance — negotiated consensus, contested numbers, governments withdrawing — (Maxwell and Hailey, 2020, 2021; Buchanan-Smith et al., 2021, 2023), for inconsistent use of household indicators (Maxwell et al., 2014; Vaitla et al., 2017; Maxwell et al., 2023; Lentz et al., 2024), for undercounting (Lentz et al., 2025), and for famine thresholds that are too high or too insensitive (Young and Jaspars, 2009; Maxwell et al., 2020; de Zwarte et al., 2026). Every one of these critiques takes the IPC's reference table as the standard and asks whether analysts apply it consistently or honestly. The IPC's own accuracy study tests internal consistency, not whether phases predict outcomes (Lentz et al., 2024). Our critique is different: even applied correctly, the scale is only loosely related to the outcomes it is used to stand for. And it concerns the Crisis and Emergency populations where most deaths occur, not famine declarations.

Counting the dead. That crisis mortality data are too sparse, late and unrepresentative is the long-standing argument of Francesco Checchi, Les Roberts and colleagues (Checchi and Roberts, 2005, 2008; Checchi et al., 2017), and recent work in Somalia shows what a mortality database can do (Ratnayake et al., 2025; Palmer et al., 2025). We restate the call for more data and add a value-of-information argument: scheduled surveys are worth paying for as an input to allocating treatment, not only as evidence for classification, advocacy or accountability. Since models built on many covariates have not predicted malnutrition well (Checchi et al., 2022), the case rests on running surveys often, not on cleverer models.

Measure the outcomes, and act on them. Our practical proposal — collect regular survey measures of the final outcomes and direct attention by them — is not new either. Nutritional

surveillance, the regular measurement of malnutrition to guide action, was set out by WHO, FAO and UNICEF in the 1970s and developed in the 1980s (Mason et al., 1984); its use in humanitarian settings has been reviewed by, among others, FSNAU's own former leaders (Bilukha et al., 2012); Darcy and Hofmann (2003) proposed malnutrition and mortality as the common measures of severity across sectors two decades ago; and Somalia's scheduled surveys are an example of the idea in operation. What is new is evidence on the cost of not doing it. The headline measure the system actually uses is only loosely related to the outcomes; cheap triggers and previous surveys carry information the phases miss; and surveys used for allocation would pay for themselves many times over. The proposal has been made; it has not been acted on, and the reason is not that its value was known to be low.

Camps and everywhere else. That the burden of crisis mortality has moved outside camps, and that refugees in camps often fare better than displaced people and residents outside them, is the argument of Salama et al. (2004) and Spiegel et al. (2010), confirmed by Heudtlass et al. (2016): across crises, refugee death rates sat near their host baselines while those of internally displaced people were 2.5 times baseline and residents' 1.5 times. The new part here is twofold. The IPC's death rate bands were calibrated on camp data and are applied to whole rural zones. And the household surveys that could measure deaths elsewhere systematically leave out insecure areas and say little about camps (Section 9).

Early warning. The anticipatory action literature asks whether forecasts of hazards and food insecurity can trigger earlier, cheaper action (Coughlan de Perez et al., 2015; Pople et al., 2026), and post-mortems of Somalia's 2011 famine concluded that early warning was not enough (Hillbruner and Moloney, 2012). We agree with both, and ask a prior question: whether what is forecast is the outcome that matters.

11 Implications

The evidence supports a modest conclusion and suggests a larger one.

The modest conclusion is that FEWS NET is not the problem. It forecasts the IPC's phases, and a companion paper finds that it does so with information others lack (Claude Opus 5.5 et al., 2026). On the same data, FEWS NET and the IPC's Famine Review Committee reached different conclusions about North Gaza in 2024 (IPC Famine Review Committee, 2024a); what binds is the classification's rules, not the forecaster. A decision on whether to keep paying for FEWS NET can rest on what it is designed to do, and it does that reasonably well.

The larger conclusion concerns the IPC and the way its phase counts are used. Four changes follow from the evidence, in increasing order of ambition.

1. **Stop converting phases into deaths.** The reference bands for malnutrition and mortality are not calibrated to phases, and measured gradients are a fraction of them. Estimates of

lives at stake built by multiplying people in Crisis or worse by the bands overstate deaths in some crises and understate them in others. Death rates, where they matter for classification, should be read against local baselines (de Zwarte et al., 2026).

2. **Report outcomes next to phases.** Each IPC analysis could report the latest measured acute malnutrition and mortality for each area, with dates, and flag areas with none. That costs little and makes clear when a phase count rests on no outcome data at all.

3. **Pay for scheduled surveys and surveillance in high-risk areas.** Somalia's scheduled rounds show what a fixed calendar buys: results that predict the next round and are not driven by where alarm happened to be raised. Our estimates put the value of such surveys, if treatment followed them, at about $1,200–2,200 per death averted. They would need to be designed for the places current surveys avoid — insecure districts, displaced people outside camps — using methods built for those settings.

4. **Target downstream outcomes.** If most crisis deaths come from disease, a system that sizes and prioritizes crises by people in food insecurity directs attention to a small part of the problem. A response organized around forecasts of malnutrition and mortality would put more weight on measles vaccination, water and sanitation, malaria and diarrhoea treatment and therapeutic feeding — the measures the public health literature has long found most effective — and could save more lives with the same budget.

The first two are technical changes the IPC partnership could make. The last two would change what the humanitarian system optimizes, and they raise questions we cannot settle here. Mortality is harder and more dangerous to measure than food consumption, especially where conflict is worst. Concentrating on deaths wherever they occur, rather than on the camps and sites that are easiest to reach and measure, is right in principle but would shift resources toward places that are hard to serve. And food insecurity is a harm in itself, not only a route to death; a system that stopped counting it would lose something. These are arguments for testing the larger proposal against the views of practitioners, not for dismissing it.

A Method

For a survey of area a in season t, with measured outcome y_{at} (acute malnutrition, or a crude or under-five death rate) and FEWS NET phase P_{at} (its forecast for the survey month, or its map), we estimate

$$y_{at} = \beta P_{at} + \mu_a + \lambda_t + \varepsilon_{at}.$$

Without the area effects μ_a, β compares different areas: are areas in worse phases worse off? With them, it compares the same area over time: when an area is one phase worse than usual, how much worse are its outcomes? The season effects λ_t remove shocks common to the country in a season. We also score FEWS NET's forecast as a predictor of y_{at} out of sample, leaving out one season at a time, against the simplest alternative: the area's own measurement in the previous round.

Somalia. We kept FSNAU's rural SMART surveys and matched each surveyed zone to FEWS NET's mapping units by livelihood zone and region. Mean acute malnutrition was 13 percent, and 33 percent of zone-seasons were at or above the critical level of 15 percent. FEWS NET's forecast was issued a median of 3 months ahead of the survey. FSNAU's analysts work alongside FEWS NET's, though the survey measurements are independent of the classifications.

Household indicators. Areas are matched to FEWS NET's map for the same month by averaging the phase over the FEWS NET units in the survey area. The IPC thresholds used for the 20 percent rule are those of the IPC reference table: borderline food consumption at Crisis, poor at Emergency or worse, and a reduced coping strategies index of 19 or more at Crisis or worse.

Survey timing. The country-year panel covers 143 countries, 1990–2023. The outcome is whether fieldwork for a DHS (including malaria and AIDS indicator surveys), a MICS or an LSMS started in the year. Conflict is battle deaths from the Uppsala Conflict Data Program, as published in the World Development Indicators (minor conflict: 25–999 deaths; war: 1,000 or more). Linear probability models with country and year effects; standard errors clustered by country. For DHS alone the effects are 4.5 (1.6) points for minor conflict, 3.9 (2.0) for war and 2.2 (0.9) for a downturn, from a base of 7.4 percent.

Survey exclusions. Reports were downloaded from the DHS Program, the World Bank and IHSN microdata catalogs and IPUMS-MICS. Sampling passages were extracted automatically and coded by a language model against a fixed codebook (areas excluded at the design stage, clusters dropped in the field and why, camps excluded or included, displacement questions), with spot checks against the text. Silence is coded as "not mentioned", not as "not excluded".




Figure 9: Within zones: when FEWS NET forecasts a Somali zone to be worse than usual, is its measured malnutrition worse than usual? Both variables are measured relative to the zone's own average and the season's average across zones. Light points: 329 zone-seasons; dark points: averages within 15 equal-sized bins of the forecast. Solid line: fitted slope. Dashed line: the slope implied by the IPC reference bands, roughly the average of the 5- and 10-point steps between band midpoints.




Figure 10: Somalia's scheduled surveys: acute malnutrition in a rural zone this round against the same zone six months earlier, 329 zone-seasons, 2011–2024. Solid line: least-squares fit; dashed line: equality.




Figure 11: The IPC's two scales for the same areas. Rows: the IPC acute food insecurity phase of the area; columns: its acute malnutrition phase for the same period (published by the IPC, or classified from published acute malnutrition prevalence using the IPC's thresholds). Each cell gives the share of the row and the number of areas; outlined cells are where the two scales agree. 1,427 areas in Afghanistan, the Central African Republic, Cameroon, the Democratic Republic of Congo, Mali, Mozambique, Nigeria, Somalia and South Sudan, 2021–2026 (years vary by country).

References

Barrett, C. B. and D. G. Maxwell (2005). *Food Aid After Fifty Years: Recasting Its Role*. London: Routledge.

Bilukha, O., C. Prudhon, G. Moloney, P. Hailey, and D. Doledec (2012). Measuring anthropometric indicators through nutrition surveillance in humanitarian settings: Options, issues, and ways forward. *Food and Nutrition Bulletin 33*(2).

Buchanan-Smith, M., J. Cocking, and Z. Moallin (2023). Independent review of the integrated food security phase classification (IPC) Somalia. Hpg commissioned report, Humanitarian Policy Group, ODI, London.

Buchanan-Smith, M., J. Cocking, and S. Sharp (2021, September). Independent review of the IPC South Sudan. Hpg commissioned report, Humanitarian Policy Group, ODI, London.

Checchi, F. et al. (2022). A method for small-area estimation of population mortality in settings affected by crises. *Population Health Metrics 20*.

Checchi, F., S. Frison, A. Warsame, K. T. Abebe, J. Achen, E. A. Ategbo, M. A. Ayoya, I. Kassim, B. Ndiaye, and M. Nyawo (2022). Can we predict the burden of acute malnutrition in crisis-affected countries? Findings from Somalia and South Sudan. *BMC Nutrition 8*, 92.

Checchi, F., C. I. Jarvis, K. van Zandvoort, and A. Warsame (2023). Mortality among populations affected by armed conflict in northeast Nigeria, 2016 to 2019. *Proceedings of the National Academy of Sciences 120*(30), e2217601120.

Checchi, F. and L. Roberts (2005). Interpreting and using mortality data in humanitarian emergencies: A primer for non-epidemiologists. HPN Network Paper 52, Humanitarian Practice Network, Overseas Development Institute, London.

Checchi, F. and L. Roberts (2008). Documenting mortality in crises: What keeps us from doing better? *PLoS Medicine 5*(7), e146.

Checchi, F. and W. C. Robinson (2013). Mortality among populations of southern and central Somalia affected by severe food insecurity and famine during 2010–2012. Technical report, FAO/FSNAU and FEWS NET.

Checchi, F., A. Warsame, V. Treacy-Wong, J. Polonsky, M. van Ommeren, and C. Prudhon (2017). Public health information in crisis-affected populations: a review of methods and their use for advocacy and action. *The Lancet 390*(10109), 2297–2313.

Claude Opus 5.5, C. Kenny, and J. Sandefur (2026). How much does early warning improve famine response? Working draft, Center for Global Development.

Coughlan de Perez, E., B. van den Hurk, M. K. van Aalst, B. Jongman, T. Klose, and P. Suarez (2015). Forecast-based financing: an approach for catalyzing humanitarian action based on extreme weather and climate forecasts. *Natural Hazards and Earth System Sciences 15*(4), 895–904.

Daher, J., F. Cassard, S. Gari, V. Sauveplane-Stirling, M. Nyawo, and P. Codjia (2018). Implementation of nutrition surveys using SMART methodology in sub-saharan africa. *Field Exchange* (58).

Darcy, J. and C.-A. Hofmann (2003). According to need? Needs assessment and decision-making in the humanitarian sector. HPG Report 15, Overseas Development Institute, Humanitarian Policy Group.

de Waal, A. (1989a). Famine mortality: a case study of Darfur, Sudan 1984–5. *Population Studies 43*(1), 5–24.

de Waal, A. (1989b). *Famine that Kills: Darfur, Sudan, 1984–1985*. Oxford: Clarendon Press. Revised edition, Oxford University Press, 2005.

de Zwarte, I., A. de Waal, and L. H. Lumey (2026). Rethinking current famine classification: insights from history. *The Lancet*. Published 17 February 2026; volume/pages unverified.

Degomme, O. and D. Guha-Sapir (2010). Patterns of mortality rates in Darfur conflict. *The Lancet 375*(9711), 294–300.

FSIN and Global Network Against Food Crises (2026). 2026 global report on food crises. Technical report, FSIN, Rome.

GiveWell (2024, March). Community-based management of acute malnutrition (CMAM).

GiveWell (2025). Taimaka – acute malnutrition treatment in Gombe state, Nigeria (november 2024) and program monitoring (march 2025).

Heudtlass, P., N. Speybroeck, and D. Guha-Sapir (2016). Excess mortality in refugees, internally displaced persons and resident populations in complex humanitarian emergencies (1998–2012) – insights from operational data. *Conflict and Health 10*, 15.

Hillbruner, C. and G. Moloney (2012). When early warning is not enough—lessons learned from the 2011 Somalia famine. *Global Food Security 1*(1), 20–28.

Howe, P. and S. Devereux (2004). Famine intensity and magnitude scales: a proposal for an instrumental definition of famine. *Disasters 28*(4), 353–372.

IPC Famine Review Committee (2022, December). Somalia: Famine review of the IPC acute food insecurity and acute malnutrition analyses (december 2022). Technical report, IPC Global Partners.

IPC Famine Review Committee (2024a, June). FEWS NET gaza analysis: Famine review committee conclusions and recommendations. Technical report, IPC Global Partners.

IPC Famine Review Committee (2024b, December). Sudan: Famine review committee report (december 2024). Technical report, IPC Global Partners.

IPC Global Partners (2019). IPC acute food insecurity reference table for area classification. Reproduced by FAO.

IPC Global Partners (2021). Integrated food security phase classification technical manual version 3.1: Evidence and standards for better food security and nutrition decisions. Technical report, IPC Global Support Unit / FAO, Rome.

IPC Global Partnership (2024, April). Exploring linkages between acute food insecurity and acute malnutrition. Ipc guidance note, IPC Global Support Unit, Rome.

IPC Nutrition Working Group (2016, June). IPC for acute malnutrition: Concepts, tools, and procedures to be used to classify areas based on acute malnutrition. Technical report, IPC Global Support Unit, Rome. Cleared by the IPC Technical Advisory Group and endorsed by the IPC Steering Committee.

Isanaka, S., C. T. Andersen, S. Cousens, M. Myatt, A. Briend, J. Krasevec, C. Hayashi, A. Mayberry, L. Mwirigi, and S. Guerrero (2021). Improving estimates of the burden of severe wasting: analysis of secondary prevalence and incidence data from 352 sites. *BMJ Global Health 6*(3), e004342.

Isanaka, S., N. Nombela, A. Djibo, M. Poupard, D. Van Beckhoven, V. Gaboulaud, P. J. Guerin, and R. F. Grais (2009). Effect of preventive supplementation with ready-to-use therapeutic food on the nutritional status, mortality, and morbidity of children aged 6 to 60 months in Niger: A cluster randomized trial. *JAMA 301*(3), 277–285.

Kenny, C. (2026, September). We don't know the impact of humanitarian aid cuts. We should fix that. Center for Global Development blog, 30 September 2026.

Langendorf, C., T. Roederer, S. de Pee, D. Brown, S. Doyon, A.-A. Mamaty, et al. (2014). Preventing acute malnutrition among young children in crises: a prospective intervention study in Niger. *PLoS Medicine 11*(9), e1001714.

Lentz, E., K. Baylis, and C. Kim (2025). Official estimates of global food insecurity undercount acute hunger. *Nature Food 6*(12).

Lentz, E., K. Baylis, H. Michelson, and C. Kim (2024). IPC accuracy study: Analyzing the internal consistency of the IPC AFI and AMN analyses. Technical report, IPC Global Support Unit.

London School of Hygiene & Tropical Medicine, UNICEF, and WHO (2023, March). From insight to action: Examining mortality in Somalia. Technical report, LSHTM, UNICEF, WHO, Federal Government of Somalia, FAO-FSNAU, SIMAD University.

Luc, G., M. Keita, B. Bazongo, B. Diarra, A. Virchenko, C. Okley, T. Seyvet, Y. Ngongang Mbunang, D. Molla, D. Dujanovic, A. Nur, A. Marshak, A. Rahim, and A. M. Osman (2024). Special focus: Wasting patterns in Somalia. Field Exchange 72, Emergency Nutrition Network.

Machefer, M., M. Ronco, A.-C. Thomas, M. Assouline, M. Rabier, C. Corbane, and F. Rembold (2025). A monthly sub-national harmonized food insecurity dataset for comprehensive analysis and predictive modeling. *Scientific Data 12*, 741. Data: Zenodo 15017473 and the Humanitarian Data Exchange.

Mason, J. B., J.-P. Habicht, H. Tabatabai, and V. Valverde (1984). *Nutritional Surveillance*. Geneva: World Health Organization.

Maxwell, D., G. Adan, P. Hailey, M. S. Day, S. B. J. Odhiambo, L. Kaindi, J. Njiru, A. Venkat, and A. Marshak (2023). Using the household hunger scale to improve analysis and classification of severe food insecurity in famine-risk conditions: Evidence from three countries. *Food Policy 118*, 102449.

Maxwell, D. and P. Hailey (2020). The politics of information and analysis in famines and extreme emergencies: Synthesis of findings from six case studies. Technical report, Feinstein International Center, Tufts University.

Maxwell, D. and P. Hailey (2021). Analysing famine: The politics of information and analysis in food security crises. *Journal of Humanitarian Affairs 3*(1), 16–27.

Maxwell, D., A. Khalif, P. Hailey, and F. Checchi (2020). Viewpoint: Determining famine: Multi-dimensional analysis for the twenty-first century. *Food Policy 92*, 101832.

Maxwell, D., B. Vaitla, and J. Coates (2014). How do indicators of household food insecurity measure up? an empirical comparison from Ethiopia. *Food Policy 47*, 107–116. DOI not verified: 10.1016/j.foodpol.2014.04.003.

Médecins Sans Frontières (1997). *Refugee Health: An Approach to Emergency Situations*. London: Macmillan.

Mokyr, J. and C. Ó Gráda (2002). What do people die of during famines: the Great Irish Famine in comparative perspective. *European Review of Economic History 6*(3), 339–363.

Moradi, S., A. Mirzababaei, H. Mohammadi, et al. (2019). Food insecurity and the risk of undernutrition complications among children and adolescents: A systematic review and meta-analysis. *Nutrition 62*, 52–60.

Ó Gráda, C. (2007). Making famine history. *Journal of Economic Literature 45*(1), 5–38.

Olin, E. et al. (2014). Funding based on needs? A study on the use of needs assessment data by a major humanitarian health assistance donor in its decisions to allocate funds. *PLoS Currents Disasters*.

Palmer, J., M. Alhaffar, B. Garba, J. Hassan, R. Ratnayake, A. Seal, M. Jelle, and F. Checchi (2025). The politics and challenges of measuring mortality for humanitarian decision-making in Somalia, the Central African Republic and Bangladesh. medRxiv preprint.

Pople, A., R. Hill, S. Dercon, and B. Brunckhorst (2026). The importance of being early? Anticipatory cash transfers for flood-affected households. *World Bank Economic Review*. Advance article; earlier CSAE WPS/2021-07.

Ratnayake, R., Y. Ouchtar, Y. A. Ahmed, J. H. Mohamoud, M. Jelle, A. Seal, N. I. Dirie, J. Palmer, and F. Checchi (2025). The Somalia Mortality Estimation Database (S-MED): a bird's eye view of mortality and its determinants. *medRxiv*. Preprint, posted 8 September 2025.

Reusken, M., C. Coffey, F. Cruijssen, B. Melenberg, and C. van Wanrooij (2025). Identification of factors associated with acute malnutrition in children under 5 years and forecasting future prevalence. *BMJ Public Health 3*, e001460.

Rogers, E., M. Myatt, S. Woodhead, S. Guerrero, and J. L. Alvarez (2015). Coverage of community-based management of severe acute malnutrition programmes in twenty-one countries, 2012–2013. *PLoS ONE 10*(6), e0128666.

Salama, P., F. Assefa, L. Talley, P. Spiegel, A. van der Veen, and C. A. Gotway (2001). Malnutrition, measles, mortality, and the humanitarian response during a famine in Ethiopia. *JAMA 286*(5), 563–571.

Salama, P., P. Spiegel, L. Talley, and R. Waldman (2004). Lessons learned from complex emergencies over past decade. *The Lancet 364*(9447), 1801–1813.

Sen, A. (1981). *Poverty and Famines: An Essay on Entitlement and Deprivation*. Oxford: Clarendon Press.

Spiegel, P. B., F. Checchi, S. Colombo, and E. Paik (2010). Health-care needs of people affected by conflict: future trends and changing frameworks. *The Lancet 375*(9711), 341–345.

Toole, M. J. and R. J. Waldman (1990). Prevention of excess mortality in refugee and displaced populations in developing countries. *JAMA 263*(24), 3296–3302.

Toole, M. J. and R. J. Waldman (1997). The public health aspects of complex emergencies and refugee situations. *Annual Review of Public Health 18*, 283–312.

Trommlerová, S. K., M. de Vries Mecheva, and M. Rieger (2026). Disease data and development: Evidence from DHS and MICS surveys. *Journal of Development Studies 62*(10), 1873–1895.

UNICEF (1990). Strategy for improved nutrition of children and women in developing countries. UNICEF Policy Review E/ICEF/1990/L.6, United Nations Children's Fund, New York.

UNICEF (2020). UNICEF conceptual framework on maternal and child nutrition. United Nations Children's Fund, New York.

UNICEF and Action Against Hunger (2016). Implementation of nutrition surveys using SMART methodology in sub-saharan africa: Analytical report. Technical report, UNICEF ESARO/WCARO and Action Against Hunger.

Vaitla, B., J. Coates, L. Glaeser, C. Hillbruner, P. Biswal, and D. Maxwell (2017). The measurement of household food security: Correlation and latent variable analysis of alternative indicators in a large multi-country dataset. *Food Policy 68*, 193–205.

Venkat, A., A. Marshak, H. Young, and E. N. Naumova (2023). Seasonality of acute malnutrition in African drylands: Evidence from 15 years of SMART surveys. *Food and Nutrition Bulletin 44*(2_suppl).

Warsame, A., S. Frison, and F. Checchi (2023). Drought, armed conflict and population mortality in Somalia, 2014–2018: A statistical analysis. *PLOS Global Public Health 3*(4), e0001136.

Young, H., A. Borrel, D. Holland, and P. Salama (2004). Public nutrition in complex emergencies. *The Lancet 364*(9448), 1899–1909.

Young, H. and S. Jaspars (1995). Nutritional assessments, food security and famine. *Disasters 19*(1), 26–36.

Young, H. and S. Jaspars (2006). The meaning and measurement of acute malnutrition in emergencies: A primer for decision-makers. HPN Network Paper 56, Humanitarian Practice Network, Overseas Development Institute, London.

Young, H. and S. Jaspars (2009, September). Review of nutrition and mortality indicators for the integrated food security phase classification (IPC): Reference levels and decision-making. Technical report, SCN Task Force on Assessment, Monitoring and Evaluation; Feinstein International Center; Humanitarian Policy Group, ODI.

Young, H. and A. Marshak (2017). Persistent global acute malnutrition: A discussion paper on the scope of the problem, its drivers, and recommendations for policy, practice, and research. Technical report, Feinstein International Center, Tufts University, Boston.