

Impact of a neurology specialty clinical pharmacist on migraine management outcomes: A prospective observational cohort study

Bethany Franz, PharmD, BCACP; Heather Dalton, PharmD, MS; Megan E. Fleischman, PharmD, BCACP; Kama Thomas, PharmD, BCACP

FROEDTERT & THE MEDICAL COLLEGE OF WISCONSIN | FROEDTERT HOSPITAL | MILWAUKEE, WI



BACKGROUND

- ❖ Chronic migraine affects approximately 2% of the world population¹
- ❖ Migraines present with a significant burden of illness as they impair a patient’s functional ability at school, work and usual daily activities²
- ❖ Calcitonin gene-related peptide (CGRP) antagonists (erenumab, fremanezumab, and galcanezumab) are monoclonal antibodies targeting the calcitonin gene-related peptide ligand or its receptor used in the treatment of chronic migraine prevention³
- ❖ CGRP antagonists are dispensed at Froedtert Specialty Pharmacy due to the high cost, initial monitoring, handling and complex care coordination
- ❖ Neurology specialty pharmacists at Froedtert Health provide comprehensive migraine medication management, handling everything from prior authorizations and financial assistance to drug interaction reviews, administration counseling, coordinated refills and personalized support as part of an integrated care team

>925
Neurology
patients

>650
Migraine
patients

PURPOSE

Evaluate the clinical impact of enhanced pharmacist-patient engagement in a neurology specialty pharmacy program via pharmacist interventions on migraine-related outcomes over six months following CGRP monoclonal antibody initiation

OBJECTIVES

Primary objective:

Analyze the impact of increased pharmacist–patient contact within a neurology specialty pharmacy program by measuring interventions made in the first 6 months of CGRP monoclonal antibody therapy.

Secondary objective:

Evaluate the impact on migraine frequency and usual daily activities from baseline to six months after CGRP monoclonal antibody initiation.

METHODS

Study Design

Prospective, single center, observational cohort study

Inclusion Criteria:

- Patient new to CGRP monoclonal antibodies
- Prescription for one of the following subcutaneous migraine medications: erenumab, fremanezumab and galcanezumab
- Prescribed by a Froedtert Hospital Neurology Clinic Provider
- Migraine ICD-10 code G43.XXX in electronic medical record
- Patient age of 18 years or older

TIMELINE

Study enrollment period:
December 2024 – March 2025

Clinical Outcomes Assessments Completed

Final Outcomes Data Analysis:
September 2025

What is the clinical impact of specialty pharmacist management on migraine patients initiating CGRP monoclonal antibodies?

WORKFLOW OUTLINE

Initial Assessment

Month 3 Assessment

Month 6 Assessment

Figure 1. Clinical Outcomes Assessment Cadence

Headache Migraine

Which follow-up is this patient being assessed?

Radio button Options: [Initial] [First 3 months] [First 6 months] [Annual] [Other]

Average number of migraines per month

0 1-5 5-10 10-15 15-20 >20

Average number of missed days of school or work or usual daily activities in the last month?

0 1-5 5-10 10-15 15-20 >20

Figure 2. Pharmacist Clinical Outcomes Outreach Assessment Tool

REFERENCES

1. May A, Schulte LH. Chronic migraine: risk factors, mechanisms and treatment. *Nat Rev Neurol*. 2016 Aug;12(8):455–64. doi: 10.1038/nrneuro.2016.93. Epub 2016 Jul 8. PMID: 27389092.
2. Ailani, J, Burch, RC, Robbins, MS; the Board of Directors of the American Headache Society. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. *Headache*. 2021; 61: 1021– 1039. <https://doi.org/10.1111/head.14153>
3. Cohen F, Yuan H, Silberstein SD. Calcitonin Gene-Related Peptide (CGRP)-Targeted Monoclonal Antibodies and Antagonists in Migraine: Current Evidence and Rationale. *BioDrugs*. 2022;36(3):341–358. doi:10.1007/s40259-022-00530-0

FUTURE DIRECTION

- ❖ Develop standardized pharmacist intervention protocols
- ❖ Integrate patient-reported outcome measures into routine specialty care
- ❖ Analyze cost-effectiveness of specialty pharmacy interventions
- ❖ Continue to follow beyond 6 months to assess sustained benefits