

POLYADP-RIBOSE POLYMERASE INHIBITOR (PARPI) THERAPY: PATIENT CHARACTERISTICS AND ACCESS

Vivian Truong, PharmD¹ | Brooke Looney, PharmD CSP² | Ryan Moore, MS³ | Autumn D. Zuckerman, PharmD² | Nisha B. Shah, PharmD²

¹Department of Pharmacology, Vanderbilt University, ²Vanderbilt Specialty Pharmacy, Vanderbilt University Medical Center, ³Department of Biostatistics, Vanderbilt University Medical Center

BACKGROUND

- PolyADP-ribose polymerase Inhibitor (PARPi) therapy is used in a variety of cancer indications but is most utilized in ovarian malignancies.¹
- PARPi therapy has been used largely in patients who are female, white, and possess a breast cancer gene (BRCA) mutation.^{2,3}
- High out-of-pocket (OOP) costs for oral cancer treatment lead to delays in starting therapy.⁴
- Integrated specialty pharmacy assistance for OOP costs can reduce financial toxicity.⁵

OBJECTIVE

To analyze disease characteristics, time to medication approval, and out-of-pocket costs for patients receiving PARPi therapy.

METHODS

- DESIGN** Single-center retrospective cohort study
- INCLUSION** Adult patients who initiated PARPi therapy through center's outpatient oncology clinic from November 2017 - December 2018
- EXCLUSION** Patients who did not initiate PARPi therapy
Patients enrolled in a clinical trial

RESULTS

Table 1. Demographics and baseline characteristics (n=15)

	n (%)
Age, years, mean ± SD	63 ± 9
Gender, female	14 (93)
Race	
White	12 (80)
African American	2 (13)
Asian	1 (7)
Body mass index, mean ± SD	31 ± 8
Disease duration, years, mean ± SD	4 ± 4
Total previous chemotherapies, median (range)	3 (1-6)
Previous platinum-based regimens, median (range)	2 (0-3)
ECOG** Performance Status	
0	3 (20)
1	12 (80)
Insurance type	
Commercial	7 (47)
Medicare	6 (40)
Tricare	1 (7)
Uninsured	1 (7)
Alcohol use	
None	9 (60)
Occasional	3 (20)
Daily	1 (7)
2-3 drinks per week	1 (7)
Not documented	1 (7)
Smoking status	
Non-smoker	8 (53)
Previous smoker	6 (40)
Current smoker	1 (7)

*Eastern Cooperative Oncology Group

RESULTS

Figure 1. BRCA, HER2* and hormone-receptor status (n=15)




Table 2. Medication and starting dose (n=15)

	Dose	Cancer type	n (%)
OLAPARIB	300 mg twice daily	Ovarian	7 (47)
		Fallopian tube	2 (13)
		Pancreatic	1 (7)
		Primary peritoneal	1 (7)
		Metastatic squamous cell	1 (7)
OLAPARIB	200 mg twice daily	Ovarian	1 (7)
RUCAPARIB	600 mg twice daily	Ovarian	2 (13)

Figure 4. Patient OOP cost for first fill (n=15)




Figure 2. Sites of metastasis (n=15)




Figure 3. Prior authorization (PA) outcomes



Table 3. Financial assistance (n=15)

	n (%)
No financial assistance	7 (47)
Manufacturer co-pay card	5 (63)
Foundation grant	1 (13)
Manufacturer assistance program	1 (13)
Institution medication assistance program	1 (13)

CONCLUSIONS

- PARPi therapy was predominantly utilized in patients with epithelial ovarian cancer, with olaparib as the most commonly used agent.
- High insurance approval rate and low OOP costs were observed, supporting the role of an integrated specialty pharmacist in reducing financial toxicity.