

Case Study: Successful Prophylactic Use of Coagulation Factor X in an Infant Patient in the Home Environment

Yvonne Lantz; Christine Miller, PharmD; Barbara Prosser, RPh; Timothy Walton, MHS, CCRP
Soleo Health, Frisco, Texas, United States



PURPOSE / OBJECTIVES

Factor X deficiency is a rare bleeding disorder that requires long-term management. The female infant discussed within this case study, Patient A, was diagnosed with severe factor X deficiency two days after birth. Therapy with prothrombin complex concentrate (PCC) was initiated until factor X concentrate was approved by the insurance provider.

After initially receiving twice-weekly factor X concentrate infusions at her local pediatric Hemophilia Treatment Center (HTC) for prophylaxis of bleeding episodes, infusion therapy was transitioned to the home environment at one month old. The purpose of this case study is to highlight the successful long-term, effective use of factor X concentrate in a female infant patient in the home environment

MATERIALS & METHODS

A retrospective review of medical records was performed by a multidisciplinary team from June 9, 2017, through January 23, 2020. The review included customized clinical assessments specific to bleeding disorders that were performed by the specialty infusion pharmacy, as reported through the proprietary clinical outcomes program SoleMetrics®. The review also included progress notes, nursing visit assessments, and hospital and physician office visit notes.

These outcomes measures were collected on 31 different occasions during Patient A’s 30-month period of service, revealing the positive results detailed within this case study prevents adverse events, and promotes positive patient outcomes.

RESULTS

The purpose of this case study is to highlight the successful long-term, effective use of factor X concentrate in a female infant patient in the home environment.

- Clinical monitoring and interventions were key in successful outcomes:
- Patient assessment at each dispense and patient interaction
 - Bleed logs
 - Catheter events
 - Barriers to care/teaching
 - Dosage/weight adjustments
 - Inventory management

The patient was able to successfully receive treatment in her home and travel with her family on several occasions. No adherence issues or interruption of services were experienced during the 30-month period of Patient A’s treatment with factor X concentrate.

Patient monitoring through therapy-specific SoleMetrics assessments were completed at each dispense.

SUMMARY/CONCLUSION

A retrospective review of medical records was performed by a multidisciplinary team from June 9, 2017, through January 23, 2020. As shown within this case study, utilizing customized, therapy-specific assessments and collecting outcomes measures regularly on patients receiving specialty therapies in the home environment supports the correct use of infusion therapy, prevents adverse events, and promotes positive patient outcomes.

Authors of this presentation disclose the following concerning possible financial or personal relationships with commercial entities that may have a direct or indirect interest in the subject matter of this presentation: Nothing to disclose.

SoleMetrics Assessment

Assessment Parameter	Frequency
Caregiver training	Independent after 60 days
Patient weight	Documented prior to fill
Dosage adjustments	7
Catheter events	4
Side effects	None
Hospitalizations/ER visits	None
Bleed documentation/logs	0 Bleeds experienced

Access Events: June 2017 to January 2020

