



# [CS 11] Prac 7r – Continued Fraction III

## Problem Statement

Given an integer  $k$ , consider the following infinite sequence of rational numbers:

$$k, k + \frac{1}{k+1}, k + \frac{1}{(k+1) + \frac{1}{k+2}}, k + \frac{1}{(k+1) + \frac{1}{(k+2) + \frac{1}{k+3}}}, \dots$$

For example, for  $k = 1$ , we get

$$1, 1 + \frac{1}{2}, 1 + \frac{1}{2 + \frac{1}{3}}, 1 + \frac{1}{2 + \frac{1}{3 + \frac{1}{4}}}, \dots$$

which simplify to

$$\frac{1}{1}, \frac{3}{2}, \frac{10}{7}, \frac{43}{30}, \frac{225}{157}, \frac{1393}{972}, \dots$$

For  $k = 2$ , we get

$$2, 2 + \frac{1}{3}, 2 + \frac{1}{3 + \frac{1}{4}}, 2 + \frac{1}{3 + \frac{1}{4 + \frac{1}{5}}}, \dots$$

which simplify to

$$\frac{2}{1}, \frac{7}{3}, \frac{30}{13}, \frac{157}{68}, \frac{972}{421}, \frac{6961}{3015}, \dots$$

Your task is to generate this infinite sequence, as rational numbers.

Submit solution [CS 11]  
Practice 7

Points: 120 (partial)

Time limit: 6.0s

Memory limit: 1G

Author: kvatienza (Kevin Atienza)

Problem type

Allowed languages  
NONE, py3

## Task Details

Your task is to implement a function called `convergents`. This function has a single parameter  $k$ , an `int`.

The function must return a *generator* that generates `Fraction`s, as described in the problem statement.

For this problem, the `Fraction` type can be imported from the `fractions` module.

Note that your generator must be **as lazy as possible**. It should yield each resulting next element as soon as it has enough information, and it should produce these results while advancing the input generators for as little as possible.

## Restrictions

(See 7a for more restrictions)

For this problem:

- Loops and lists are allowed.
- Up to 8 function definitions are allowed.
- Recursion is **disallowed**. (The recursion limit has been greatly reduced.)
- Sets and dictionaries are allowed.
- Generators and comprehensions are allowed.
- The source code limit is 1000.

## Example Calls

`print(*take(6, convergents(1)))` Copy

`1/1 3/2 10/7 43/30 225/157 1393/972` Copy

## Example 2 Function Call

`print(*take(6, convergents(2)))` Copy

`2/1 7/3 30/13 157/68 972/421 6961/3015` Copy

## Example 2 Output

## Constraints

When your program is run:

- The function `convergents` will be called at most 6 times.
- At most 100 elements will be consumed from the returned generator.
- $1 \leq k \leq 100$ .

## Scoring

- You get 120 points if you solve all test cases.

## Clarifications

[Report an issue](#)

No clarifications have been made at this time.