

## Quiz 11 - Asymptotics II

---

Due Date ..... March 4  
Name ..... **Julia Troni**  
Student ID ..... **109280095**

### Contents

<b>1</b>	<b>Instructions</b>	<b>1</b>
<b>2</b>	<b>Standard 13- Asymptotics II</b>	<b>2</b>

### 1 Instructions

- The solutions **should be typed**, using proper mathematical notation. We cannot accept hand-written solutions. Here's a short intro to  $\text{\LaTeX}$ .
- You should submit your work through the **class Canvas page** only. Please submit one PDF file, compiled using this  $\text{\LaTeX}$  template.
- You may not need a full page for your solutions; pagebreaks are there to help Gradescope automatically find where each problem is. Even if you do not attempt every problem, please submit this document with no fewer pages than the blank template (or Gradescope has issues with it).
- You **may not collaborate with other students. Copying from any source is an Honor Code violation. Furthermore, all submissions must be in your own words and reflect your understanding of the material.** If there is any confusion about this policy, it is your responsibility to clarify before the due date.
- Posting to **any** service including, but not limited to Chegg, Discord, Reddit, StackExchange, etc., for help on an assignment is a violation of the Honor Code.

## 2 Standard 13- Asymptotics II

**Problem 1.** Let  $f(n) = (2n)!$  and  $g(n) = 5^n$ . Determine the relationship that best applies:  $f(n) \in O(g(n))$ ,  $f(n) \in \Theta(g(n))$ , or  $f(n) \in \Omega(g(n))$ . Prove your answer. Note that you are expected to spell out **all Calculus details** at the level of Calculus I-II. You may want to use the Sterling's approximation  $n! \sim \left(\frac{n}{e}\right)^n \sqrt{2\pi n}$ , i.e.,  $\lim_{n \rightarrow \infty} \frac{n!}{\left(\frac{n}{e}\right)^n \sqrt{2\pi n}} = 1$ , which implies  $n! = \Theta\left(\left(\frac{n}{e}\right)^n \sqrt{2\pi n}\right)$ .

Apply Limit Comparison Test

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{(2n)!}{5^n}$$

Apply ratio test

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$$

$$\lim_{n \rightarrow \infty} \left[ \frac{\frac{(2(n+1))!}{5^{n+1}}}{\frac{(2n)!}{5^n}} \right]$$

$$\lim_{n \rightarrow \infty} \left( \frac{(2n+2)!}{5^n \cdot 5} \cdot \frac{5^n}{(2n)!} \right)$$

$$\lim_{n \rightarrow \infty} \left( \frac{(2n+2)(2n)! \cdot 5^n}{5^n \cdot 5 \cdot (2n)!} \right)$$

$$\lim_{n \rightarrow \infty} \left( \frac{2n+2}{5} \right) = \infty$$

As the  $L = \infty$  the series diverges. Thus  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$  and so by limit comparison test we have that

$f(n) \in \bigcup (g(n))$ , so we can say

$f(n)$  grows faster than  $g(n)$

Proof.

Another method using Sterling Approximation:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{(2n)!}{5^n} =$$

$$= \lim_{n \rightarrow \infty} \frac{\left(\frac{2n}{e}\right)^{2n} \sqrt{2\pi \cdot 2n}}{5^n} \quad \text{By Sterling's Approximation}$$

$$= \lim_{n \rightarrow \infty} \frac{\left(\frac{4n^2}{e^2}\right)^n \cdot 2\sqrt{\pi} n^{1/2}}{5^n}$$

$$= \lim_{n \rightarrow \infty} \frac{2\sqrt{\pi} \cdot (4n^2)^n \cdot n^{1/2}}{(e^2)^n \cdot 5^n}$$

apply root test

$$= 2\sqrt{\pi} \cdot \lim_{n \rightarrow \infty} \left( \frac{(4n^2)^n \cdot n^{1/2}}{(e^2)^n \cdot 5^n} \right)^{\frac{1}{n}}$$

$$= 2\sqrt{\pi} \cdot \lim_{n \rightarrow \infty} \left( \frac{4n^2 \cdot n^{\frac{1}{2n}}}{5e^2} \right)$$

$$= \frac{8\sqrt{\pi}}{5e^2} \lim_{n \rightarrow \infty} \left( n^2 \cdot n^{\frac{1}{2n}} \right) = \infty$$

As the  $L = \infty$  the series diverges. Thus  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$  and so by Limit comparison test we have that

$f(n) \in \bigcup (g(n))_{n^2}$  so we can say

$f(n)$  grows faster than  $g(n)$

□