

The histogram displays the frequency of days post-injury for 100 subjects. The x-axis, labeled 'Days Post-Injury', ranges from 0 to 60. The y-axis represents the frequency, ranging from 0 to 100. The distribution is highly skewed to the right, with a peak frequency of approximately 100 occurring between 2 and 4 days post-injury. The frequency drops sharply after 4 days, with most subjects reporting an injury within the first 10 days.

| Days Post-Injury (Bin) | Frequency |
|------------------------|-----------|
| 0 - 2 | 10 |
| 2 - 4 | 100 |
| 4 - 6 | 10 |
| 6 - 8 | 10 |
| 8 - 10 | 10 |
| 10 - 12 | 10 |
| 12 - 14 | 10 |
| 14 - 16 | 10 |
| 16 - 18 | 10 |
| 18 - 20 | 10 |
| 20 - 22 | 10 |
| 22 - 24 | 10 |
| 24 - 26 | 10 |
| 26 - 28 | 10 |
| 28 - 30 | 10 |
| 30 - 32 | 10 |
| 32 - 34 | 10 |
| 34 - 36 | 10 |
| 36 - 38 | 10 |
| 38 - 40 | 10 |
| 40 - 42 | 10 |
| 42 - 44 | 10 |
| 44 - 46 | 10 |
| 46 - 48 | 10 |
| 48 - 50 | 10 |
| 50 - 52 | 10 |
| 52 - 54 | 10 |
| 54 - 56 | 10 |
| 56 - 58 | 10 |
| 58 - 60 | 10 |

