Tetraplegia Female Male

-		19% (n	= 15)		2019 _		81% (n =	= 63)
-		23% (n =	= 35)		2018 _	77	% (n = 1	16)
-		24% (n =	= 56)		2017 _	76	% (n = 1	82)
-		21% (n	= 48)		2016 _	79	9% (n = 1	179)
-		21% (n	= 33)		2015 _	79	9% (n = 1	126)
-		26% (n =	39)		2014 _	74%	% (n = 11	1)
-		17% (ı	n = 28)		2013 _	3	33% (n =	133)
-		19% (n	= 26)		2012 _	8	1% (n =	110)
-		18% (r	n = 26)		2011 _	8	2% (n =	115)
-		26% (n =	32)		2010 _	74	1% (n = 9	92)
-		21% (n	= 30)		2009 _	79	9% (n = 1	10)
-		17% (n = 20)		2008 _		33% (n =	101)
-		18% (r	1 = 28		2007	8	32% (n =	128)
-		27% (n =	43)		2006 _	73%	% (n = 11	7)
-		23% (n :	= 32)		2005 _	77	% (n = 1	09)
-		29% (n = 2)	27)		2004 _	719	% (n = 66)	3)
-		23% (n =	= 11)		2003 _	7	7% (n =	37)
-		28% (n =	11)		2002 _	72	% (n = 2	9)
-		27% (n =			2001 _	7	'3% (n =	8)
100	75	50	25	0	() 25	50	75