

J. Andrés^{1,2}, I. Marzáñ¹, P. Ayarza², D. Martí¹, I. Palomeras², R. Carbonell¹

¹Institute of Earth Science Jaume Almera, ICTJA-CSIC, Barcelona, Spain. ²Department of Geology, University of Salamanca, Salamanca, Spain.

1 Introduction

Thermal conductivity (k) determine the rate of heat transport by conduction from the lithosphere to the surface. Heat flow (HF) in the surface is related to the thermal gradient (grdT) and thermal conductivity in steady-state by the Fourier Law,

$$q = -k \cdot \text{grdT}$$


where, q represents the surface heat flow, k is the thermal conductivity and grdT is the thermal gradient.

When modelling thermal data, k is usually given as a fixed value in the range of $(2.5 \text{ to } 5 \text{ W m}^{-1} \text{ K}^{-1})$, where $2.5 \text{ W m}^{-1} \text{ K}^{-1}$ and $3.2 \text{ W m}^{-1} \text{ K}^{-1}$ are the typical values assigned to the crust and mantle respectively. Measurements of k are restricted to shallow layers of the crust, are discrete, and often do not illustrate correctly its depth and lateral distribution.

Can we estimate the thermal conductivity of the Iberian Peninsula?

2 Data

Data used in this study is calculated in accordance to the depth of the Curie isotherm (CDP) (Andrés et al., 2017). Thus, the data is referred to a layer of heterogeneous thickness. For onshore areas the CDP is always above the Moho. For offshore areas, it is almost always below the Moho, so upper mantle is included.



Bibliography

Andrés, J., Marzáñ, I., Ayarza, P., Martí, D., Palomeras, I., Campbell, S., Carbonell, R. (2017), Curie Depth Point of the Iberian Microplate. A thermal, compositional and tectonic perspective of its evolution. *Earth and Planetary Science Letters*. Submitted.

Palomeras, I., Villaseñor, A., Thurner, S., Lavender, A., Gallart, J. and Harratt, M. (2017), Lithospheric structure of Iberia and Morocco using finite-frequency Rayleigh wave tomography from earthquakes and seismic ambient noise. *Geochim. Geophys. Geosyst.*. Accepted Author Manuscript. doi:10.1002/2016GC006567

Fraley, C., and Raftery, A. E. (2002), Model-based Clustering, Discriminant Analysis and Density Estimation. *Journal of the American Statistical Association* 97:611-631.

3 Methodology

The proposed methodology (Fig 1) is based on determining a k map from the correlation between V_s map and grdT map determined by calculating the depth of the Curie isotherm for the Iberian Peninsula.

- The data is cross-plotted (Fig. 3), and a statistical classification method based on a non-supervised algorithm, the expectation-maximization (EM) algorithm is used to classify populations within the cross-plot (Fraley et al. 2002) (Fig. 3). The linear regressions of each population is the relationship between thermal gradient and thermal conductivity.
- Calculate k assigning a known mean HF value for each area
- Calculate the HF from the calculated k and grdT
- Calculate differences between calculated HF and observed HF and iterate until the minimum misfit is obtained.




Figure 3. Cross-plot of V_s and grdT with colored populations and linear regressions overlapped.

Methodology





Figure 4. Map of distribution of clusters. Geographically, the populations correspond with different tectonic domains in Iberia. 1. Variscan domain, 2. Alpine domain, 3. Mediterranean, 4. Gibraltar Arc.

4 Results

- Differentiation between the Variscan domain, values above 2.8 W/mK with maximums up to 3.2 W/mK , and the Alpine domain, mean value of 2.6 W/mK (Fig. 5).
- Offshore areas have values up to 3.6 W/mK with mean value of 3 W/mK .
- Gibraltar Arc region presents very low values between 1.4 W/mK and 2.4 W/mK , with an average thermal conductivity of 2 W/mK .
- Good fit between calculated and observed HF offshore and onshore despite the poor coverage inland (Fig. 6).
- Gibraltar Arc abrupt change is captured




Figure 5. Mean thermal conductivity calculated.




Figure 6. Calculated HF with overlapped observed HF.




Figure 7. Differences between observed and calculated HF. a) Histogram of the differences, b) plot.

5 Conclusions and future work

- Differentiation between different tectonic domains.
- Offshore areas have values up to 3.6 W/mK with mean value of 3 W/mK .

Annowledgements

We acknowledge research support from projects CGL2014-56548-P, 2009-SGR-1595, CGL2013-47412-C2-1-P

Get this poster!