


a

	Top-1 accuracy			
Baseline	0.689	0.682	0.527	0.724
HostMix-TOO	0.728	0.756	0.698	0.455
Site-specific mixtures	0.714	0.758	0.690	0.435
Normal classes	0.675	0.709	0.487	0.397
Linear deconvolution	0.709	0.723	0.536	0.812
Native-organ masking	0.600	0.637	0.347	0.209
Gene removal 20%	0.677	n/c	n/c	n/c
Site-specific + removal 20%	0.709	n/c	n/c	n/c
Site-specific + removal 40%	0.705	n/c	n/c	n/c
Gene sets	0.568	0.596	0.302	0.060
Gene sets + mixtures	0.670	0.727	0.505	0.229
Gene sets + host correction	0.579	0.678	0.295	0.013
Gated model	0.735	0.775	0.550	0.683
22-tissue model	0.705	0.746	0.534	0.539
Perceptron + mixtures	0.696	0.750	0.520	0.535
	MET500 (n = 437)	POG570 (n = 512)	Auxiliary RNA-seq (n = 729)	Microarray (n = 536)



b

	Host-attraction rate			
	0.136	0.228	0.204	0.256
	0.044	0.024	0.056	0.101
	0.030	0.024	0.049	0.111
	0.119	0.180	0.171	0.203
	0.089	0.143	0.145	0.010
	0.000	0.000	0.000	0.000
	0.119	n/c	n/c	n/c
	0.030	n/c	n/c	n/c
	0.028	n/c	n/c	n/c
	0.141	0.246	0.152	0.048
	0.044	0.037	0.061	0.087
	0.053	0.074	0.094	0.005
	0.053	0.042	0.061	0.097
	0.047	0.048	0.056	0.188
	0.030	0.024	0.044	0.053
	MET500 (n = 361)	POG570 (n = 378)	Auxiliary RNA-seq (n = 427)	Microarray (n = 207)

