
a
muscle WT, 6 vs 6
kidney WT, 6 vs 6
kidney 16/23
muscle 19/23
KO − WT (Δlog10 hazard units)
1.00
0.75
0.50
0.25
0.00
-0.25
black
blue
brown
cyan
darkgreen
darkgrey
darkorange
darkred
green
greenyellow
grey60
lightgreen
lightyellow
magenta
midnightblue
orange
pink
red
salmon
skyblue
tan
turquoise
yellow
b
80% power |ρ|, α = 0.05
adjusted Spearman ρ
0.4
0.2
0.0
-0.2
-0.4
ATI
ATII
B
Ciliated
DC
Endothelial
Macrophage
Monocyte
NK
Stromal
T
FDR < 0.10
composite 0.21
adjusted Spearman ρ
0.4
0.2
0.0
-0.2
0
10
20
modules ordered
by adjusted ρ
c
recovery of expected shift (%)
100
80
60
40
20
0
-20
whole lung
detected genes
expression only
composition only
AT2
Alv. mac
Ciliated
d
sum of contributions: mouse 0.72, HLCA 1.00
other
ATI
Airway epi
Ciliated
DC
ATII
Stromal
T
Macrophage
other
vascular endothelial cells
Mesothelial cells
Interstitial Fibroblast
Club cells
Interstitial macrophages
Type 2 pneumocytes
Alveolar macrophage
Ciliated cells
mouse lung
HLCA
-0.4
-0.2
0.0
0.2
0.4
0.6
0.8
1.0
contribution (fraction of the expression-only effect)