

# Contents

<b>1</b>	<b>Introduction and theory overview</b>	<b>1</b>
1.1	Introduction . . . . .	1
1.2	Theory overview . . . . .	2
1.2.1	Heavy Vector Triplet model . . . . .	2

# List of Figures

# List of Tables

# Chapter 1

## Introduction and theory overview

### 1.1 Introduction

This thesis presents the analysis details and the results of the search for heavy resonances decaying into a Z boson and a Higgs boson (h) at the center-of-mass energy of 8 TeV, using  $19.7 \text{ fb}^{-1}$  p-p collision data. In turn, the Z boson is identified through its leptonic decays (Leptons often refer to  $e$  and  $\mu$  only in experiments.  $l = e, \mu$ ). The Higgs boson h is expected to hadronically decay into a pair of b-quarks. The investigated final states consist of two charged leptons which are identified in the detector and limit the presence of the background, and two b-quarks from the hadronic Higgs decay which collects the largest possible fraction of Higgs events.

This thesis is organised as follows. In the latter part of this chapter, the heavy resonances model is introduced, including the expected cross section and the specification of model parameters. In chapter 2, the LHC and the CMS experiment are introduced, including the information of each sub-detector and the trigger system of the CMS. The details of the analysis are shown in chapter 3. This chapter reveals the way to reconstruct physical objects in CMS. By adding some proper kinematic selections on those physics objects, the interested event in data collected by the CMS detector can be selected. Moreover, this chapter shows the comparison between data and simulation. In the last chapter, the results of the search and the conclusion are shown.

## 1.2 Theory overview

Although the Higgs boson discovery by the ATLAS and CMS collaborations [1–3] imposes strong constraints on theories beyond the Standard Model, the extreme fine tuning in quantum corrections required to have a light fundamental Higgs boson with mass close to 125 GeV [4–7] suggests that the Standard Model may be incomplete, and not valid beyond a scale of a few TeV. Various dynamical electroweak symmetry breaking scenarios which attempt to solve this naturalness problem, such as Minimal Walking Technicolor [8], Little Higgs [9–11], or composite Higgs models [12–14] predict the existence of new resonances decaying to a vector boson plus a Higgs boson.

### 1.2.1 Heavy Vector Triplet model

We introduce a model-independent strategy which applies a heavy vector triplet [15] of the Standard Model group to study the new resonances. It will build a simplified phenomenological Lagrangian which reproduces a large class of explicit models.

# Bibliography

- [1] ATLAS Collaboration. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phy. Lett. B*, 716:1, 2012.
- [2] CMS Collaboration. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phy. Lett. B*, 716:30, 2012.
- [3] CMS Collaboration. Observation of a new boson with mass near 125 GeV in pp collisions at  $\sqrt{s} = 7$  and 8 TeV. *JHEP*, 06:081, 2013.
- [4] CMS Collaboration. Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV. *Eur. Phys. J. C*, 75:212, 2015.
- [5] ATLAS Collaboration. Measurement of the Higgs boson mass from the  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4l$  channels in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector. *Phys. Rev. D*, 90:052004, 2014.
- [6] ATLAS Collaboration. Evidence for the spin-0 nature of the Higgs boson using ATLAS data. *Phys. Lett. B*, 726:120, 2013.
- [7] CMS and ATLAS Collaboration. Combined Measurement of the Higgs Boson Mass in pp Collisions at  $\sqrt{s} = 7$  and 8 TeV with the ATLAS and CMS Experiments. *Phys. Rev. Lett.*, 114:191803, 2015.
- [8] Mads T. Frandsen. Minimal Walking Technicolor. 2007.
- [9] B. McElrath T. Han, H. E. Logan and L.-T. Wang. Phenomenology of the little Higgs model. *Phys. Rev. D*, 67:095004, 2003.
- [10] M. Schmaltz and D. Tucker-Smith. LITTLE HIGGS THEORIES. *Annual Review of Nuclear and Particle Science*, 55:no. 1, 229–270, 2005.
- [11] M. Perelstein. Little Higgs models and their phenomenolog. *Progress in Particle and Nuclear Physics*, 58:no. 1, 247–291, 2007.
- [12] D. Marzocca R. Contino, D. Pappadopulo and R. Rattazzi. On the effect of resonances in composite Higgs phenomenology. *Journal of High Energy Physic*, 2011:no. 10, 1–50, 2011.

- [13] M. Serone D. Marzocca and J. Shu. General composite Higgs model. *Journal of High Energy Physics*, 2012:no. 8, 1–52, 2012.
- [14] C. Cski B. Bellazzini and J. Serra. Composite Higgses. *Eur. Phys. J.*, C74:no. 5, 2766, 2014.
- [15] R. Torre D. Pappadopulo, A. Thamm and A. Wulzer. Heavy vector triplets: bridging theory and data. *Journal of High Energy Physics*, 2014:no. 9, 1–50, 2014.