

# Use case: leukemia signatures (Den Boer, 2009)

Analyse Statistique des données Génomiques (ASG)

Jacques van Helden

2020-03-06



# Identification of signatures for subtypes of leukemia




Figure 1: Identification of a gene-expression signature enabling classification of paediatric ALL

**Figure 1:** Figure 1 from Den Boer et al. (2009).

# Hierarchical clustering of genes and samples





Figure 2: Clustering of ALL subtypes by gene-expression profiles

Hierarchical clustering of patients from the COALL (left) and DCOG (right) studies with 110 gene-probe sets selected to classify paediatric ALL. Heat map shows which gene-probe sets are overexpressed (in red) and which gene probe sets are underexpressed (in green) relative to mean expression of all gene-probe sets (see scale bar).

\*Patients with E2A-rearranged subclone (15–26% positive cells). †Right column of grey bar denotes BCR-ABL1-like cases.

**Figure 2:** Figure 2 from Den Boer et al. (2009).

# Survival curves



**Figure 3: Kaplan-Meier estimates for the probability of disease-free survival (pDFS) in children with precursor B-ALL**

The COALL precursor B-ALL cohort (left) comprised 145 patients from COALL-92/97 and nine from DCOG ALL-9 treated at Sophia Children's Hospital. As reference, data for 22 patients with BCR-ABL1-positive disease enrolled in the COALL-92/97 protocol were included. Univariate analysis of pDFS comparing 30 BCR-ABL1-like with 119 remaining precursor B-ALL cases (excluding five BCR-ABL1-positives):  $p=0.012$ . Curves were almost the same when the nine DCOG patients were excluded. The DCOG precursor B-ALL cohort (right) comprised 92 children from DCOG-ALL8. As reference, data for 25 BCR-ABL1-positive cases enrolled in DCOG-ALL-7, 8, and 9 were included. Univariate analysis of pDFS comparing 14 BCR-ABL1-like and 77 remaining precursor B-ALL cases (excluding BCR-ABL1-positive case):  $p=0.026$ .

**Figure 3: Figure 3 from Den Boer et al. (2009).**