

Compactness

Definition 1. (Compact) Let (X, d) be a metric space and let $K \subseteq X$. K is said to be compact if every open cover of K has a finite subcover. That is, if $\{O_\alpha\}_{\alpha \in \Lambda}$ is any open cover of K , then

$$\exists \alpha_1, \dots, \alpha_n \text{ such that } K \subseteq O_{\alpha_1} \cup \dots \cup O_{\alpha_n}$$

Example. Let (X, d) be a metric space and let $E \subseteq X$.
If E is finite, then E is compact.

The reason is as follows:

Let $\{O_\alpha\}_{\alpha \in \Lambda}$ be any open cover of E . Our goal is to show that this open cover has a finite subcover.

If $E = \emptyset$, there is nothing to prove.

If $E \neq \emptyset$, denote the elements of E by x_1, \dots, x_n :

$$E = \{x_1, \dots, x_n\}$$

. We have:

$$x_1 \in E \subseteq \bigcup_{\alpha \in \Lambda} O_\alpha \implies \exists \alpha_1 \in \Lambda \text{ such that } x_1 \in O_{\alpha_1}$$

$$x_2 \in E \subseteq \bigcup_{\alpha \in \Lambda} O_\alpha \implies \exists \alpha_2 \in \Lambda \text{ such that } x_2 \in O_{\alpha_2}$$

\vdots

$$x_n \in E \subseteq \bigcup_{\alpha \in \Lambda} O_\alpha \implies \exists \alpha_n \in \Lambda \text{ such that } x_n \in O_{\alpha_n}$$

Hence,

$$E = \{x_1, \dots, x_n\} \subseteq O_{\alpha_1} \cup \dots \cup O_{\alpha_n}$$

So, $O_{\alpha_1}, \dots, O_{\alpha_n}$ is a finite subcover of E .

Q.E.D.

Example. Consider $(\mathbb{R}, ||)$ and let $E = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$.

Prove that E is compact. (In general, if $a_n \rightarrow a$ in \mathbb{R} then $F = \{a_n : n \in \mathbb{N}\} \cup \{a\}$ is compact.)

Proof. Let $\{O_\alpha\}_{\alpha \in \Lambda}$ be any open cover of E . Our goal is to show that this open cover has a finite subcover.

$$\left. \begin{array}{l} 0 \in E \\ E \subseteq \bigcup_{\alpha \in \Lambda} O_\alpha \end{array} \right\} \implies 0 \in \bigcup_{\alpha \in \Lambda} O_\alpha \implies \exists \alpha_0 \in \Lambda \text{ such that } 0 \in O_{\alpha_0} \quad (\text{I})$$

$$\left. \begin{array}{l} 0 \in O_{\alpha_0} \\ O_{\alpha_0} \text{ is open} \end{array} \right\} \implies \exists \epsilon > 0 \text{ such that } (-\epsilon, \epsilon) \subseteq O_{\alpha_0}$$

By the archimedean property of \mathbb{R} ,

$$\exists m \in \mathbb{N} \text{ such that } \frac{1}{m} < \epsilon$$

so

$$\forall n \geq m \quad \frac{1}{n} < \epsilon.$$

Hence

$$\forall n \geq m \quad \frac{1}{n} \in (-\epsilon, \epsilon) \subseteq O_{\alpha_0} \quad (\text{II})$$