

FOUR PROBLEMS FROM ERICH FRIEDMAN’S *MATH MAGIC*

VAMSHI JANDHYALA

ABSTRACT. Erich Friedman’s *Math Magic* pages pose a problem each month and keep a list of the questions that remained open. We settle four of them. Two exact values for armies of bishops, $B(3, 5) = 5$ and $B(3, 6) = 8$, follow from a one-line lemma and an exhaustive count. Three colours of king with touch counts $(1, 3, 6)$, $(1, 6, 3)$ or $(2, 3, 4)$ admit no finite arrangement (unsolved problem 24), by a reduction to finite formulas with checkable refutations. Three of the 6-point capture digraphs of unsolved problem 21 are not realised by any chess position. And one of the heptominoes of unsolved problem 15 can never be arranged in a square with equal row and column counts, by an explicit weighting. Every computation is reproduced by programs published with the paper.

1. INTRODUCTION

Since 1998 Erich Friedman’s *Math Magic* has posed one problem a month, published the solutions readers sent in, and collected the questions they could not answer on a list of unsolved problems [1]. The problems are recreational in flavour and often easy to state, which makes the open ones tantalising. This note settles four of them. As of 9 October 2026 the list still shows problems 15, 21 and 24 as open, and the March 2005 page on armies of bishops still asks for the two values we prove [2].

The results are of two kinds, and we label them. A *proved* statement has a proof a reader can follow by hand. A *computer-certified* statement rests on a finite computation: an exhaustive count, or a solver’s refutation of a finite formula. For each certified statement we say what was computed, why the finite computation implies the statement, and how it was checked independently. The programs are in the `verify` folder that accompanies this paper; each section names the subfolder.

Problem	Source	Result	Kind
Armies of bishops	March 2005	$B(3, 5) = 5$, $B(3, 6) = 8$	computer-certified
Touch cycles (no. 24)	March 2015	three triples impossible	computer-certified
Capture digraphs (no. 21)	October 2013	diagrams q , r , s impossible	computer-certified
Magic polyomino squares (no. 15)	May 2007	heptomino A impossible	proved

2. ARMIES OF BISHOPS

Place C armies of k bishops each on an $N \times N$ board so that no bishop attacks a bishop of another army. A bishop attacks along diagonals and is blocked by the first piece in its way, of either army. Let $B(C, N)$ be the largest such k . Friedman’s table [2] gives lower bounds for small C and N , and asks whether $B(3, 5) = 5$ and $B(3, 6) = 8$. Figure 1 shows three armies of eight on the 6×6 board. The question is whether nine is possible.

The figure suggests the right way to look at a placement: by diagonals, not by bishops.

Lemma 1 (proved). *A placement of armies is valid if and only if every diagonal containing bishops contains bishops of only one army.*

Date: Preprint, 9 October 2026. Not peer reviewed.




FIGURE 1. Three armies of eight bishops (black, white, grey) on the 6×6 board. Every diagonal that carries bishops carries only one army.

Proof. If a diagonal carries bishops of two armies, walk along it: somewhere a bishop of one army is followed, with nothing in between, by a bishop of the other, and these two attack each other. Conversely, an attack always runs along a diagonal between two bishops on it, so if every diagonal carries one army no bishop attacks another army. $\square$

So blocking plays no part, and a placement is determined, up to which cells are used, by a label on each diagonal: the army it carries, or “unused”. A bishop of army i can stand on a cell exactly when both diagonals through the cell carry label i . Taking every such cell gives a valid placement by Lemma 1, and removing bishops keeps a placement valid. Write $u_i(\lambda)$ for the number of cells whose two diagonals both carry label i in the labelling λ . Hence

$$B(C, N) = \max_{\lambda} \min_{1 \leq i \leq C} u_i(\lambda).$$

The right side is the size of the smallest army when every usable cell is filled, maximised over all ways of labelling the diagonals. A light square and a dark square never share a diagonal, so the light and dark diagonals can be labelled separately: for each colour we list the vectors of army sizes that its labellings achieve, then combine one vector from each colour.

Theorem 2 (computer-certified). $B(3, 5) = 5$ and $B(3, 6) = 8$.

Method. The lower bounds are the placements of Figure 1 and of its 5×5 analogue in `verify/bishops/positions.json`, checked by a move-by-move attack simulator. For the upper bounds, with $C = 3$ each diagonal takes one of four labels. On the 5×5 board the light squares lie on ten diagonals and the dark squares on eight, giving $4^{10} + 4^8$ labellings; on the 6×6 board each colour has eleven, giving $2 \cdot 4^{11}$. The program `verify/bishops/check.py` lists them all and evaluates the formula above. The maxima are 5 and 8. The same program reproduces $B(3, 3) = 2$ and $B(3, 4) = 4$, and with two armies the values $B(2, 5) = 12$ and $B(2, 6) = 18$ in Friedman’s table. In `mutants.py` a recoloured bishop is caught by the simulator, and a mutant that lets a bishop stand on a cell when only one of its diagonals matches its army gives a larger value for $B(3, 5)$, so the count does depend on the lemma. $\square$

3. TOUCH CYCLES (UNSOLVED PROBLEM 24)

Place kings of three colours A , B , C on the cells of the plane, at most one per cell. Two cells touch when they are king moves apart, that is, when their coordinates differ by at most one in each direction. Friedman [5] asks for which triples (a, b, c) there is a nonempty finite arrangement in which every A touches exactly a kings of colour B , every B touches exactly b of colour C , and every C touches exactly c of colour A . Figure 2 shows one for $(1, 1, 6)$. The unsolved list gives $(1, 3, 6)$, $(1, 6, 3)$ and $(2, 3, 4)$ as unknown and conjectures that they do not exist.




FIGURE 2. A finite arrangement for $(a, b, c) = (1, 1, 6)$: every A (white) touches one B (black), every B touches one C (grey), and every C touches six A 's.

Theorem 3 (computer-certified). *There is no nonempty finite arrangement for $(a, b, c) = (1, 3, 6)$, $(1, 6, 3)$ or $(2, 3, 4)$. Nor is there one for $(1, 4, 5)$.*

Method. The idea is to look at an extreme king, where the arrangement must stop. Fix a direction n , namely $n = (0, 1)$ for the first, second and fourth triples and $n = (1, 1)$ for $(2, 3, 4)$. Given a finite arrangement, choose the occupied cell p that maximises $n \cdot p$, breaking ties by the larger first coordinate, and translate so that p is the origin. Then every cell q with $n \cdot q > 0$, or with $n \cdot q = 0$ and positive first coordinate, is empty.

Now restrict attention to the window $[-r, r]^2$. Every cell strictly inside it, with both coordinates of absolute value less than r , has all eight neighbours in the window, so the counting rule at that cell refers only to cells of the window. The restriction of the arrangement is therefore a solution of a finite problem: colour each window cell A , B , C or empty, with the origin occupied, the half-plane beyond it empty, and the counting rule imposed at every occupied interior cell. That problem is unsatisfiable for $r = 6, 6, 8$ and 4 for the four triples in the order listed. So no finite arrangement exists.

Each finite problem is written as a formula in conjunctive normal form, and the folder `verify/touch-cycles/certificates` holds the formula with a DRUP refutation found by a SAT solver. The checker `verify_drup.py`, which uses only the Python standard library, confirms all four refutations in about thirteen seconds. Trust then moves to the encoding, and that was checked separately: `touch_extreme.py` is an independently written encoding of the same window problem for the Z3 solver. It reports the four problems unsatisfiable at the same radii, and satisfiable for the controls $(1, 2, 3)$ and $(2, 2, 2)$, which do have finite arrangements. $\square$

For $(1, 4, 5)$, which is not on the list, there is also a proof by hand (`HUMAN-PROOF-145.md`). For the three listed triples finiteness is essential: each has a doubly periodic arrangement on the whole plane, with periods 6×6 , 8×8 and 6×6 (`touch_torus.py`). A proof of Theorem 3 must therefore use the boundary of the arrangement, as the extreme-king argument does.

4. CAPTURE DIGRAPHS (UNSOLVED PROBLEM 21)

Put chess pieces (kings, queens, rooks, bishops and knights, any number of each, colours ignored) on a board, and draw an arrow from piece X to piece Y when X attacks Y ; sliding pieces are blocked by the first piece in their way. Friedman [4] asks which regular digraphs arise this way. For six vertices with two arrows in and two out at every vertex there are 23 digraphs up to isomorphism, which his page labels a to w . Readers found positions for eighteen of them, DeVincentis showed that f cannot be realised, and the author found a position for u on a 3×4 board (Figure 3, left). The diagrams q , r and s remained open, and the unsolved list conjectures that they are not capture graphs.




FIGURE 3. Left: a position on a 3×4 board whose capture graph is the 2-regular digraph u ; each arrow is an attack. Right: the digraph q , which no position realises.

Theorem 4 (computer-certified). *The digraphs q , r and s are not the capture graph of any position, on a board of any size.*

Method. A position on any finite board is also a position on the infinite board $\mathbb{Z}^2$ with the same attacks, so it is enough to rule out $\mathbb{Z}^2$. The program `src/digraphs.py` lists the 2-regular digraphs on six vertices up to isomorphism and finds 23, matching Friedman’s count. To tie the solver’s numbering to his letters without reading arrows off pictures, the eighteen published positions were transcribed (`out/published_solutions.txt`) and their capture graphs computed; they cover eighteen distinct classes, and the five left over are exactly the five diagrams f , q , r , s , u without a published position.

Two models then decide every class. The first (`src/realize.py`) gives each piece unbounded integer coordinates and asks Z3, in linear integer arithmetic, for a position with exactly the required arrows; it enumerates the assignments of piece types to vertices, discarding only assignments that cannot work for a stated reason (two pieces of the same type attack each other symmetrically, for example). The second (`verify/second.py`) is written independently for the cvc5 solver, keeps the piece types as solver variables, and does no pruning. Both models find positions for 19 classes and prove the other four unrealisable: f , q , r and s . Every position the solvers return is re-checked by a square-by-square simulator (`verify/check.py`), and the encodings are tested against that simulator on random positions, with deliberate mutants caught (`verify/test_*.py`). $\square$

The result for f agrees with DeVincentis. A proof of Theorem 4 by hand would still be welcome: after the pruning there are only 240, 60 and 60 type assignments to rule out for q , r and s .

5. A HEPTOMINO THAT NEVER BALANCES (UNSOLVED PROBLEM 15)

Friedman’s May 2007 problem [3] asks how few copies of a given polyomino can be placed, without overlapping, in a square so that every row and every column contains the same positive number of covered cells. Rotations and reflections are allowed. For most small polyominoes some arrangement works. Unsolved problem 15 lists heptominoes for which none is known and conjectures that none exists. We settle the first of them, the shape A of Figure 4.

Theorem 5 (proved). *No arrangement of one or more copies of A in an $n \times n$ square has the same positive number of covered cells in every row and every column.*

Proof. Give each column x a weight w_x and each row y the same weight w_y , and give the cell (x, y) the weight $w_x + w_y$. The weight of a set of cells is then the sum, over columns, of the




FIGURE 4. The heptomino A with its row counts $P = (2, 1, 1, 1, 2)$ and column counts $Q = (1, 5, 1)$.

column's weight times the number of cells in it, plus the same sum over rows. For a balanced arrangement with c covered cells in every row and column this is $2c \sum_x w_x$.

Now weigh a single copy. The row counts of A are $P = (2, 1, 1, 1, 2)$ and its column counts are $Q = (1, 5, 1)$. Both are palindromes, so in every rotation and reflection one direction sees P across five consecutive lines and the other sees Q across three. A copy therefore weighs $P \cdot (w_i, \dots, w_{i+4}) + Q \cdot (w_j, \dots, w_{j+2})$ for some i and j . Call these the P -windows and Q -windows of w . If every P -window is at least -1 and every Q -window is at least 1 , each copy weighs at least 0 , and so does any arrangement. If moreover $\sum w < 0$, a balanced arrangement would weigh $2c \sum w < 0$. It remains to find such weights for every $n \geq 5$; a square of side less than five cannot hold A at all.

For $5 \leq n \leq 9$ exact rational weights are listed in `verify/heptomino/exact-weights-A.jsonl`. For larger n write $n = 5K + m$ with $K \geq 1$ and $5 \leq m \leq 9$, and build w from K blocks of length five followed by a core:

$$w = (a_K Z + d, \dots, a_1 Z + d, C_m), \quad Z = (-5, 1, 0, -1, 5), \quad d = \left(-\frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2}\right),$$

with $a_1 = \frac{7}{2}$, $a_{k+1} = 5a_k$, and fixed cores C_m of negative sum given in `shapeA_family.json`. Each block sums to zero, since Z and d do, so $\sum w = \sum C_m < 0$. A window lying inside one block does not depend on its scale: $P \cdot Z = -10 + 1 + 0 - 1 + 10 = 0$, and the three Q -windows of Z are $-5 + 5 + 0$, $1 + 0 - 1$ and $0 - 5 + 5$, all zero. Such windows meet the bounds through d . A window across the boundary between blocks of scales $5s$ and s has the form $\alpha s + \delta$ with $\alpha > 0$, and it meets its bound at $s = a_1$, hence at every larger scale. The windows that touch the core take finitely many values and are checked directly. $\square$

The checker `verify/heptomino/check_shapeA.py` uses exact rational arithmetic and only the standard library. It verifies the window bounds and the negative sum explicitly for every side from 5 to 400 , checks the induction step symbolically, and rejects four mutants: a wrong core, the ratio 4 in place of 5 , no offset d , and a wrong profile. The other heptominoes on the list remain open. For them the weighting method fails: a fractional relaxation of the placement problem is feasible on some larger squares, so no weights of this kind can exist there.

VERIFICATION AND USE OF AI TOOLS

The searches, certificates and parts of the proofs were produced with AI coding assistants, Claude (Anthropic) and Codex (OpenAI), and then checked as each section describes. In particular the extreme-king reduction and DRUP certificates for touch cycles, and the finite weights for the heptomino on small squares, were first found by Codex. The author takes responsibility for every statement. All programs and data are in the folder that accompanies this paper; each section names its subfolder, and `README.md` gives the commands and running times.

REFERENCES

- [1] E. Friedman, *Math Magic: unsolved problems*, <https://erich-friedman.github.io/mathmagic/unsolved.html>, accessed 9 October 2026.
- [2] E. Friedman, *Math Magic*, problem of the month, March 2005 (armies of bishops), <https://erich-friedman.github.io/mathmagic/0305.html>, accessed 9 October 2026.
- [3] E. Friedman, *Math Magic*, problem of the month, May 2007 (polyominoes in squares), <https://erich-friedman.github.io/mathmagic/0507.html>, accessed 9 October 2026.
- [4] E. Friedman, *Math Magic*, problem of the month, October 2013 (regular digraphs of chess pieces), <https://erich-friedman.github.io/mathmagic/1013.html>, accessed 9 October 2026.
- [5] E. Friedman, *Math Magic*, problem of the month, March 2015 (touching letters), <https://erich-friedman.github.io/mathmagic/0315.html>, accessed 9 October 2026.