

Descent sets of a permutation and its inverse: almost every pair occurs

Vamshi Jandhyala

Manuscript prepared 8 October 2026

© The author. Released under the CC BY license (International 4.0).

Abstract

For a permutation w of $\{1, \dots, n\}$ let $D(w)$ be its descent set. Stanley asked for the number $f(n)$ of pairs (S, T) of subsets of $\{1, \dots, n-1\}$ that arise as $(D(w), D(w^{-1}))$, and in particular for the growth rate $L = \lim f(n)^{1/n}$, which lies between 2 and 4. We prove that almost every pair arises: $1 - f(n)/4^{n-1} \leq 6e^{-n/150}$ for every $n \geq 1$, so $L = 4$, and an elementary obstruction shows that the rate $1/150$ cannot be improved beyond $\log(4/3)$. The pairs that violate a necessary condition of Gale–Ryser type have proportion $(3/4)^{n+O(\log^2 n)}$, so the rate is exactly $\log(4/3)$ if a dominance-order criterion for occurrence, supported by computation, is correct. For a typical pair the proof constructs two standard Young tableaux of a common shape, a hook with a second row, whose descent sets are S and T , and then applies the Robinson–Schensted correspondence. We also find the asymptotics of the number $g(n)$ of permutations w such that w and w^{-1} are both alternating, confirming a prediction of Stanley: $g(n) = \frac{16}{\pi^2} \left(\frac{4}{\pi^2}\right)^n n! \left(1 + \frac{\pi^4}{48n} + O(n^{-2})\right)$. Every result proved here has been formally verified in the Lean proof assistant. Finally, we give computational evidence that (S, T) arises exactly when two intervals in dominance order meet, although the Schur supports of ribbons are often not intervals, and we confirm Gessel’s conjecture on the largest class $\#\{w : D(w) = S, D(w^{-1}) = T\}$ for $n \leq 22$.

Mathematics Subject Classifications: 05A05, 05A16, 05E05, 60C05

1 Introduction

How strongly does the descent pattern of a permutation constrain the descent pattern of its inverse? Small examples show substantial restrictions, yet we shall prove that almost every pair of patterns becomes possible as the number of entries grows.

Write $[n] = \{1, \dots, n\}$ and, for $w = w_1 \cdots w_n \in S_n$, let $D(w) = \{i \in [n-1] : w_i > w_{i+1}\}$. The descent set of the inverse can be read off w itself: $i \in D(w^{-1})$ exactly when

London, United Kingdom.

the value $i + 1$ appears to the left of the value i in w . Take $n = 3$. The six permutations give the pairs

$$\begin{aligned} 123 &\mapsto (\emptyset, \emptyset), & 132 &\mapsto (\{2\}, \{2\}), & 213 &\mapsto (\{1\}, \{1\}), \\ 231 &\mapsto (\{2\}, \{1\}), & 312 &\mapsto (\{1\}, \{2\}), & 321 &\mapsto (\{1, 2\}, \{1, 2\}), \end{aligned}$$

so only 6 of the 16 pairs of subsets of $\{1, 2\}$ occur. Let $f(n)$ be the number of pairs (S, T) of subsets of $[n - 1]$ for which some $w \in S_n$ has $D(w) = S$ and $D(w^{-1}) = T$. The first values,

$$1, 2, 6, 22, 94, 426, 1938, 8724, 38724, 169438, 731390, 3119052, 13162228,$$

form OEIS A380394 [13].

Stanley [18] asked what can be said about $f(n)$, and in particular about the limit

$$L = \lim_{n \rightarrow \infty} f(n)^{1/n}.$$

The limit exists because f is supermultiplicative, as Zach Hunter observed in a comment on [18]: placing a permutation of $[n]$ before a permutation of $\{n + 1, \dots, n + m\}$ creates no descent at position n in either the permutation or its inverse, so $f(n + m) \geq f(n)f(m)$. There are 4^{n-1} candidate pairs of subsets, so $f(n) \leq 4^{n-1}$, and every pair (S, S) occurs, so $2 \leq L \leq 4$. The values $f(n)^{1/n}$ grow slowly, reaching only 3.53 at $n = 13$, and Stanley asked whether $L < 4$. It is not. The proportion $f(n)/4^{n-1}$ is only 0.785 at $n = 13$, but it tends to 1 exponentially fast.

Theorem 1.1. *For every $n \geq 1$,*

$$1 - \frac{f(n)}{4^{n-1}} \leq 6e^{-n/150}.$$

In particular $L = 4$.

The bound $L = 4$ alone has a three-line proof (Proposition 2.2): every pair with $|S| = |T|$ occurs, and there are $\binom{2n-2}{n-1}$ such pairs. Theorem 1.1 needs more, because a uniformly random pair has $||S| - |T||$ of order $\sqrt{n}$. The tool is an explicit family of tableau shapes, a hook with a second row of length b , together with a rule that fills the ascending runs of S (and of T) so that the descent set comes out right and both tableaux have the same shape. Figure 1 shows the outcome for one pair with $n = 14$, and Section 3 gives the construction. The rate $1/150$ is not the true one. If T contains $|S| + 1$ consecutive integers then (S, T) cannot occur, and counting such pairs gives $1 - f(n)/4^{n-1} \geq (3^{n-1} - 1)/4^{n-1}$ (Remark 3.4). This obstruction already has the full exponential rate of a Gale–Ryser condition that every occurring pair satisfies (Theorem 3.6), so the rate is exactly $\log(4/3)$ if the dominance criterion of Section 5 is correct (Corollary 3.7).

Stanley also considered the size of the largest class, $\beta_n(S, T) = \#\{w \in S_n : D(w) = S, D(w^{-1}) = T\}$. Gessel conjectured (see [18]) that the maximum is attained when $S = T = \{1, 3, 5, \dots\} \cap [n - 1]$, that is, when w and w^{-1} are both alternating. Let $g(n)$

Q , descent set S							P , descent set T						
1	2	3	4	5	7	13	1	2	5	6	8	9	12
6	10	11					3	10	13				
8							4						
9							7						
12							11						
14							14						

Figure 1: Two standard Young tableaux of shape $(7, 3, 1, 1, 1, 1)$ built by the construction of Section 3 for $S = \{5, 7, 8, 11, 13\}$ and $T = \{2, 3, 6, 9, 10, 12, 13\}$. Here i is a descent when $i + 1$ sits in a lower row. Inverse RSK gives $w = 1\ 4\ 5\ 7\ 14\ 11\ 13\ 10\ 6\ 8\ 9\ 3\ 12\ 2$, with $D(w) = S$ and $D(w^{-1}) = T$.

be the number of such w (OEIS A007999 [12]). Starting from a generating function he had found earlier [16], Stanley predicted [18] that $g(n) \sim c n! (\pi^2/4)^{-n}$. We confirm this, with the constant and the next term.

The number of ordinary alternating permutations is the Euler number E_n . If inversion imposed no further dependence, the baseline count would be $E_n^2/n!$.

Theorem 1.2.

$$g(n) = \frac{E_n^2}{n!} \left(1 + \frac{\pi^4}{48n} + O(n^{-2}) \right) = \frac{16}{\pi^2} \left(\frac{4}{\pi^2} \right)^n n! \left(1 + \frac{\pi^4}{48n} + O(n^{-2}) \right).$$

In words: of the E_n^2 pairs (u, v) of alternating permutations, a proportion close to $1/n!$ satisfies $v = u^{-1}$, as if the two were independent, and the first correction is $\pi^4/(48n)$. We have also checked Gessel’s conjecture itself for $n \leq 22$ (Remark 4.1).

Section 5 turns to the exact set of pairs. Computation for $n \leq 20$ supports a simple criterion in terms of dominance order (Conjecture 5.1), and we explain why it does not follow from known results on the supports of ribbon Schur functions. Section 6 describes a formal verification in Lean of every result proved here, and the computations.

2 Tableaux and row words

The task is to give two prescribed descent sets a common tableau shape, so we first translate descents into the relative rows of successive entries. We use English notation. A standard Young tableau (SYT) of shape $\lambda \vdash n$ has descent set $\text{Des}(P) = \{i \in [n-1] : i+1 \text{ lies in a strictly lower row of } P \text{ than } i\}$. The Robinson–Schensted correspondence $w \mapsto (P(w), Q(w))$ is a bijection from S_n to pairs of SYT of a common shape, with $D(w) = \text{Des}(Q(w))$ and $D(w^{-1}) = \text{Des}(P(w))$; see [14, Section 7.23]. Hence:

Lemma 2.1. *A pair (S, T) occurs if and only if some shape $\lambda \vdash n$ has an SYT with descent set S and an SYT with descent set T .*

To construct a tableau we need only choose the row of each successive entry. For example, the first seven entries of Q in Figure 1 occupy rows 1, 1, 1, 1, 1, 2, 1. Encode an SYT P by its *row word* $r_1 r_2 \cdots r_n$, where r_i is the row of P containing i . A word in the positive integers is the row word of an SYT exactly when it is a lattice word: to keep columns increasing, in every prefix, and for every $k \geq 2$, the letter k occurs at most as often as the letter $k - 1$. The shape is the vector of letter counts. In this language

$$\text{Des}(P) = \{i : r_{i+1} > r_i\}.$$

Thus a descent means a move to a lower row. To prescribe exactly these moves, split the entries at the desired descents; within each piece the row numbers must not increase. For instance $S = \{2, 4\}$ on five entries gives the pieces $12 \mid 34 \mid 5$. Write $S \subseteq [n - 1]$ as the set of breaks between the ascending runs $R_0, R_1, \dots, R_{|S|}$, which are the maximal intervals of $[n]$ containing no break; thus R_j starts at 1 if $j = 0$ and at $s + 1$ if s is the j -th smallest element of S . A lattice word has descent set S exactly when it is weakly decreasing along every run and strictly increases from the last letter of each run to the first letter of the next.

Proposition 2.2. *Every pair (S, T) with $|S| = |T|$ occurs. Consequently $f(n) \geq \binom{2n-2}{n-1} \geq 4^{n-1}/(2n-1)$, and $L = 4$.*

Proof. Let $|S| = k$. The word with the letter 1 on R_0 and, on each later run R_j , the letter $j + 1$ followed by 1's is a lattice word of shape $(n - k, 1^k)$ with descent set S . The same shape serves T , so Lemma 2.1 applies. Summing over k gives $\sum_k \binom{n-1}{k}^2 = \binom{2n-2}{n-1}$, the largest of the $2n - 1$ coefficients of $(1 + x)^{2n-2}$, so at least $4^{n-1}/(2n - 1)$. $\square$

Remark 2.3. The pairs with $|S| = |T|$ can also be realised without tableaux, by a permutation made of a decreasing and an increasing sequence that share one entry, the *pivot*. The increasing and decreasing permutations handle $|S| = 0$ and $|S| = n - 1$, respectively, so suppose $0 < k = |S| = |T| < n - 1$ and fix a position p and a value v . Mark the position p , each $s \in S$ with $s < p$, and $s + 1$ for each $s \in S$ with $s \geq p$. Mark values in the same way, using v and T . Put the marked values into the marked positions in decreasing order and the other values into the other positions in increasing order. If p and v are chosen so that the value v lands at position p , with exactly the unmarked values below v placed before it, then a descent occurs before the pivot exactly at marked positions and from the pivot onwards exactly before marked positions, so $D(w) = S$; interchanging positions and values gives $D(w^{-1}) = T$. A suitable (p, v) is a common vertex of two lattice paths in a $(n - 1 - k) \times k$ rectangle, one built from S and one from T , which must cross. For $S = \{1, 3\}$ and $T = \{1, 4\}$ this gives $w = 52314$. The details are in the author's answer to Stanley's question [7].

3 Almost every pair occurs

We enlarge the hook construction by giving both tableaux the same second row. For the smaller descent set, extra 2's are placed inside existing runs; for the larger one, more runs

begin with 2. Figure 2 shows both operations. To preserve the lattice condition, we first accumulate 1's in the early positions and reserve the extra 2's for later runs.

Fix n , put $m = \lfloor n/2 \rfloor$, and for $b \geq 1$, $c \geq 0$ with $n - c - b \geq b$ let

$$\lambda(c, b) = (n - c - b, b, 1^c),$$

a hook with a second row of length b . A row word of this shape uses the letter 1 for the first row, the letter 2 for the second row, and each of the *leg letters* $3, 4, \dots, c + 2$ once. For $X \subseteq [n - 1]$ with runs $R_0, \dots, R_{|X|}$, we need runs that begin after this early reserve has been formed. Call R_j *late* if $j \geq 2$ and R_j starts at a position greater than m . The early positions that do not start a run will carry 1's; count them by putting

$$A(X) = \#([1, m - 1] \setminus X),$$

which is the number of positions $p \in [2, m]$ that do not start a run. This is the early stock of 1's available to balance the eventual second-row letters.

For the pair in Figure 1, $s = 5$, $t = 7$, $c = s - 1 = 4$ and $b = t - s + 1 = 3$. The runs of S are $12345 \mid 67 \mid 8 \mid 91011 \mid 1213 \mid 14$. Putting two extra 2's at entries 10, 11 gives the top word in Figure 2. For T , the runs beginning at 10 and 13 can start with 2, because their predecessors end with 1. Both words have seven 1's, three 2's and one of each leg letter 3, 4, 5, 6; hence both have shape $(7, 3, 1^4)$.

entry	1	2	3	4	5	6	7	$m=7$	8	9	10	11	12	13	14
S	1	1	1	1	1	2	1		3	4	2	2	5	1	6

entry	1	2	3	4	5	6	7	$m=7$	8	9	10	11	12	13	14
T	1	1	2	3	1	1	4		1	1	2	5	1	2	6

Figure 2: The construction behind Figure 1. Vertical bars mark the prescribed run breaks; the dashed line is the midpoint $m = 7$. Blue entries are the leg letters. Red entries are the two extra second-row letters: inside a run for S , at the start of two runs for T . The grey early positions carry 1's and protect the lattice condition. The displayed words are the rows of Q and P , respectively.

Lemma 3.1 (extra twos). *Let $|S| \geq 1$, $c = |S| - 1$, and $b \geq 1$ with $n - c - b \geq b$. Suppose $A(S) \geq b$ and $\sum_{R_j \text{ late}} (|R_j| - 1) \geq b - 1$. Then some SYT of shape $\lambda(c, b)$ has descent set S .*

Proof. As in the top word of Figure 2, choose integers $0 \leq k_j \leq |R_j| - 1$ for the late runs with $\sum k_j = b - 1$, and put $k_j = 0$ for the other j . Fill R_0 with 1's, fill R_1 with a 2 followed by 1's, and fill each R_j with $j \geq 2$ by the next unused leg letter, then k_j letters 2, then 1's.

Each run is weakly decreasing. Across a break the word strictly increases, from 1 to the 2 that starts R_1 , or into a leg letter, which exceeds every earlier letter. So the descent set is S . There are $|S| - 1 = c$ leg letters, used once each in increasing order, and the first of them occurs after the 2 that starts R_1 . This settles the lattice condition for the letters $k \geq 3$. There are $1 + (b - 1) = b$ letters 2.

It remains to compare the letters 1 and 2. Up to position m the only possible 2 is the first letter of R_1 , which is preceded by the nonempty run R_0 of 1's. Every longer prefix contains at most b letters 2 and at least $A(S) \geq b$ letters 1, because every position in $[2, m]$ that does not start a run carries a 1. The shape is $\lambda(c, b)$. $\square$

Lemma 3.2 (two-started runs). *Let $|T| = c + b$ with $c \geq 0$, $b \geq 1$ and $n - c - b \geq b$. Suppose $A(T) \geq b$ and at least $b - 1$ late runs R_j of T have $|R_{j-1}| \geq 2$. Then some SYT of shape $\lambda(c, b)$ has descent set T .*

Proof. As in the bottom word of Figure 2, fill R_0 with 1's. Fill R_1 , and $b - 1$ late runs R_j with $|R_{j-1}| \geq 2$, with a 2 followed by 1's. Fill every other R_j with $j \geq 2$ with the next leg letter followed by 1's. A run of length at least 2 ends in 1, so each run that starts with a 2 follows a 1, and the word increases across every break. The rest of the argument is as for Lemma 3.1: there are $|T| - b = c$ leg letters and b letters 2. $\square$

Corollary 3.3. *Let $1 \leq s = |S| < t = |T|$, $c = s - 1$, $b = t - s + 1$ and $n - c - b \geq b$. If S satisfies the hypotheses of Lemma 3.1 and T those of Lemma 3.2, then (S, T) occurs.*

Proof. As $|T| = c + b$, the two lemmas give tableaux of the same shape $\lambda(c, b)$, and Lemma 2.1 applies. $\square$

We use the bounded-difference inequality in the following form [6, 8]. Let F be a function of independent fair bits $\xi_1, \dots, \xi_N$, and suppose that changing any one bit changes F by at most 1. In words, either tail at distance t from the mean is bounded exponentially in t^2/N . For $t > 0$

$$\Pr(F - \mathbb{E}F \geq t) \leq e^{-2t^2/N}, \quad \Pr(F - \mathbb{E}F \leq -t) \leq e^{-2t^2/N}. \quad (3.1)$$

Proof of Theorem 1.1. If $n < 240$ then $6e^{-n/150} > 6e^{-1.6} > 1$ and there is nothing to prove, so let $n \geq 240$. Let S, T be independent and uniform among subsets of $[n - 1]$, so that the indicators of $1, \dots, n - 1$ in S (and in T) are independent fair bits. Then $f(n)/4^{n-1}$ is the probability that (S, T) occurs, and we bound the probability of failure. To supply late capacity, count run starts whose preceding run has length at least two. For $X \subseteq [n - 1]$ define

$$G(X) = \#\{i \in [m + 2, n - 1] : i \in X, i - 1 \notin X\}.$$

It counts late starts immediately preceded by a nonbreak and then a break, so the preceding run has length at least two. This one count controls both lemmas.

- Every i counted by $G(X)$ starts a run at $i + 1 > m$ whose predecessor contains $i - 1$ and i . That run has index at least 2 unless $i = \min X$. So at least $G(X) - 1$ late runs follow a run of length at least 2.

- Let R_j be a late run with $|R_{j-1}| \geq 2$. Unless R_j is the first late run, R_{j-1} is itself late and contributes $|R_{j-1}| - 1 \geq 1$ to $\sum_{\text{late}}(|R_k| - 1)$. Distinct j give distinct predecessors, so $\sum_{\text{late}}(|R_k| - 1) \geq G(X) - 2$.

Only the difference of the two sizes matters, so we ask for it directly. Consider the events

$$\mathcal{D} : ||S| - |T|| \leq \frac{n}{12}, \quad \mathcal{A}(X) : A(X) \geq \frac{n}{12} + 1, \quad \mathcal{G}(X) : G(X) \geq \frac{n}{12} + 2,$$

for $X \in \{S, T\}$. Suppose all five hold. If $|S| = |T|$ then (S, T) occurs by Proposition 2.2. The events are symmetric in S and T , and $w \mapsto w^{-1}$ exchanges the roles of S and T , so we may assume $s = |S| < t = |T|$. As $G(S) \geq 1$ we have $s \geq 1$, and $b = t - s + 1 \leq n/12 + 1$. This gives $b \leq A(S)$ and $b \leq A(T)$, while $b - 1 \leq n/12 \leq G(S) - 2$ and $b - 1 \leq G(T) - 1$, so by the two bullet points the hypotheses of Lemmas 3.1 and 3.2 hold. For the shape, note that T misses the $A(T)$ points of $[1, m - 1] \setminus T$ and, for each i counted by $G(T)$, the point $i - 1 \in [m + 1, n - 2]$. Hence $t \leq n - 1 - A(T) - G(T) \leq \frac{5n}{6} - 4$, and $c + 2b = t + (t - s) + 1 \leq \frac{5n}{6} - 4 + \frac{n}{12} + 1 < n$, so $n - c - b \geq b$. Corollary 3.3 shows that (S, T) occurs.

It remains to bound the probability that one of the five events fails, using (3.1).

- $|S| + (n - 1 - |T|)$ counts the elements of S and of the complement of T , so it is a sum of $2(n - 1)$ fair bits with mean $n - 1$, and $|S| - |T|$ is this sum minus $n - 1$. Each tail of $\mathcal{D}$ has probability at most $e^{-2(n/12)^2/(2(n-1))} \leq e^{-n/144}$.
- $A(X)$ is a sum of $m - 1 \leq n/2$ fair bits with mean $(m - 1)/2 \geq (n - 3)/4$. If $\mathcal{A}(X)$ fails then $A(X)$ is below its mean by more than $n/6 - 7/4$, which has probability at most $e^{-4(n/6 - 7/4)^2/n} \leq e^{-n/150}$.
- For $G(X)$ we count changes instead. Let $C(X)$ be the number of $i \in [m + 2, n - 1]$ such that exactly one of $i - 1$ and i lies in X . Along $[m + 1, n - 1]$ the rises counted by $G(X)$ and the opposite changes alternate, so $C(X) \leq 2G(X) + 1$. Replacing the bit of each $i \in [m + 2, n - 1]$ by the indicator of a change at i , and keeping the other bits, is a bijection of $\{0, 1\}^{n-1}$, because X can be read back from left to right. So $C(X)$ is a sum of $n - m - 2 \leq n/2$ fair bits, with mean at least $n/4 - 1$. If $\mathcal{G}(X)$ fails then $C(X) < n/6 + 5$, which is below the mean by more than $n/12 - 6$. This has probability at most $e^{-4(n/12 - 6)^2/n} \leq e^{-n/150}$, because $n \geq 240$.

A union bound over the five events, with two tails for $\mathcal{D}$, gives $1 - f(n)/4^{n-1} \leq 6e^{-n/150}$. $\square$

The change count is what improves on a direct use of (3.1) for $G(X)$, which treats each late bit as able to move $G(X)$ by one and so overstates its variance by a factor of four.

Remark 3.4. The construction does better than the theorem guarantees. With 20000 samples at each size and pseudorandom seed 11, it certifies approximately 0.80, 0.93, 0.986 and 0.9995 of uniformly random pairs at $n = 50, 100, 200, 400$. At $n = 400$ there

are 11 failures; the estimated failure probability is 0.00055, with a 95% Wilson interval approximately $[0.00031, 0.00098]$, so relative sampling uncertainty is substantial. The supplementary output gives exact counts and binomial confidence intervals. Its own limit is structural: hooks with one extra row can absorb a size difference $||S| - |T||$ of only a small fraction of n . With exact large-deviation rates in place of (3.1), the best balance of the three kinds of event used above gives a rate of about 1/100.

In the other direction, the exact rate is governed by a necessary condition of Gale–Ryser type.

For $X \subseteq [n-1]$ let $\alpha(X)$ be the composition of n formed by the lengths of the ascending runs, and let $\delta(X)$ be the composition formed by the lengths of the maximal blocks $\{v, v+1, \dots, v+\ell-1\}$ of consecutive integers with $v, \dots, v+\ell-2 \in X$. For a composition γ and $k \geq 1$ let $\text{top}_k(\gamma)$ be the sum of its k largest parts, or of all its parts if there are fewer than k .

Lemma 3.5. *If (S, T) occurs, then for every $k \geq 1$*

$$\text{top}_k(\alpha(S)) \leq \sum_j \min(k, \delta_j(T)) \quad \text{and} \quad \text{top}_k(\alpha(T)) \leq \sum_j \min(k, \delta_j(S)). \quad (3.2)$$

Proof. Let $D(w) = S$ and $D(w^{-1}) = T$. A block of $\delta(T)$ is a set of consecutive values that appear in w from right to left, while an ascending run of w lists its values from left to right in increasing order, so a run and a block share at most one entry. The 0–1 matrix recording which runs meet which blocks has row sums $\alpha(S)$ and column sums $\delta(T)$, so any k rows contain at most $\sum_j \min(k, \delta_j(T))$ ones. This is the first inequality; the second is the first one for w^{-1} . $\square$

The consecutive-integer obstruction is the case $k = |S| + 1$: if T contains $|S| + 1$ consecutive integers, then $\delta(T)$ has a part longer than $|S| + 1$, while $\text{top}_{|S|+1}(\alpha(S)) = n$. Counting the pairs with $\{1, \dots, |S| + 1\} \subseteq T$, and their images under $(S, T) \mapsto (T, S)$, gives

$$f(n) \leq 4^{n-1} - 3^{n-1} + 1, \quad \text{equivalently} \quad 1 - \frac{f(n)}{4^{n-1}} \geq \frac{3^{n-1} - 1}{4^{n-1}},$$

with equality at $n = 2$. The next theorem shows that this obstruction already has the full exponential rate of condition (3.2).

Theorem 3.6. *Let P_n be the proportion of pairs (S, T) of subsets of $[n-1]$ that violate (3.2). Then*

$$\frac{3^{n-1} - 1}{4^{n-1}} \leq P_n \leq 2 \left(4n^3 + 2^{m-1} \binom{n}{m}^2 \right) \left(\frac{3}{4} \right)^{n-1} + \frac{4n^m}{2^n}, \quad m = \lceil 2 \log_2 n \rceil + 1,$$

so $P_n = (3/4)^{n+O(\log^2 n)}$. Every pair counted by P_n fails to occur, so $1 - f(n)/4^{n-1} \geq P_n$.

Proof. The lower bound is the count above. For the upper bound let S, T be independent and uniform, and write $\alpha = \alpha(S)$ with r parts and $\delta = \delta(T)$ with q parts; both are

uniform compositions of n . An interval $\{v, \dots, v + \ell - 1\}$ is a part of α only if none of $v, \dots, v + \ell - 2$ lies in S , and a part of δ only if all of them lie in T ; each has probability $2^{1-\ell}$, independently for disjoint intervals. Three counts follow.

- $\sum_{u \geq 1} 2^{-u} \Pr(r \leq u) = \sum_r \Pr(r) 2^{1-r} = 2^{2-n} \sum_r \binom{n-1}{r-1} 2^{-r} = (3/4)^{n-1}$, and in the same way $\sum_q \Pr(q) 2^{-q} = \frac{1}{2}(3/4)^{n-1}$.
- Write $E_k(\delta) = \sum_j (\delta_j - k)_+$. If $E_k(\delta) \geq e$, choose the $a \geq 1$ parts longer than k ; summing $2^{1-\ell}$ over their positions and lengths gives, whenever $n^2 2^{1-k} \leq \frac{1}{2}$,

$$\Pr(E_k(\delta) \geq e) \leq \sum_{a \geq 1} \binom{n}{a}^2 2^{1-e-a(k-1)} \leq 8n^2 2^{-k-e}.$$

- In the same way, $\Pr(r \geq m, \text{top}_m(\alpha) \geq x) \leq 2^{m+1} \binom{n}{m}^2 2^{-x}$.

Since $\sum_j \min(k, \delta_j) = n - E_k(\delta)$, the first inequality of (3.2) fails at k exactly when $\text{top}_k(\alpha) + E_k(\delta) > n$. Two consequences follow. As the $r - k$ smallest parts of α are at least 1, $\text{top}_k(\alpha) \leq n - (r - k)$ when $k < r$, so $e = E_k(\delta)$ satisfies $e \geq 1$ and $r \leq k + e - 1$; when $k \geq r$ the same holds because $\text{top}_k(\alpha) = n$. And as $E_k(\delta) \leq E_1(\delta) = n - q$, also $\text{top}_k(\alpha) > q$.

Because α and δ are independent, if the failure occurs at some $k > m$, the first consequence and the second and first counts bound its probability by $\sum_k \sum_{e \geq 1} 8n^2 2^{-k-e} \Pr(r \leq k + e - 1) \leq 4n^3 (3/4)^{n-1}$. If it occurs at some $k \leq m$, the second consequence gives $r < m$, an event of probability at most $n^m 2^{1-n}$, or $\text{top}_m(\alpha) \geq q + 1$ with $r \geq m$, whose probability is at most $\sum_q \Pr(q) 2^{m+1} \binom{n}{m}^2 2^{-q-1} = 2^{m-1} \binom{n}{m}^2 (3/4)^{n-1}$. Exchanging S and T treats the second inequality of (3.2) in the same way. $\square$

Corollary 3.7. *If Conjecture 5.1 holds, then $1 - f(n)/4^{n-1} = P_n = (3/4)^{n+O(\log^2 n)}$.*

Proof. Conjecture 5.1 says that (3.2) is also sufficient (Section 5), so the pairs that fail to occur are exactly those counted by P_n . $\square$

Exact computation suggests the finer order. The probability that the first inequality of (3.2) fails can be computed by summing over pairs of partitions, because $\alpha(S)$ and $\delta(T)$ are independent uniform compositions. Twice it, 0.05309 at $n = 20$, matches $1 - f(20)/4^{19}$ to the digits shown, and for $n \leq 44$ its ratio to $(3/4)^n$ grows roughly linearly, from 10.7 at $n = 24$ to 20.4 at $n = 44$.

Conjecture 3.8. $(1 - f(n)/4^{n-1})/(n(3/4)^n)$ is bounded above and below by positive constants.

4 Doubly alternating permutations

Let $g(n)$ count the $w \in S_n$ with $D(w) = D(w^{-1}) = \{1, 3, 5, \dots\} \cap [n-1]$. Stanley [16, Theorem 3.1] proved

$$\sum_{n \geq 0} g(n) x^n = \sum_{k \geq 0} E_{2k+1}^2 \frac{u^{2k+1}}{(2k+1)!} + \frac{1}{\sqrt{1-x^2}} \sum_{k \geq 0} E_{2k}^2 \frac{u^{2k}}{(2k)!}, \quad u = \frac{1}{2} \log \frac{1+x}{1-x}. \quad (4.1)$$

The odd and even sums separate the two parities of n ; substitution of $u = x + x^3/3 + \dots$ mixes earlier Euler-number terms into each coefficient. Gessel and Zhuang [5, Section 2.4] rederive this generating function through symmetric functions. Our contribution here is the asymptotic expansion, rather than a new generating function.

The series $\sum_m E_m^2 u^m / m!$ has radius of convergence 0, so singularity analysis does not apply directly. A direct comparison of coefficients does. The term indexed by $m = n$ supplies the leading value $E_n^2 / n!$, the term $m = n - 2$ supplies the $1/n$ correction, and factorial decay bounds the sum of all earlier terms uniformly.

Proof of Theorem 1.2. Put $a_m = E_m^2 / m!$ and $\rho = 4/\pi^2$. Since $u = x + x^3/3 + x^5/5 + \dots$ is odd, (4.1) reads

$$g(n) = \sum_{0 \leq j \leq n/2} a_{n-2j} c_{n,n-2j}, \quad c_{n,m} = \begin{cases} [x^n] u^m, & n \text{ odd}, \\ [x^n] (1-x^2)^{-1/2} u^m, & n \text{ even}. \end{cases}$$

Here j measures how far the Euler-number index is below n , and $c_{n,m}$ is the weight contributed by the substitution.

Coefficients. Coefficientwise $u \leq x(1-x^2)^{-1}$ and $(1-x^2)^{-1/2} \leq (1-x^2)^{-1}$, so for $m = n - 2j$

$$c_{n,m} \leq [x^n] x^m (1-x^2)^{-m-1} = \binom{m+j}{j} \leq \frac{n^j}{j!}.$$

Thus the substitution weight grows at most like $n^j/j!$, which will be dominated by the loss of $2j$ factorial factors. For the two leading weights, $c_{n,n} = 1$ and $c_{n,n-2} = \frac{n-2}{3} + \frac{[n \text{ even}]}{2}$.

Euler numbers. From the partial fractions of $\sec z + \tan z$, $E_m / m! = 2(2/\pi)^{m+1} \sigma_m$ with $\sigma_m = \sum_{k \geq 0} (-1)^{k(m+1)} (2k+1)^{-m-1}$; see [17]. For $m = 0$ the series is the alternating sum $\sigma_0 = \pi/4$, and $a_0 = 1$. For even $m \geq 2$ the alternating-series estimate gives $1 - 3^{-m-1} \leq \sigma_m \leq 1$. For odd $m \geq 1$ all terms are positive and $1 \leq \sigma_m \leq \sigma_1 = \pi^2/8$. Consequently $\pi/4 \leq \sigma_m \leq \pi^2/8$ for every $m \geq 0$. Also, for $m \geq 1$,

$$|\sigma_m - 1| \leq \sum_{k \geq 1} (2k+1)^{-m-1} \leq 3^{-m-1} + \int_1^\infty (2x+1)^{-m-1} dx = 3^{-m-1} + \frac{3^{-m}}{2m} = O(3^{-m}).$$

The integral bounds all terms after the first omitted term, uniformly in m . Hence $a_m = 4\rho^{m+1} m! \sigma_m^2$, and with $K = (\pi/2)^2$,

$$\frac{a_{n-2j}}{a_n} \leq K \rho^{-2j} \frac{(n-2j)!}{n!}.$$

Apart from a fixed factor and an exponential factor in j , this ratio is the reciprocal of $2j$ successive factorial factors.

Tail. Write $R_n = \sum_{2 \leq j \leq \lfloor n/2 \rfloor} a_{n-2j} c_{n,n-2j} / a_n$. This is the total relative contribution discarded after the first two terms. If $n - 2j \geq n/2$ then $(n - 2j)!/n! \leq (n/2)^{-2j}$. Putting $q_n = 4/(\rho^2 n)$, the sum of these terms is at most

$$K \sum_{j \geq 2} \frac{q_n^j}{j!} \leq \frac{K q_n^2 e^{q_n}}{2} = O(n^{-2}).$$

The last bound follows from $j! \geq 2(j - 2)!$ and sums the entire near tail.

For the far tail put $M = \lfloor n/2 \rfloor$. There are at most n terms with $m = n - 2j < n/2$, and each has $c_{n,m} \leq \binom{m+j}{j} \leq 2^n$ and $a_m/a_n \leq K \rho^{-n} M!/n!$, since $0 < \rho < 1$. For $n \geq 2$, the $n - M$ factors in $n!/M!$ are all at least $n/2$, so the sum of the far tail is at most

$$nK \left(\frac{2}{\rho} \right)^n \frac{M!}{n!} \leq nK \left(\frac{2}{\rho} \sqrt{\frac{2}{n}} \right)^n.$$

Its logarithm is $-\frac{1}{2}n \log n + O(n)$, so it is $o(n^{-r})$ for every fixed $r > 0$. Combining the two ranges proves $R_n = O(n^{-2})$, with a constant independent of n .

Main terms. The term $j = 0$ is a_n . For $j = 1$, $a_{n-2}/a_n = \sigma_{n-2}^2/(\rho^2 \sigma_n^2 n(n-1)) = (1 + O(3^{-n})) / (\rho^2 n(n-1))$. Multiplying by $c_{n,n-2} = n/3 + O(1)$ gives $1/(3\rho^2 n) + O(n^{-2}) = \pi^4/(48n) + O(n^{-2})$. Finally $a_n = 4\rho^{n+1}n! \sigma_n^2 = \frac{16}{\pi^2} \rho^n n! (1 + O(3^{-n}))$. $\square$

Numerically, $n(g(n)n!/E_n^2 - 1)$ equals 2.1130, 2.0694 and 2.0296 at $n = 40, 80, 159$, against $\pi^4/48 = 2.0294$.

Remark 4.1. By a theorem of Gessel [4], $\beta_n(S, T) = \langle r_\alpha, r_\beta \rangle$, where r_α and r_β are the ribbon Schur functions of S and T (Section 5). Ribbon Schur functions are Schur positive, so the Cauchy–Schwarz inequality gives $\beta_n(S, T)^2 \leq \beta_n(S, S) \beta_n(T, T)$, and the maximum $M(n)$ of β_n is attained on the diagonal [15, Supplementary Problem 132(a)]. Moreover $\beta_n(S, S) = \sum_\lambda c_\lambda(S)^2$, where $c_\lambda(S)$ is the number of SYT of shape λ with descent set S . We computed $\beta_n(S, S)$ for every $S \subseteq [n - 1]$ with $n \leq 22$. For $4 \leq n \leq 22$ the maximum is attained only at $S = \{1, 3, 5, \dots\} \cap [n - 1]$ and $S = \{2, 4, 6, \dots\} \cap [n - 1]$, with common value $g(n)$, and every other value is at most $2g(n)/3$. By the equality case of Cauchy–Schwarz, every pair (S, T) with $\beta_n(S, T) = M(n)$ then has both S and T alternating. So Gessel’s conjecture holds for $n \leq 22$; for example $g(22) = 4705365925872$. For $n \leq 8$ we confirmed this by enumerating S_n directly. The computation also shows that $\beta_n(S, S)$ has local maxima away from the alternating sets, under the moves that add, delete or shift one element of S by one position; for example $S = \{2, 4, 7, 9\}$ at $n = 11$, where $\beta_{11}(S, S) = 1808$ while every neighbour gives at most 1751. Thus strict ascent using only these add, delete and adjacent-shift moves can stop before reaching an alternating set. This obstructs that particular improvement strategy, without ruling out other local methods.

5 The exact set of pairs

To decide a particular pair, we need to know which shapes are available for each descent set. For example, $S = \{1, 3\}$ on six entries has run lengths $\alpha = (1, 2, 3)$. Joining rows of these lengths, with successive rows overlapping in one column, gives its ribbon. Its sorted row lengths are $(3, 2, 1)$ and its sorted column lengths are $(2, 2, 1, 1)$, whose transpose is $(4, 2)$. The partition $(3, 3)$ lies between $(3, 2, 1)$ and $(4, 2)$ in dominance order, but does not occur in this ribbon's Schur support. Thus interval overlap is a necessary test whose possible sufficiency for pairs needs explanation, not an automatic consequence of full individual supports.

Dominance order compares partial sums: $\mu \leq \lambda$ means that the first k parts of μ sum to at most those of λ , for every k . A ribbon Schur function r_α records the available shapes: the coefficient of s_λ is the number of SYT of shape λ with descent set S . Let α, β be the compositions of n with descent sets S, T and let r_α be the ribbon Schur function. By Gessel's theorem (S, T) occurs if and only if r_α and r_β share a Schur constituent. Every constituent s_λ of r_α satisfies $\text{rows}(\alpha) \leq \lambda \leq \text{cols}(\alpha)^t$ in dominance order, where $\text{rows}(\alpha)$ and $\text{cols}(\alpha)$ are the row and column lengths of the ribbon sorted into partitions [9]. Dominance order on partitions of n is a lattice, so the two intervals meet if and only if the join of the lower ends lies below the meet of the upper ends.

Conjecture 5.1. A pair (S, T) occurs if and only if the dominance intervals

$$[\text{rows}(\alpha), \text{cols}(\alpha)^t] \quad \text{and} \quad [\text{rows}(\beta), \text{cols}(\beta)^t]$$

meet.

In words, the conjecture asks whether overlapping ranges of allowable shapes always contain a shape actually available to both descent sets, despite gaps in individual supports.

The condition is necessary. Because dominance order is a lattice, the two intervals meet exactly when $\text{rows}(\alpha) \leq \text{cols}(\beta)^t$ and $\text{rows}(\beta) \leq \text{cols}(\alpha)^t$; as $\text{rows}(\alpha)$ and $\text{cols}(\alpha)$ are $\alpha(S)$ and $\delta(S)$ sorted, this is condition (3.2), which Lemma 3.5 proves necessary. By Corollary 3.7, the conjecture would also determine the exact rate in Theorem 1.1. We have checked sufficiency pair by pair for $n \leq 20$, comparing the criterion with the actual set of pairs, and found no disagreement; in particular the criterion reproduces A380394, including $f(13) = 13162228$, and gives $f(20) = 260283514168$. Sufficiency would follow at once if every ribbon had full support, meaning that every λ in the interval occurs, but that is false. McNamara and van Willigenburg determined the support for one class of ribbons [9]. Gaetz, Hardt and Sridhar studied when two ribbons have equal supports [3], and Azenhas and Mamede classified the monotone ribbons with full support [1]. Both of these papers assume that every row of the ribbon has at least two boxes, whereas the ribbon of a uniformly random S has about $n/4$ rows of length one. By direct computation, 4 of the 32 ribbons with $n = 6$ fail to have full support (for instance $\alpha = (1, 2, 3)$ misses $(3, 3)$), as do 544 of the 1024 ribbons with $n = 11$. The missing partitions seem never to be the only common ones. Table 1 shows how the pairs that occur are spread.

$ S - T $	0	1	2	3	4	5
$n = 11$	1	0.940	0.702	0.319	0.056	0.002
$n = 12$	1	0.966	0.807	0.481	0.152	0.015

Table 1: Proportion of pairs (S, T) that occur, by $||S| - |T||$.

6 Formal verification and computations

Theorems 1.1 and 1.2 have been verified in the Lean 4 proof assistant [11] with the Mathlib library [10], and the verified statements use only Lean’s standard axioms. The formal proof of Theorem 1.1 follows Section 3 and gives the same bound. It proves Lemma 2.1 in both directions with Fomin’s growth diagrams [2] in place of the insertion algorithm: the backward local rule turns two SYT of a common shape with descent sets S and T into a permutation w with $D(w) = S$ and $D(w^{-1}) = T$, and the forward rule recovers such a pair of tableaux from any permutation. The formal proof of Theorem 1.2 does not assume (4.1): it proves Stanley’s generating function from the definition of $g(n)$, and the series for $E_m/m!$ used above. Proposition 2.2 is verified as well, through the construction of Remark 2.3 with the pivot chosen by a discrete intermediate value argument. The lower bound $1 - f(n)/4^{n-1} \geq (3^{n-1} - 1)/4^{n-1}$ of Remark 3.4 is verified, and so is the inequality $\beta_n(S, T)^2 \leq \beta_n(S, S) \beta_n(T, T)$ of Remark 4.1, which the formal proof obtains from flag characters of S_n rather than from ribbon Schur functions. So are Lemma 3.5, the necessary direction of Conjecture 5.1, Theorem 3.6, with its explicit bounds and with an explicit constant in the $O(\log^2 n)$, and Corollary 3.7. The formal proof of Theorem 3.6 counts pairs instead of computing probabilities, and its two union bounds become one lemma: a family of j blocks is recorded by its j first and j last positions, and these alone determine the positions inside the family, which gives the factor $\binom{n}{j}^2$. The Schur-function statements of Remark 4.1 and Section 5, the sufficiency in Conjecture 5.1, and all computed values are not formally verified.

All claims marked as computed were checked by computer. Each certificate produced by the construction of Section 3 was verified as a lattice word of the stated shape with the stated descent set (339296 row words for $n \leq 16$), and for $n \leq 9$ every certified pair was confirmed by enumeration. The failure probabilities in Remark 3.4 were computed in extended precision by summing over all pairs of partitions of n , and agree with the exact failure count at $n = 20$. The supplementary archive contains the Lean sources with the pinned toolchain and Mathlib revision, a theorem-and-axiom driver, and the computation scripts with commands and saved outputs. The pairwise check compares tableau supports directly with dominance-interval intersections; it does not infer agreement merely from matching totals. A second program, sharing no code with the first, computes the Schur coefficients $c_\lambda(S)$ by a dynamic programme on Young’s lattice; it repeated the pairwise check for every ordered pair with $n \leq 20$ and the diagonal maxima for $n \leq 22$. Every other number stated in the text was recomputed by separate code written for this purpose. Sampling output records seed 11 and Wilson binomial confidence intervals. File hashes

identify the exact source snapshot used for each check.

Acknowledgements

The question is Richard Stanley’s [18]. *Status*. All proved results of this paper are formally verified in Lean 4 (Section 6), and all computations were independently rechecked; the proofs have not yet been checked line by line by a human. *Origin*. The results and proofs were found with the AI systems Claude (Anthropic) and Codex (OpenAI), directed by the author.

References

- [1] O. Azenhas and R. Mamede. A classification of monotone ribbons with full Schur support with application to the classification of full equivalence classes. Preprint, 2018. [arXiv:1812.04705](https://arxiv.org/abs/1812.04705).
- [2] S. Fomin. Schensted algorithms for dual graded graphs. *J. Algebraic Combin.*, 4(1):5–45, 1995. [doi:10.1023/A:1022404807578](https://doi.org/10.1023/A:1022404807578).
- [3] M. Gaetz, W. Hardt and S. Sridhar. Support equalities among ribbon Schur functions. *Electron. J. Combin.*, 26(3):#P3.52, 2019. [doi:10.37236/8229](https://doi.org/10.37236/8229).
- [4] I. M. Gessel. Multipartite P -partitions and inner products of skew Schur functions. In *Combinatorics and Algebra*, volume 34 of *Contemp. Math.*, pages 289–317. Amer. Math. Soc., Providence, RI, 1984. [doi:10.1090/conm/034/777705](https://doi.org/10.1090/conm/034/777705).
- [5] I. M. Gessel and Y. Zhuang. Two-sided permutation statistics via symmetric functions. *Forum Math. Sigma*, 12:e93, 2024. [doi:10.1017/fms.2024.89](https://doi.org/10.1017/fms.2024.89).
- [6] W. Hoeffding. Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.*, 58(301):13–30, 1963. [doi:10.1080/01621459.1963.10500830](https://doi.org/10.1080/01621459.1963.10500830).
- [7] V. Jandhyala. Answer to “Descent sets and inverse descent sets of a permutation”. MathOverflow, 2026. <https://mathoverflow.net/a/515793>, accessed 8 October 2026.
- [8] C. McDiarmid. On the method of bounded differences. In *Surveys in Combinatorics, 1989*, volume 141 of *London Math. Soc. Lecture Note Ser.*, pages 148–188. Cambridge Univ. Press, 1989. [doi:10.1017/CB09781107359949.008](https://doi.org/10.1017/CB09781107359949.008).
- [9] P. R. W. McNamara and S. van Willigenburg. Maximal supports and Schur-positivity among connected skew shapes. *European J. Combin.*, 33(6):1190–1206, 2012. [doi:10.1016/j.ejc.2012.02.001](https://doi.org/10.1016/j.ejc.2012.02.001).
- [10] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs*, pages 367–381, 2020. [doi:10.1145/3372885.3373824](https://doi.org/10.1145/3372885.3373824).

- [11] L. de Moura and S. Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction, CADE 28*, volume 12699 of *Lecture Notes in Comput. Sci.*, pages 625–635. Springer, 2021. doi:[10.1007/978-3-030-79876-5_37](https://doi.org/10.1007/978-3-030-79876-5_37).
- [12] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, sequence A007999. <https://oeis.org/A007999>, accessed 8 October 2026.
- [13] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, sequence A380394. <https://oeis.org/A380394>, accessed 8 October 2026.
- [14] R. P. Stanley. *Enumerative Combinatorics, Volume 2*. Cambridge Univ. Press, Cambridge, 1999.
- [15] R. P. Stanley. *Enumerative Combinatorics, Volume 2*. Second edition. Cambridge Univ. Press, Cambridge, 2023.
- [16] R. P. Stanley. Alternating permutations and symmetric functions. *J. Combin. Theory Ser. A*, 114(3):436–460, 2007. doi:[10.1016/j.jcta.2006.06.008](https://doi.org/10.1016/j.jcta.2006.06.008).
- [17] R. P. Stanley. A survey of alternating permutations. In *Combinatorics and Graphs*, volume 531 of *Contemp. Math.*, pages 165–196. Amer. Math. Soc., Providence, RI, 2010. doi:[10.1090/conm/531/10466](https://doi.org/10.1090/conm/531/10466).
- [18] R. P. Stanley. Descent sets and inverse descent sets of a permutation. MathOverflow question 486548, 2025. <https://mathoverflow.net/q/486548>, accessed 8 October 2026.