

DOES THE SMALLEST ENCLOSING COPY FIT? RANDOM POINTS IN A CONVEX POLYGON

VAMSHI JANDHYALA

ABSTRACT. Choose n independent uniform points in a convex polygon K and let Q_n be a smallest similar copy of K , of any orientation, that contains them all. We show that the probability that Q_n can be chosen inside K converges, as $n \rightarrow \infty$, to an explicit constant $p(K)$, and we give a formula for $p(K)$ as an integral over a Poisson limit model. For an equilateral triangle $p = 13/48$ and for a square $p = 1/4$; numerically, p ranges over $[13/48, 25/72)$ for triangles, and we conjecture that the equilateral triangle is the minimiser. For regular q -gons the constants are not monotone in q (the computed values for $q = 5, 7$ lie below their even neighbours) and appear to settle near 0.146 (computed up to $q = 200$), far from the value $1/2$ that a Mathematics Stack Exchange answer establishes for the disk. The answer is governed by the freedom to rotate the enclosing copy, which has no analogue for the disk.

Status of this preprint. Every theorem, lemma and corollary is formally verified in Lean 4 (Section 7), using only Lean’s standard axioms. The exposition, the numerical results of Section 8 and Conjecture 4 are not formally verified, and this version has not yet been independently reviewed.

1. INTRODUCTION

Drop n random points into a disk and draw the smallest circle containing them. Does that circle lie inside the disk? A question on Mathematics Stack Exchange [7] asked for the limiting probability, and an answer there shows that it is exactly $1/2$. A follow-up on MathOverflow [4] asked whether the same holds when the disk and the circle are replaced by a regular polygon and a smallest enclosing regular polygon of the same kind, which may be rotated. The asker’s simulations for two points suggested a family resemblance with the disk.

The answer is no, and the reason is instructive. A smallest enclosing polygon wants to turn slightly: turning by an angle of order $1/n$ lets it hug the outermost points better, and the same turn pushes its corners out by a distance of order $1/n$, the same order as the gaps between the outermost points and the sides. Whether the copy fits is decided by a competition between these two effects, and it can be computed exactly.

Results. Let K be a convex polygon and $X_1, \dots, X_n$ independent uniform points in K . A *copy* of K is a set $\lambda R_\theta K + c$ with $\lambda > 0$, R_θ a rotation and $c \in \mathbb{R}^2$. Let E_n be the event that some copy of minimal λ containing all X_j lies in K .

Theorem 1. $\mathbb{P}(E_n)$ converges to the probability $p(K)$ of the event E of the limit model of Section 2, and $p(K)$ is given by the formula of Theorem 10.

Corollary 2 (Triangles). *For a triangle with sides L_1, L_2, L_3 ,*

$$p = \frac{1}{2 \sum_i L_i^2} \sum_{k=1}^3 L_k^2 g\left(\frac{L_i^2}{L_k^2}, \frac{L_j^2}{L_k^2}\right), \quad g(\alpha, \beta) = \mathbb{E}[(1 - (\alpha U_1 + \beta U_2)^2)_+],$$

Date: Preprint, version 1, 8 October 2026. Not peer reviewed.

2020 Mathematics Subject Classification. 60D05; 52A22; 60G55.

Key words and phrases. random polygons, smallest enclosing copy, Poisson limit, linear programming, Mecke equation, formal verification.




FIGURE 1. Forty uniform points in an equilateral triangle (solid). The smallest enclosing equilateral triangle (dashed) is rotated by about 0.08 radians and pokes out across the right side, near the lower right corner, although it is smaller than the outer triangle.

where $\{i, j, k\} = \{1, 2, 3\}$ and U_1, U_2 are independent and uniform on $[-1, 1]$. In particular $p = 13/48$ for the equilateral triangle, $7/24$ for the right isosceles triangle and $29/96$ for the 30-60-90 triangle; $p \rightarrow 1/3$ for thin right triangles and for needle-shaped isosceles triangles (apex angle tending to 0), and $p \rightarrow 25/72$ for flat isosceles triangles (apex angle tending to π).

Corollary 3 (Regular polygons). *For the regular q -gon,*

$$p_q = q \tan \frac{\pi}{q} \mathbb{E}[\text{Vol}(T) \mathbf{1}\{0 \in \text{int } T\}] + \mathbf{1}\{q \text{ even}\} \frac{8}{3q^2},$$

where T is the tetrahedron spanned by four independent points, each chosen by picking one of the q vertical edges of the prism $P_q \times [-1, 1]$ uniformly at random (P_q the regular q -gon inscribed in the unit circle) and a uniform height on it. In particular $p_3 = 13/48$ and $p_4 = 1/4$ (Section 6), and numerically p_q is close to 0.146 for large q (Table 1).

Conjecture 4. *Among all triangles, p is minimised by the equilateral triangle; the range of p over triangles is $[13/48, 25/72]$.*

The formula of Corollary 2 is explicit enough to evaluate on a fine grid: over 28,800 triangle shapes the minimum is attained at the equilateral triangle and p increases away from it (Section 8).

The limit model is a Poisson process along each side, as in Groeneboom's study of convex hulls of uniform samples from a polygon [1, 2], but at a different scale: the relevant points lie within distance $O(1/n)$ of the sides along their whole length, not near the vertices. Uniform samples from polygons are also the setting of Sylvester's problem, the probability that the n points are in convex position, whose asymptotics Morin obtained for regular [5] and for general convex polygons [6]; that event depends on the whole sample, whereas E_n depends only on the points within distance $O(1/n)$ of the boundary.

2. THE LIMIT MODEL

Fix the polygon K , scaled to area 1, with sides $1, \dots, m$. Side i has outward unit normal u_i , support number h_i (with the origin inside K), tangent $t_i = R_{90^\circ} u_i$, and is the set of points x with

$x \cdot u_i = h_i$ and $s = x \cdot t_i \in [a_i, b_i]$; its length is $L_i = b_i - a_i$. Two identities are used repeatedly:

$$(1) \quad \sum_i L_i u_i = 0, \quad \sum_i L_i h_i = 2, \quad \sum_i L_i b_i = \frac{1}{2} \sum_i L_i^2 = - \sum_i L_i a_i.$$

The last holds because $L_i(a_i + b_i)/2 = (b_i^2 - a_i^2)/2$ and $b_i^2 - a_i^2 = |v_{i+1}|^2 - |v_i|^2$ telescopes over the vertices v_i .

Heuristic. A point x at depth $d = h_i - x \cdot u_i$ below side i and position $s = x \cdot t_i$ has support $x \cdot R_\theta u_i = h_i - d + \theta s + O(\theta^2)$ in the rotated normal direction. A copy $(1 + \varepsilon)R_\theta K + c$ has support $(1 + \varepsilon)h_i + c \cdot u_i + O(\theta|c|)$ in the same direction, and its vertex at the end of side i pokes out across the line of side i unless $\varepsilon h_i + c \cdot u_i \leq \min(\theta a_i, \theta b_i)$. With $\Theta = n\theta$, $C = nc$, $D = nd$ and ε replaced by $n\varepsilon$, all quantities are of order one.




FIGURE 2. The limit model on one side, drawn with depth D downwards. The copy's side corresponds to the solid line $D = \Theta s - H_i$; every point must lie on or below it (deeper), which is $H_i \geq M_i(\Theta)$, and the marked point is tight. The copy fits at the end a_i when its line lies on or below the dashed corner line $D = \Theta(s - a_i)$; here it lies above, so the copy pokes out at that corner.

Definition. Let $\Pi_1, \dots, \Pi_m$ be independent Poisson processes of intensity 1 on the strips $[a_i, b_i] \times [0, \infty)$, and $M_i(\Theta) = \sup_{(s,D) \in \Pi_i} (\Theta s - D)$. The *limit LP* is

minimise ε over $(\varepsilon, C, \Theta) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}$ subject to $H_i := \varepsilon h_i + C \cdot u_i \geq M_i(\Theta)$ ($i = 1, \dots, m$), and E is the event that some optimal (ε, C, Θ) satisfies $H_i \leq \min(\Theta a_i, \Theta b_i)$ for all i . The model is scale free: changing the intensity rescales $D, \varepsilon, C, \Theta$ together and leaves E unchanged.

Lemma 5. *Almost surely each M_i is finite, convex and piecewise linear, and the optimal set Z^* of the limit LP is nonempty and compact.*

Proof. Each strip contains only finitely many points with $D \leq T$ for every T . On $|\Theta| \leq B$, fix any point (s_0, D_0) of Π_i ; since $|s| \leq S_i := \max(|a_i|, |b_i|)$, every point with $D > D_0 + 2BS_i$ has $\Theta s - D < \Theta s_0 - D_0$ for all such Θ , so only the finitely many points with $D \leq D_0 + 2BS_i$ can contribute, and M_i is a maximum of finitely many affine functions on any bounded Θ -interval. The point $z = 0$ is feasible, since $M_i(0) = -\inf D \leq 0$, so $\varepsilon^* \leq 0$. By (1), $\sum_i L_i H_i = 2\varepsilon$ for every (ε, C, Θ) , so feasibility gives $2\varepsilon \geq F(\Theta) := \sum_i L_i M_i(\Theta)$. Fix $\delta > 0$ with $\delta \text{per}(K) < \frac{1}{4} \sum_i L_i^2$ and, on each side, a point with $s \geq b_i - \delta$ and a point with $s \leq a_i + \delta$. For $\Theta > 0$ the first kind give $F(\Theta) \geq \Theta(\sum_i L_i b_i - \delta \text{per}(K)) - O(1) \geq \frac{\Theta}{4} \sum_i L_i^2 - O(1)$ by (1), and for $\Theta < 0$ the second kind give the same with $|\Theta|$. Hence $\{\varepsilon \leq 0\}$ forces $|\Theta|$ bounded, and ε is bounded below there. For such (ε, Θ) each $C \cdot u_i = H_i - \varepsilon h_i$ is bounded below; since $\sum_i L_i C \cdot u_i = 0$ with all $L_i > 0$, each is also bounded above, and the u_i span $\mathbb{R}^2$, so C is bounded. The feasible set intersected with $\{\varepsilon \leq 0\}$ is therefore closed, bounded and nonempty, and the minimum of ε over it is attained on a compact set. $\square$

3. CONVERGENCE

The proof compares the finite problem with the limit model on the same marked points near the boundary. Three facts make the comparison exact rather than approximate. First, in tangent coordinates (Lemma 6) the constraint that a copy contains a sample point is exactly the constraint of the limit LP; the nonlinearity moves into the objective, the physical scale of the copy. Second, at a nondegenerate optimum of the limit LP the tilt is controlled by the excess of ε , so for large n the physical scale and ε have the same minimisers over the common feasible set (Lemma 8). Third, the fit condition of the finite problem is that of the limit model plus a term of order $1/n$, which cannot change the outcome when the limit optimum fits strictly or fails strictly, and almost surely one of the two happens (Lemma 7). The probabilistic input is a Poisson approximation of the points near the boundary in total variation, uniformly over events (Step 3).

Notation. The event E_n is invariant under similarities, so K has area 1; we list its vertices counterclockwise in strictly convex position (a vertex at which the boundary is straight can be deleted). Fix r, R_K with $B(0, r) \subseteq K \subseteq B(0, R_K)$, and write $S_K = \max_i \max(|a_i|, |b_i|)$, $h_{\max} = \max_i h_i$, $Q = \sum_i L_i b_i = \frac{1}{2} \sum_i L_i^2$ and $\text{per} = \sum_i L_i$. For $x \in K$ and a side i let $d_i(x) = h_i - x \cdot u_i \geq 0$ be its depth and $s_i(x) = x \cdot t_i$ its position. Throughout, $q = 1/n$.

Tangent coordinates. For $z = (\varepsilon, C, \Theta) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}$ with $1 + q\varepsilon > 0$ let

$$\sigma_{q,z}(x) = \frac{1}{1 + q^2\Theta^2} T_{q\Theta}((1 + q\varepsilon)x + qC), \quad T_\tau(w) = w + \tau R_{90^\circ} w.$$

Since $T_\tau = \sqrt{1 + \tau^2} R_{\arctan \tau}$, the map $\sigma_{q,z}$ is a similarity with rotation angle $\arctan(q\Theta)$ and scale

$$\lambda_q(z) = \frac{1 + q\varepsilon}{\sqrt{1 + q^2\Theta^2}},$$

and every similarity with positive scale and rotation angle in $(-\pi/2, \pi/2)$ is $\sigma_{q,z}$ for exactly one such z . We call $\sigma_{q,z}(K)$ the copy z .

Lemma 6. *Let $1 + q\varepsilon > 0$ and write $H_i = \varepsilon h_i + C \cdot u_i$.*

(a) *A point x lies in the copy z if and only if, for every side i ,*

$$(2) \quad x \cdot u_i + q\Theta x \cdot t_i \leq (1 + q\varepsilon)h_i + qC \cdot u_i.$$

For a point at depth $d_i(x) = D/n$ and position $s = s_i(x)$ this is $D \geq \Theta s - H_i$, the constraint of the limit LP.

(b) *There is $r_K > 0$, depending only on the angles of K , such that if $|q\Theta| \leq r_K$ the copy z lies in K if and only if, for every side i and $s \in \{a_i, b_i\}$,*

$$\phi_{q,i,s}(z) := \Theta s - H_i + q(\varepsilon\Theta s + \Theta C \cdot t_i + \Theta^2 h_i) \geq 0.$$

At $q = 0$ this is the condition $H_i \leq \min(\Theta a_i, \Theta b_i)$ that defines E .

Proof. (a) Since $T_{-\tau}T_\tau = (1 + \tau^2) \text{id}$, the inverse of $\sigma_{q,z}$ is $x \mapsto (1 + q\varepsilon)^{-1}(T_{-q\Theta}x - qC)$, and $y \in K$ if and only if $y \cdot u_i \leq h_i$ for all i . As $(R_{90^\circ}x) \cdot u_i = -x \cdot t_i$, we have $T_{-\tau}x \cdot u_i = x \cdot u_i + \tau x \cdot t_i$, which gives (2). Substituting $x \cdot u_i = h_i - D/n$ and $x \cdot t_i = s$ and multiplying by n gives $D \geq \Theta s - H_i$. (b) The copy is a convex polygon whose sides are those of K turned by $\arctan(q\Theta)$. If this angle is less than half the least exterior angle of K , the support of the copy in the direction u_i is attained at an endpoint of its side i , so the copy lies in K if and only if both endpoints of its side i satisfy $y \cdot u_i \leq h_i$, for every i . The endpoint $h_i u_i + s t_i$ of side i is mapped to y with $(1 + q^2\Theta^2) y \cdot u_i = (1 + q\varepsilon)h_i + qC \cdot u_i - q\Theta((1 + q\varepsilon)s + qC \cdot t_i)$, and $y \cdot u_i \leq h_i$ is $\phi_{q,i,s}(z) \geq 0$ after dividing by q . $\square$

So the finite problem and the limit LP share their constraints. The finite problem minimises $\lambda_q(z)$, which is $1 + q\varepsilon + O(q^2)$ on bounded sets, and its fit condition is $\phi_q \geq 0$, which is the limit condition up to $O(q)$.

Proof of Theorem 1. Step 0: the rotation localises. Let $\Lambda(S, \theta)$ be the least λ with $S \subseteq \lambda R_\theta K + c$ for some c . Two facts about it. First, if $S' \subseteq S + B(0, \delta)$ then $\Lambda(S', \theta) \leq \Lambda(S, \theta) + \delta/r$, because $B(0, \delta) \subseteq (\delta/r)R_\theta K$ and $\lambda A + \mu A = (\lambda + \mu)A$ for convex A . Second, since $|R_{\theta'}x - R_\theta x| \leq R_K|\theta - \theta'|$ on K , we have $R_{\theta'}K \subseteq R_\theta K + B(0, R_K|\theta - \theta'|) \subseteq (1 + R_K|\theta - \theta'|/r)R_\theta K$, so $\theta \mapsto \Lambda(S, \theta)$ is continuous. Let G_K be the finite group of angles g with $R_g K = K + t_g$ for some t_g . If $K \subseteq \lambda R_\theta K + c$ then $1 \leq \lambda^2$ by area, and $\lambda = 1$ forces $K = R_\theta K + c$ (a convex body strictly inside another has strictly smaller area), so $\Lambda(K, \theta) \geq 1$ with equality only on G_K . Fix $\eta > 0$ with $\tan \eta \leq \min(r_K, Q/4)$ and smaller than half the least gap between elements of G_K ; by continuity and compactness there is $\beta > 0$ with $\Lambda(K, \theta) \geq 1 + \beta$ whenever θ is at distance at least η from G_K . Cover K by finitely many balls $B(y_k, \delta/2)$, $y_k \in K$, with $\delta = \beta r/2$; each meets K in positive area, so with probability tending to 1 each contains a sample point, and then $K \subseteq S + B(0, \delta)$. On that event $\Lambda(S, \theta) \geq 1 + \beta/2 > 1 \geq \Lambda(S, 0)$ for every such θ , so every smallest copy has its rotation within η of G_K . For $g \in G_K$ we have $\lambda R_{\theta+g}K + c = \lambda R_\theta K + c + \lambda R_\theta t_g$, so rotations θ and $\theta + g$ give the same family of sets. Hence every smallest copy is a copy z with $|q\Theta| \leq \tan \eta$, and it has scale $\lambda_q(z) \leq 1$ because K itself contains the sample.

Step 1: witnesses confine the copies. Let $\delta_K = Q/(2\text{per})$ and, for each side, take the windows $W_i^+ = \{x : s_i(x) \in [b_i - w_i, b_i - w_i/2]\}$ and $W_i^- = \{x : s_i(x) \in [a_i + w_i/2, a_i + w_i]\}$ near its two ends, with $w_i = \min(\delta_K, L_i)/2$. Call points of the sample in $W_i^\pm$ at depth $d_i \leq y/n$ *witnesses*. The part of $W_i^\pm$ at depth at most y/n has area $w_i y/(2n)$, so the probability that some window has no witness is at most $2m \max_i (1 - w_i y/(2n))^n \leq 2me^{-\min_i w_i y/2}$, small for large y , uniformly in n .

Suppose every side has witnesses $x_i^\pm$ at scaled depths $D_i^\pm = nd_i(x_i^\pm) \leq M$, and let z satisfy their constraints, with $\lambda_q(z) \leq 1$ and $|q\Theta| \leq Q/4$. By Lemma 6(a), $D_i^+ \geq \Theta s_i^+ - H_i$. Multiply by L_i and sum: the terms in C cancel because $\sum_i L_i u_i = 0$, and $\sum_i L_i h_i = 2$, so $\Theta \sum_i L_i s_i^+ \leq 2\varepsilon + \sum_i L_i D_i^+$. Here $\sum_i L_i s_i^+ \geq \sum_i L_i (b_i - \delta_K) = Q/2$ by (1), and $\lambda_q(z) \leq 1$ gives $1 + q\varepsilon \leq \sqrt{1 + q^2\Theta^2} \leq 1 + q^2\Theta^2/2$, so $\varepsilon \leq q\Theta^2/2 \leq Q|\Theta|/8$. For $\Theta > 0$ this yields $\Theta Q/4 \leq M$ per, and the witnesses x_i^- give the same bound for $\Theta < 0$. With $|\Theta|$ bounded, the same sum bounds ε below, each $C \cdot u_i = H_i - \varepsilon h_i \geq \Theta s_i^+ - D_i^+ - \varepsilon h_i$ is bounded below, hence also above since $\sum_i L_i C \cdot u_i = 0$, and two of the u_i are not parallel, so C is bounded. So there is $R = R(M, K)$ such that every such z lies in the box $B_R = \{|\varepsilon|, |C_1|, |C_2|, |\Theta| \leq R\}$, for every n . The same argument, with $\varepsilon \leq 0$ in place of $\lambda_q(z) \leq 1$, shows that every z feasible for the witnesses in the limit LP with $\varepsilon \leq 0$ lies in B_R .

Step 2: only the boundary window matters. There is $T_c = T_c(R, K)$ with $\Theta s_i(x) - H_i \leq T_c$ for every $x \in K$, $z \in B_R$ and side i . So, by Lemma 6(a), a point with $nd_i(x) > T_c$ satisfies the side- i constraint of every copy in B_R . Fix $T \geq T_c$. Let $\mathcal{O}_n$ be the set of points of K within T/n of two sides; it has area $O(T^2/n^2)$, so with probability tending to 1 it contains no sample point. Mark each sample point x with $d_i(x) \leq T/n$ by $(i, s_i(x), nd_i(x))$, and let μ_n be the resulting configuration. For a configuration μ let $\mathcal{G}(\mu) \subseteq B_R$ be the set of copies in B_R satisfying the constraints $D \geq \Theta s - H_i$ of all its marks, and define the events

$$\begin{aligned} F_n(\mu) : & \mu \text{ has witnesses at depth } \leq M, \text{ and some minimiser of } \lambda_q \text{ over } \mathcal{G}(\mu) \\ & \text{satisfies } \phi_q \geq 0, \\ F(\mu) : & \mu \text{ has witnesses at depth } \leq M, \text{ and some minimiser of } \varepsilon \text{ over } \mathcal{G}(\mu) \\ & \text{satisfies } H_i \leq \min(\Theta a_i, \Theta b_i) \text{ for all } i. \end{aligned}$$

Let $\mathcal{A}_n$ be the event that the δ -net of Step 0 is hit, that every side has witnesses at depth at most $y/n \leq M/n$, and that no sample point lies in $\mathcal{O}_n$. On $\mathcal{A}_n$, and for $n > R/r_K$ (so that $1 + q\varepsilon > 0$ and $|q\Theta| \leq r_K$ on B_R), E_n holds if and only if $F_n(\mu_n)$ holds. Indeed, by Step 0 every smallest copy is a copy z with $|q\Theta| \leq \tan \eta$ and $\lambda_q(z) \leq 1$, so by Step 1 it lies in B_R , and the smallest copies are the minimisers of λ_q over the copies in B_R containing the sample; a copy in B_R contains the

sample if and only if it lies in $\mathcal{G}(\mu_n)$, because the constraints of a point for sides it is not within T/n of are slack (Step 2) and the others are exactly the constraints of its marks (Lemma 6(a)); and the fit condition is $\phi_q \geq 0$ by Lemma 6(b). Both events are measurable: “some minimiser of a continuous function over a compact set lies in a closed set” is a projection along a compact factor.

Step 3: the boundary window is Poisson. Let Π^T be the limit model truncated at depth T : independent Poisson processes of intensity 1 on the rectangles $[a_i, b_i] \times [0, T]$. Then

$$\sup_F |\mathbb{P}(F(\mu_n)) - \mathbb{P}(F(\Pi^T))| \rightarrow 0,$$

the supremum over all measurable events F of configurations. The set of points of K within T/n of the sides has area per $T/n + O(T^2/n^2)$, so the number of sample points in it is binomial with mean per $T + O(1/n)$, and its law tends in total variation to the Poisson law with mean per T (the law of rare events). Given this number, the points are independent and uniform on the window, and the marking map has Jacobian n on the strip along each side; the strips differ from the rectangles $[a_i, b_i] \times [0, T]$, in the marked coordinates, by triangles at the ends and by the overlaps $\mathcal{O}_n$, of total area $O(T^2/n)$. So the conditional law of the marks is within $O(1/n)$ in total variation of the normalised Lebesgue measure on the rectangles, the mark law of Π^T . Mixing over the number of points gives the bound, uniformly in F . Consequently, if events F_n and F satisfy $F_n(\Pi^T) \Leftrightarrow F(\Pi^T)$ for all large n almost surely, then $\mathbb{P}(F_n(\mu_n)) \rightarrow \mathbb{P}(F(\Pi^T))$: the difference $\mathbb{P}(F_n(\mu_n)) - \mathbb{P}(F_n(\Pi^T))$ tends to 0 by the bound, and $\mathbb{P}(F_n(\Pi^T)) \rightarrow \mathbb{P}(F(\Pi^T))$ by dominated convergence.

Step 4: stability. By Lemma 8 below, almost surely $F_n(\Pi^T) \Leftrightarrow F(\Pi^T)$ for all large n . So $\mathbb{P}(F_n(\mu_n)) \rightarrow \mathbb{P}(F(\Pi^T))$.

Step 5: removing the cutoff. Let E^T be the event E for Π^T : some copy minimising ε over all copies feasible for Π^T satisfies $H_i \leq \min(\Theta a_i, \Theta b_i)$. If Π^T has witnesses at depth at most M , every feasible copy with $\varepsilon \leq 0$ lies in B_R (Step 1), and 0 is feasible, so the minimisers over all feasible copies and over $\mathcal{G}(\Pi^T)$ coincide, and $F(\Pi^T) \Leftrightarrow E^T$. The probability that Π^T has no witness in some window at depth at most $M \leq T$ is at most $2me^{-\min_i w_i M/2}$, by the void probability of Poisson processes (each window has area $w_i M/2$ in the marked coordinates). Finally $\mathbb{P}(E^T) \rightarrow \mathbb{P}(E) = p(K)$ as $T \rightarrow \infty$: Π^T is the restriction of the limit model to depths at most T , and by Lemma 5 and the witness bound the optimal set of the limit model lies in a bounded box whose lines are below a finite depth, almost surely, so $E^T = E$ for all large T .

Assembly. Given $\epsilon > 0$ choose y so that the sample has witnesses at depth at most y/n with probability at least $1 - \epsilon$ for all n (Step 1), then $M \geq y$ with $2me^{-\min_i w_i M/2} \leq \epsilon$, then R, η, δ and $T \geq T_c$ with $|\mathbb{P}(E^T) - p(K)| \leq \epsilon$ (Steps 0, 1, 2 and 5). For large n , $\mathbb{P}(\mathcal{A}_n^c) \leq 2\epsilon$ (Steps 0, 1, 2), so $|\mathbb{P}(E_n) - \mathbb{P}(F_n(\mu_n))| \leq 2\epsilon$; then $|\mathbb{P}(F_n(\mu_n)) - \mathbb{P}(F(\Pi^T))| \leq \epsilon$ (Steps 3 and 4), $|\mathbb{P}(F(\Pi^T)) - \mathbb{P}(E^T)| \leq \epsilon$ (Step 5) and $|\mathbb{P}(E^T) - p(K)| \leq \epsilon$. So $|\mathbb{P}(E_n) - p(K)| \leq 5\epsilon$ for all large n . $\square$

Nondegeneracy and stability. Write $a_j = (h_i, u_i, -s_j)$ for the gradient of the limit constraint $g_j(z) = \varepsilon h_i + C \cdot u_i - \Theta s_j + D_j \geq 0$ of a point j on side i , and $\tau = (0, t_k, 0)$. A feasible z is optimal for the limit LP if and only if $e_1 = \sum_{j \in B} \lambda_j a_j$ for some $\lambda_j \geq 0$, where B is the set of constraints tight at z .

Lemma 7. *For the limit model, and for Π^T at each cutoff T , the following hold almost surely whenever the optimal set Z^* is nonempty.*

- (a) *Either (V) $Z^* = \{z^*\}$, exactly four constraints are tight at z^* , their gradient matrix A_B is invertible and $A_B^{-\top} e_1$ is positive; or (S) exactly three constraints are tight on Z^* , from points j_1, j_2 on a side k and j_3 on the parallel side $\bar{k}$ with $s_{j_1} < -s_{j_3} < s_{j_2}$, and $Z^* = \{z^* + y\tau : \ell \leq y \leq r\}$ with $\ell < r$.*
- (b) *There is D_0 such that every feasible z satisfies $|\Theta(z) - \Theta^*| \leq D_0(\varepsilon(z) - \varepsilon^*)$, where Θ^* is the tilt of Z^* (constant on Z^*) and ε^* the optimal value.*

- (c) *If some $z \in Z^*$ satisfies $H_i \leq \min(\Theta a_i, \Theta b_i)$ for all i , then some $z \in Z^*$ satisfies these inequalities strictly.*

Proof. Every failure is contained in an event of the following form: there are distinct points $x_1, \dots, x_r$ of the process and an index l such that, for fixed values of all the other coordinates, the depth of x_l satisfies a relation $\alpha D_l + \beta = 0$ with $\alpha \neq 0$, where α, β are measurable functions of the other coordinates of $x_1, \dots, x_r$ and of the rest of the process. By the multivariate Mecke equation [3, Thm. 4.4] the probability of such an event is at most the integral, over $(x_1, \dots, x_r)$, of the probability that the relation holds when $x_1, \dots, x_r$ are added to the process; for fixed other coordinates the relation picks out one value of D_l , a Lebesgue null set, so the integral vanishes. We list the failures.

If the rows of a tight set are linearly dependent, the depths of its points satisfy an affine relation whose coefficient on some depth is nonzero. So almost surely the tight rows are independent at every z : at most four constraints are tight, and never three from one side (their rows span only two dimensions). Let $e_1 = \sum_{j \in B} \lambda_j a_j$ with all $\lambda_j > 0$ and B minimal, so $\sum \lambda_j h_{i(j)} = 1$, $\sum \lambda_j u_{i(j)} = 0$ and $\sum \lambda_j s_j = 0$. If $|B| \leq 2$, the condition $\sum \lambda_j u_{i(j)} = 0$ forces one point on each of two antiparallel sides with equal weights, and then $\sum \lambda_j s_j = 0$ says $s_j = -s_{j'}$, a null event (apply the argument to a position instead of a depth). If $|B| = 3$: three pairwise non-parallel sides fix λ up to scale through the u_i , and $\sum \lambda_j s_j = 0$ is a nontrivial linear equation in the positions, a null event; two points on a side k and one on a side not parallel to it, or one point each on $k, \bar{k}$ and a third side, are incompatible with $\sum \lambda_j u_{i(j)} = 0$ and $\lambda > 0$; two points on k and one on $\bar{k}$ is case (S), and then $\lambda_{j_1} + \lambda_{j_2} = \lambda_{j_3}$ together with $\sum \lambda_j s_j = 0$ forces $s_{j_1} < -s_{j_3} < s_{j_2}$ (equality is a null event). If $|B| = 4$ the rows are independent and $\lambda = A_B^{-\top} e_1$ is positive by minimality: case (V). In case (S) every row in B is orthogonal to τ , so Z^* contains a segment in direction τ , bounded because sides with $u_j \cdot t_k > 0$ and with $u_j \cdot t_k < 0$ both exist; its endpoints are where one further constraint becomes tight. A further tight point at z^* in case (V), or a fifth constraint tight at an endpoint in case (S), is an affine relation in that point's depth with coefficient 1, so null. If no optimal point carries a set of type (S), Z^* is a single point, since a face of positive dimension would contain a point whose tight rows span fewer than four dimensions. This proves (a).

For (b): in case (V), $e_1 = A_B^\top \lambda$ and $g_B(z^*) = 0$ give $\varepsilon(z) - \varepsilon^* = \sum_{j \in B} \lambda_j g_j(z) \geq \lambda_{\min} |g_B(z)|_1$ for feasible z , and $|z - z^*| \leq \|A_B^{-1}\| |g_B(z)|_1$, so $|\Theta - \Theta^*| \leq (\|A_B^{-1}\| / \lambda_{\min})(\varepsilon - \varepsilon^*)$. In case (S) the same identity with the three multipliers controls the coordinates of z other than the one along τ , and Θ is among them.

For (c), let $\mu(z) = \min_i (\min(\Theta a_i, \Theta b_i) - H_i(z))$, concave and piecewise affine, so that the hypothesis is $\max_{Z^*} \mu \geq 0$; it suffices that $\max_{Z^*} \mu \neq 0$ almost surely. The maximum is attained at z^* , at an endpoint of Z^* , or at a kink of μ on Z^* , and on the event that B is the tight set it is in each case an affine function, with nonzero coefficient, of one depth: z^* is affine in D_B with invertible linear part in case (V), and $(\varepsilon, \Theta, C \cdot u_k)$ is affine in D_B with Jacobian $(s_{j_2} - s_{j_1}) w_k \neq 0$ in case (S), where $w_k = h_k + h_{\bar{k}}$. For a kink: along the segment each piece of μ has the form $\alpha_i + c_i y$ with α_i affine in $(\varepsilon, \Theta, C \cdot u_k)$; two pieces from the same side cross only where $\Theta = 0$, a null event, and two pieces from sides $i \neq i'$ cross at the value $(c_i \alpha_{i'} - c_{i'} \alpha_i) / (c_i - c_{i'})$, which vanishes identically only if (h_i, u_i) is proportional to $(h_{i'}, u_{i'})$, impossible for distinct sides of a convex polygon. So each case is null. $\square$

Lemma 8. *Fix T, M and R as in the proof of Theorem 1. Almost surely, $F_n(\Pi^T) \Leftrightarrow F(\Pi^T)$ for all large n .*

Proof. Work on the almost sure event of Lemma 7, with all depths nonnegative, and assume Π^T has witnesses at depth at most M (otherwise both events fail). The set $\mathcal{G} = \mathcal{G}(\Pi^T)$ is compact and contains 0. Let $\varepsilon^* = \min_{\mathcal{G}} \varepsilon$ and $Z^* = \{z \in \mathcal{G} : \varepsilon(z) = \varepsilon^*\}$, compact. By Step 1 every feasible

copy with $\varepsilon \leq 0$ lies in B_R , so Z^* is the optimal set of the limit LP for Π^T , and Lemma 7 applies to it.

The minimisers coincide. Let $z \in \mathcal{G}$ and $z^* \in Z^*$. If $\varepsilon(z) = \varepsilon^*$ then $\Theta(z) = \Theta^*$ by Lemma 7(b), so $\lambda_q(z) = \lambda_q(z^*)$. If $\varepsilon(z) > \varepsilon^*$, write $\Delta = \varepsilon(z) - \varepsilon^*$; then $|\Theta(z) - \Theta^*| \leq D_0 \Delta$, and since the derivative of $\tau \mapsto (1 + \tau^2)^{-1/2}$ has absolute value at most $|\tau|$, and $|q\Theta| \leq qR$ and $|\varepsilon^*| \leq R$ on $\mathcal{G}$,

$$\lambda_q(z) - \lambda_q(z^*) \geq \frac{q\Delta}{\sqrt{1 + q^2 R^2}} - (1 + qR)qR \cdot qD_0 \Delta,$$

which is positive once $qRD_0(1 + qR)\sqrt{1 + q^2 R^2} < 1$. So for all large n the minimisers of λ_q over $\mathcal{G}$ are exactly Z^* .

The fit conditions agree. If some $z \in Z^*$ satisfies $H_i \leq \min(\Theta a_i, \Theta b_i)$ for all i , then by Lemma 7(c) some $z' \in Z^*$ satisfies them strictly, so $\phi_{0,i,s}(z') > 0$ for all i and $s \in \{a_i, b_i\}$, and by continuity $\phi_{q,i,s}(z') > 0$ for all small q . If no $z \in Z^*$ satisfies them, then no $z \in Z^*$ satisfies $\phi_q \geq 0$ for small q : otherwise there would be $q_k \rightarrow 0$ and $z_k \in Z^*$ with $\phi_{q_k}(z_k) \geq 0$, and a limit point of (z_k) in the compact set Z^* would satisfy $\phi_0 \geq 0$, by joint continuity of ϕ in (q, z) . In both cases $F_n(\Pi^T) \Leftrightarrow F(\Pi^T)$ for all large n . $\square$

4. COMPUTING THE LIMIT

Lemma 9 (void probability). *If $H_i \leq \min(\Theta a_i, \Theta b_i)$ for all i , the probability that no point of any Π_i satisfies $D < \Theta s - H_i$ (that is, lies strictly above the line in the orientation of Figure 2) is $\exp(2\varepsilon)$, where ε is defined by $H_i = \varepsilon h_i + C \cdot u_i$.*

Proof. Under the hypothesis each line is nonnegative on $[a_i, b_i]$, so the region between $D = 0$ and the line (above it in Figure 2) has area $\int_{a_i}^{b_i} (\Theta s - H_i) ds = \Theta(b_i^2 - a_i^2)/2 - L_i H_i$. Summing over i , the first terms telescope to 0 and $\sum L_i H_i = \varepsilon \sum L_i h_i + C \cdot \sum L_i u_i = 2\varepsilon$ by (1). $\square$

The telescoping is the heart of the computation: the rotation costs nothing on average, and only the scale enters.

What remains is to count optima. The limit LP has four unknowns, $z = (\varepsilon, C, \Theta)$, so a generic optimum is a vertex fixed by four tight points. When K has a pair of parallel sides there is a second possibility: an optimum fixed along a segment by three tight points, two on one side and one on the side parallel to it. Theorem 10 has one term for each case. In the vertex term, $V(K)$ integrates over scale and tilt the area of translations for which the copy fits, weighted by the void probability of Lemma 9, and $T_{\text{nd}}(K)$ is the density of four-point configurations that can be optimal. The quantities w_k , S_k and σ_k play the same roles in the segment term. For triangles everything becomes explicit (Section 5).

For $t \geq 0$ and $\Theta \in \mathbb{R}$ let

$$R(t, \Theta) = \{C \in \mathbb{R}^2 : C \cdot u_i \leq t h_i + \min(\Theta a_i, \Theta b_i) \text{ for all } i\},$$

$$V(K) = \int_{\mathbb{R}} \int_0^\infty e^{-2t} \text{area } R(t, \Theta) dt d\Theta.$$

Generic optima of the limit LP are vertices determined by four tight points. For a choice of sides $i_1, \dots, i_4$ and positions $s_1, \dots, s_4$, let A be the 4×4 matrix with rows $(h_{i_r}, u_{i_r}, -s_r)$, and let

$$T_{\text{nd}}(K) = \frac{1}{24} \sum_{i_1, \dots, i_4} \int \cdots \int |\det A| \mathbf{1}\{e_1 \in \text{int cone}(\text{rows of } A)\} ds_1 \cdots ds_4,$$

the integrals over the sides, where tuples containing two indices of a side k and one of a side parallel to it are excluded. (The cone condition says that the origin lies inside the tetrahedron

with vertices $(u_{i_r}, -s_r)/h_{i_r}$, which sit over the vertices of the polar polygon of K .) For a pair of parallel sides $k, \bar{k}$ let $w_k = h_k + h_{\bar{k}}$, $S_k = \sum_{j: u_j \cdot t_k > 0} L_j u_j \cdot t_k$ and

$$\sigma_k = \int_{I_k} \frac{1}{2}(s - a_k)(b_k - s)(b_k - a_k) ds,$$

where I_k is the intersection of side k with the projection of side $\bar{k}$ (in the coordinate $x \cdot t_k$).

Theorem 10. *For a convex polygon K of area one,*

$$p(K) = V(K) T_{\text{nd}}(K) + \sum_{(k, \bar{k})} \sigma_k \int_{\mathbb{R}} \int_0^\infty e^{-2t} w_k [\text{width}_{u_k} R(t, \Theta) + S_k \text{area } R(t, \Theta)] dt d\Theta,$$

the sum over ordered pairs of parallel sides (empty when K has no parallel sides).

In words, the first term is the vertex case, a fitting volume times a configuration density, and the sum is the segment case, which occurs only when K has parallel sides.

Proof. By Lemma 7 the limit LP is almost surely nondegenerate, so E splits into the disjoint events E_v (vertex optimum) and $E_s^{(k)}$ (segment optimum with the pair of tight points on side k and one on the parallel side $\bar{k}$). Write $\Pi = \bigcup_i \Pi_i$, viewed as a Poisson process on the disjoint union $\mathcal{S}$ of the strips, with intensity $ds dD$ on each.

Vertex optima. $\mathbb{P}(E_v) = \mathbb{E} \sum \mathbf{1}\{x_1, \dots, x_4 \text{ are the tight points and } E \text{ holds}\}$, the sum over sets of four distinct points of Π . By the multivariate Mecke equation [3, Thm. 4.4] this equals

$$\frac{1}{24} \int_{\mathcal{S}^4} \mathbb{P}(\{x_1, \dots, x_4\} \text{ is the tight set of the optimum of}$$

$$\Pi \cup \{x_1, \dots, x_4\}, \text{ and } E \text{ holds}) dx_1 \cdots dx_4,$$

the factor $1/24$ accounting for the orderings. Fix the sides i_r and positions s_r of the x_r , with $\det A \neq 0$. The map $(D_1, \dots, D_4) \mapsto z = (\varepsilon, C, \Theta)$ defined by making the four constraints tight, $D_r = \Theta s_r - \varepsilon h_{i_r} - C \cdot u_{i_r}$, is affine with Jacobian $|\det A|$. Given z , the four points form the tight set of an optimal vertex iff (a) the KKT condition $e_1 \in \text{int cone}(\text{rows of } A)$ holds, which depends only on the s_r , and (b) no point of Π violates a constraint at z . On the event E , which at z reads $C \in R(-\varepsilon, \Theta)$ (that is, $H_i \leq \min(\Theta a_i, \Theta b_i)$; the set $R(t, \cdot)$ is empty for $t < 0$ because $\sum L_i H_i = 2\varepsilon$), every line $D = \Theta s - H_i$ is nonnegative on its side, so the tight points have $D_r \geq 0$ and (b) has probability $e^{2\varepsilon}$ by Lemma 9. Hence

$$\mathbb{P}(E_v) = \frac{1}{24} \sum_{i_1, \dots, i_4} \int |\det A| \mathbf{1}\{\text{KKT}\} ds \times \int e^{2\varepsilon} \mathbf{1}\{C \in R(-\varepsilon, \Theta)\} d\varepsilon dC d\Theta = T_{\text{nd}}(K) V(K).$$

Tuples with two points on a side k and one on a parallel side $\bar{k}$ have the origin on a face of the tetrahedron (the three rows are linear combinations of $(h_k, u_k, 0)$, $(h_{\bar{k}}, -u_k, 0)$ and $(0, 0, 1)$), so they never give an interior cone point and are excluded at no cost.

Segment optima. Now apply the Mecke equation with three points: x_1, x_2 on side k with $s_1 < s_2$ (this ordering replaces the factor $1/2$) and x_3 on $\bar{k}$. The tight equations $D_r = \Theta s_r - H_k$ ($r = 1, 2$) and $D_3 = \Theta s_3 - H_{\bar{k}}$ determine $(\Theta, H_k, H_{\bar{k}})$ with Jacobian $s_2 - s_1$, and $H_k = \varepsilon h_k + w$, $H_{\bar{k}} = \varepsilon h_{\bar{k}} - w$ with $w = C \cdot u_k$ has Jacobian $h_k + h_{\bar{k}} = w_k$. KKT requires multipliers $\mu_1, \mu_2, \mu_3 > 0$ with $\mu_1 + \mu_2 = \mu_3$, $(\mu_1 + \mu_2)w_k = 1$ and $\mu_1 s_1 + \mu_2 s_2 + \mu_3 s_3 = 0$, which is possible iff $s_1 < -s_3 < s_2$: the third point projects between the first two (note $x \cdot t_{\bar{k}} = -x \cdot t_k$).

The remaining coordinate $y = C \cdot t_k$ does not enter the three tight constraints. For every other side j , $H_j(y) = \varepsilon h_j + w u_j \cdot u_k + y c_j$ with $c_j = u_j \cdot t_k$. The vertex constraints hold exactly for y in the chord $Y_E = [\alpha, \beta]$ of $R(-\varepsilon, \Theta)$ at $C \cdot u_k = w$. Feasibility of side j is $M_j(\Theta) \leq H_j(y)$: a lower bound on y if $c_j > 0$, an upper bound if $c_j < 0$. Let ℓ and r be the largest lower and the smallest

upper bound; they are independent. For $y \in Y_E$ all lines are nonnegative on their sides, so by the computation in Lemma 9

$$F(y) := \mathbb{P}(\ell \leq y) = \prod_{c_j > 0} e^{-\text{void}_j(y)}, \quad G(y) := \mathbb{P}(r \geq y) = \prod_{c_j < 0} e^{-\text{void}_j(y)},$$

with $\text{void}_j(y) = \Theta(b_j^2 - a_j^2)/2 - L_j H_j(y)$. Since $\sum_j L_j c_j = 0$ (and $c_k = c_{\bar{k}} = 0$), the product FG is constant on Y_E , and $F' = S_k F$. The segment optimum meets Y_E iff $\ell \leq \beta$, $r \geq \alpha$ and $\ell \leq r$, so

$$\mathbb{P}(\exists y \in Y_E \text{ feasible}) = F(\alpha)G(\alpha) + \int_{\alpha}^{\beta} G dF = FG(1 + S_k(\beta - \alpha)).$$

Multiplying by the void probabilities of sides k and $\bar{k}$ gives $e^{2\varepsilon}(1 + S_k|Y_E|)$ by Lemma 9. Integrating over w , $\int \mathbf{1}\{Y_E \neq \emptyset\} dw = \text{width}_{u_k} R$ and $\int |Y_E| dw = \text{area } R$; integrating over $s_1 < s_2$ and s_3 the factor $(s_2 - s_1)\mathbf{1}\{s_1 < -s_3 < s_2\}$ gives σ_k , because $\int_{s_1 < x < s_2} (s_2 - s_1) ds_1 ds_2 = \frac{1}{2}(x - a_k)(b_k - x)(b_k - a_k)$. Summing over ordered pairs $(k, \bar{k})$ proves the theorem. $\square$

5. TRIANGLES

For a triangle the map $(\varepsilon, C) \mapsto (H_1, H_2, H_3)$ is linear and invertible, so the LP becomes: minimise $\sum L_i H_i$ subject to $H_i \geq M_i(\Theta)$, that is, choose $H_i = M_i(\Theta)$ and minimise $F(\Theta) = \sum_i L_i M_i(\Theta)$, a convex piecewise linear function. At the minimiser one side carries two tight points $s_1 < s_2$ and the others one each, and optimality reads $L_k s_1 + \Sigma' \leq 0 \leq L_k s_2 + \Sigma'$ with $\Sigma' = \sum_{j \neq k} L_j s_j$. By Lemma 9 and the identity $\sum L_i a_i = -\frac{1}{2} \sum L_i^2$, the integral over Θ and the heights equals $1/(Q L_1 L_2 L_3)$ for each sign of Θ , where $Q = \frac{1}{2} \sum L_i^2$; the region R is a homothetic copy of K , and $V(K) = 1/\sum L_i^2$. The positions contribute

$$I_k = \frac{L_k}{2} \iint ((x - a_k)(b_k - x))_+ ds_i ds_j, \quad x = -\frac{L_i s_i + L_j s_j}{L_k},$$

so $p = 2 \sum_k I_k / (Q L_1 L_2 L_3)$. Writing x as the midpoint of side k plus a sum of two independent uniform variables of half-widths $L_i^2/(2L_k)$ and $L_j^2/(2L_k)$ gives Corollary 2.

For the equilateral triangle $\alpha = \beta = 1$, $U_1 + U_2$ has density $(2 - |w|)/4$, and $g(1, 1) = 2 \int_0^1 (1 - w^2)(2 - w)/4 dw = 13/24$, so $p = \frac{1}{6} \cdot 3 \cdot \frac{13}{24} = \frac{13}{48}$. If $\alpha + \beta \leq 1$ (the angle opposite side k is obtuse or right) the positive part is inactive and $g = 1 - (\alpha^2 + \beta^2)/3$.

6. REGULAR POLYGONS

Take the regular q -gon of area one, centred at the origin, with inradius r and half side $c = r \tan(\pi/q)$, so that $q \tan(\pi/q) r^2 = 1$. Its normals are $u_k = (\cos \frac{2\pi k}{q}, \sin \frac{2\pi k}{q})$, $h_k = r$, and every side has $[a_k, b_k] = [-c, c]$. We evaluate the three factors of Theorem 10; write P for the regular q -gon of inradius 1, which has area $q \tan(\pi/q)$.

The vertex term. Since $\min(\Theta a_k, \Theta b_k) = -c|\Theta|$, the region $R(t, \Theta)$ is ρP with $\rho = rt - c|\Theta|$ when $\rho > 0$, and is empty when $\rho < 0$. With $\alpha = c|\Theta|/r$,

$$V(K) = q \tan \frac{\pi}{q} \int_{\mathbb{R}} \int_{\alpha}^{\infty} e^{-2t} r^2 (t - \alpha)^2 dt d\Theta = q \tan \frac{\pi}{q} \frac{r^2}{4} \int_{\mathbb{R}} e^{-2c|\Theta|/r} d\Theta = \frac{q \tan(\pi/q) r^3}{4c}.$$

In $T_{\text{nd}}(K)$ substitute $s_r = -c y_r$ with $y_r \in [-1, 1]$. The rows of A become $(r, u_{i_r}, c y_r)$, so $|\det A| = rc |\det M|$, where M has rows $(1, P_r)$ with $P_r = (u_{i_r}, y_r)$ a point on the vertical edge over the vertex u_{i_r} of the prism $P_q \times [-1, 1]$ of Corollary 3; and $|\det M| = 6 \text{Vol}(T)$ for the tetrahedron T with vertices $P_1, \dots, P_4$. The cone condition says that 0 is a combination of the P_r with positive

weights, which when $\det M \neq 0$ means $0 \in \text{int } T$; when $\det M = 0$ both $\text{Vol}(T)$ and the interior of T vanish. Averaging over the q^4 choices of edges and the density $1/2$ of each height,

$$T_{\text{nd}}(K) = \frac{1}{24} c^4 \cdot 6rc \cdot 16q^4 \mathbb{E}[\text{Vol}(T) \mathbf{1}\{0 \in \text{int } T\}] = 4rc^5 q^4 \mathbb{E}[\text{Vol}(T) \mathbf{1}\{0 \in \text{int } T\}].$$

Since $r^4 c^4 q^4 = (r^2 q \tan(\pi/q))^4 = 1$, the vertex term is $V(K) T_{\text{nd}}(K) = q \tan(\pi/q) \mathbb{E}[\text{Vol}(T) \mathbf{1}\{0 \in \text{int } T\}]$.

The segment term. The normals u_k and $u_{\bar{k}} = -u_k$ are both present exactly when q is even, with $\bar{k} = k + q/2$, giving q ordered pairs, and none when q is odd. For each pair $w_k = 2r$, $I_k = [-c, c]$ and $\sigma_k = \int_{-c}^c \frac{1}{2}(c^2 - s^2) 2c ds = \frac{4}{3}c^4$. The region $R(t, \Theta) = \rho P$ has width 2ρ in the direction u_k and area $q \tan(\pi/q) \rho^2$, and $S_k = 2c \sum_{j=1}^{q/2-1} \sin(2\pi j/q) = 2c \cot(\pi/q) = 2r$. Hence

$$\int_{\mathbb{R}} \int_0^\infty e^{-2t} w_k [2\rho_+ + S_k q \tan \frac{\pi}{q} \rho_+^2] dt d\Theta = 2r \cdot \frac{r}{c} \left(\frac{2r}{4} + \frac{2r \cdot q \tan(\pi/q) r^2}{4} \right) = \frac{2r^3}{c},$$

and each ordered pair contributes $\frac{4}{3}c^4 \cdot \frac{2r^3}{c} = \frac{8}{3}r^3 c^3 = \frac{8}{3q^3}$, since $r^3 c^3 = (r^2 q \tan(\pi/q))^3 / q^3$. The q pairs give $8/(3q^2)$, which proves Corollary 3.

For $q = 3$, Corollary 2 gives $p_3 = 13/48$. For $q = 4$ only tuples with one point on each edge of the prism contribute to the expectation, and the integral factorises over the two pairs of opposite edges; it gives $\mathbb{E}[\text{Vol}(T) \mathbf{1}\{0 \in \text{int } T\}] = 1/48$, so $p_4 = 4 \cdot \frac{1}{48} + \frac{1}{6} = \frac{1}{4}$.

q	3	4	5	6	7	8	10	12	20	200
p_q	$\frac{13}{48}$	$\frac{1}{4}$	0.1732	0.1935	0.1585	0.1730	0.1632	0.1586	0.1506	0.1464

TABLE 1. Limit probabilities for regular q -gons (Monte Carlo evaluation of Corollary 3, standard error $\leq 3 \times 10^{-4}$).

The values are not monotone: apart from $q = 3, 4$ (where $p_3 > p_4$), the computed odd values $q = 5, 7$ lie below their even neighbours. They appear to settle near 0.146 for large q (we computed up to $q = 200$), and do not approach the disk value $1/2$. For the disk the enclosing circle has no rotational freedom and its size is governed by curvature; for a polygon with many sides the rotation still costs a corner poke of order $1/n$ while the side lengths shrink, and the two limits $n \rightarrow \infty$ and $q \rightarrow \infty$ do not commute.

7. FORMAL VERIFICATION

Every theorem and corollary above has been checked in the Lean 4 proof assistant (Lean 4.34 with Mathlib), in about 20,000 lines that depend only on Lean's three standard axioms (propositional extensionality, choice and quotient soundness). The formal statements are about the objects of the paper, not about a model of them: the sample is n independent points with the uniform distribution on the polygon (Lebesgue measure restricted to it), and the event is that some copy $\lambda R_\theta K + c$ of least λ containing the sample lies in K .

- Theorem 1, for every polygon of area one with vertices listed counterclockwise in strictly convex position: $\mathbb{P}(E_n)$ tends to the value of Theorem 10, which is the probability of E in the limit model. The proof follows Section 3; the Poisson processes, the multivariate Mecke equation and the Poisson approximation of Step 3 are developed from scratch, since Mathlib has no point processes.
- Lemmas 7, 8 and 9, the coercivity part of Lemma 5 (bounded tilt, and deep points never binding; existence of an optimum is not needed), and Theorem 10 for every polygon whose sides satisfy the nondegeneracy conditions used here (unit normals, positive width across parallel sides), which every strictly convex polygon of area one does.

- Corollary 2 for every triangle of area one, including the values $13/48$, $7/24$, $29/96$ and the limits $25/72$ and $1/3$; on the way the formalisation found the closed form $g(\alpha, \beta) = (\Phi(\alpha + \beta) - \Phi(\alpha - \beta)) / (2\alpha\beta)$ with $\Phi(x) = x^2/2 - x^4/12$ for $|x| \leq 1$ and $\Phi(x) = 2|x|/3 - 1/4$ for $|x| \geq 1$.
- Corollary 3 for every $q \geq 3$, with Vol the Lebesgue measure of the convex hull and int the topological interior; and $p_4 = 1/4$ in the limit model.

Not formalised: the reduction to area one by scaling, the numerical values of Section 8 (including p_q for $q \geq 5$), and Conjecture 4. Each formal statement was also tested against deliberately wrong variants (a changed constant, a dropped hypothesis, a reversed inequality); all of them fail to check. The Lean source, the simulation programs and an independent recheck of the triangle values are in the folder `smallest-enclosing-copy` of <https://github.com/jvvk/mathematics>.

8. NUMERICAL VERIFICATION

Every value in this paper was checked by at least two of three independent programs: (i) exact simulation of the finite problem, minimising over rotations with a linear program at each angle; (ii) Monte Carlo simulation of the limit LP with Poisson strips; (iii) evaluation of the formulas above. For regular q -gons with $q \geq 5$ the check is weaker: the values in Table 1 come from two separate implementations of Corollary 3 (for instance $p_5 = 0.1732$ and 0.17345 ± 0.00042), and limit-LP simulations confirm them only to about ± 0.01 .

shape	formula	limit LP (MC)	finite n
equilateral triangle	$13/48 = 0.2708$	0.2722 ± 0.0017	0.29 ± 0.03 ($n = 500$)
square	$1/4$	0.2515 ± 0.0016	0.30 ± 0.04 ($n = 500$)
30-60-90 triangle	$29/96 = 0.3021$	0.299 ± 0.009	0.30 ± 0.04 ($n = 400$)
20-40-120 triangle	0.3249	0.335 ± 0.009	
2×1 rectangle	0.2039	0.205 ± 0.008	0.16 ± 0.03 ($n = 400$)
regular hexagon	0.1935	0.189 ± 0.006	0.18 ± 0.04 ($n = 500$)
irregular centrally symmetric hexagon	0.1832	0.183 ± 0.008	

TABLE 2. Three independent checks. Errors are one standard error.

Direct simulation of the original problem. Program (i) was also run at scale for the equilateral triangle and the square: for each sample of n uniform points the smallest enclosing copy is found by exact geometry, with the best rotation located numerically, and we record whether some smallest copy lies inside K (Table 3). The pooled estimates are within 1.3 standard errors of $13/48$ and $1/4$; the single row furthest off, the triangle at $n = 5,000$, is about 2.4 standard errors low, which is unremarkable among seven rows.

The triangle formula was evaluated on a grid of 28,800 shapes (angle steps below 1°); its minimum is $13/48$ at the equilateral triangle, it increases away from it, and it tends to $25/72$ for flat isosceles triangles and $1/3$ for thin right and needle-shaped isosceles triangles. For two points the finite simulation reproduces the asker's values $2/3$ (square) and $3/4$ (triangle) [4].

9. OPEN PROBLEMS

Prove Conjecture 4. Find closed forms for p_q , $q \geq 5$; show that p_q converges as $q \rightarrow \infty$ and identify the limit (numerically about 0.146; $p_{200} = 0.1464$). Extend to smooth convex bodies other than the disk, and to higher dimensions, where the rotation group is larger.

shape	n	trials	fits	estimate
equilateral triangle	200	100,000	27,016	0.2702 ± 0.0014
	1,000	100,000	27,273	0.2727 ± 0.0014
	5,000	145,498	38,992	0.2680 ± 0.0012
	20,000	29,150	7,847	0.2692 ± 0.0026
	pooled	374,648	101,128	0.2699 ± 0.0007
square	200	100,000	24,981	0.2498 ± 0.0014
	1,000	100,000	25,053	0.2505 ± 0.0014
	5,000	79,200	19,907	0.2514 ± 0.0015
	pooled	279,200	69,941	0.2505 ± 0.0008

TABLE 3. Direct simulation of the finite problem (one standard error). The limits are $13/48 = 0.2708$ and $1/4$.

ACKNOWLEDGEMENT

The derivations, text, simulation programs and Lean formalisation were developed with the AI assistants Claude (Anthropic) and Codex (OpenAI), under the author's direction. The author is responsible for the contents.

REFERENCES

- [1] P. Groeneboom, Limit theorems for convex hulls, *Probab. Theory Related Fields* **79** (1988), no. 3, 327–368. <https://doi.org/10.1007/BF00342231>
- [2] P. Groeneboom, Convex hulls of uniform samples from a convex polygon, *Adv. Appl. Probab.* **44** (2012), no. 2, 330–342. <https://doi.org/10.1239/aap/1339878714>
- [3] G. Last and M. Penrose, *Lectures on the Poisson Process*, Institute of Mathematical Statistics Textbooks 7, Cambridge University Press, 2017. <https://doi.org/10.1017/9781316104477>
- [4] Dan, Question about n random points in a regular polygon, and a limiting probability, MathOverflow question 458571, 16 November 2023. <https://mathoverflow.net/q/458571> (accessed 8 October 2026).
- [5] L. Morin, Probability that n points are in convex position in a regular κ -gon: asymptotic results, *Adv. Appl. Probab.* **57** (2025), no. 3, 811–870. <https://doi.org/10.1017/apr.2024.63>
- [6] L. Morin, Probability that n points are in convex position in a general convex polygon: asymptotic results, *J. Appl. Probab.* **63** (2026), no. 3, 969–994. <https://doi.org/10.1017/jpr.2025.10066>
- [7] Dan, Intuition is silent: find the probability that the smallest circle enclosing n random points on a disk lies completely on the disk, as $n \rightarrow \infty$, Mathematics Stack Exchange question 4799757, 3 November 2023, with the accepted answer by abacaba, 8 November 2023. <https://math.stackexchange.com/q/4799757> (accessed 8 October 2026).