Racial and Ethnic Disparities in Emergency Department Analgesic Prescription

Joshua H. Tamayo-Sarver, PhD, Susan W. Hinze, PhD, Rita K. Cydulka, MD, MS, and David W. Baker, MD, MPH

Racial/ethnic disparities in the prescription of analgesics appear widespread¹ and are evident in fracture treatment, ^{2,3} cancer pain, ⁴ and postoperative pain. ⁵ In a particularly striking series of studies, Todd et al. ^{2,3,6} demonstrated that African Americans and Latinos were significantly less likely to receive analgesia in the emergency department (ED) for isolated long bone fractures than were Whites, despite the fact that physicians rated patients' pain as similar in severity. These studies raise concerns that patients may be receiving inadequate pain control, and this suffering may fall disproportionately on minorities. ⁷

Previous studies provide support for several hypotheses to explain why racial/ethnic differences in prescription of analgesics might occur.8-10 Minority patients tend to be less assertive and less active in the physicianpatient interaction⁸ and less satisfied with their ability to communicate with their respective physicians¹¹ than Whites. Physicians' perceptions of patients also vary by race/ ethnicity. 10 Communication has been shown to be less effective when social distance exists between the physician and the patient, 9,12-15 so some of the racial/ethnic differences may reflect more frequent communication difficulties that result from the generally lower socioeconomic status (SES) of minority patients or the general underrepresentation of minorities among physicians. These communication difficulties may lead to a physician being less likely to appreciate a patient's pain and less likely to prescribe analgesics.

If some of the racial/ethnic differences in analgesic use are caused by racial/ethnic variations in physician—patient communication, then disparities in the prescription of analgesics should be greatest when the clinician must rely on the medical history to determine the cause and severity of a patient's pain. Thus, differences should be least for conditions with unequivocal objective findings, such

Objectives. We examined racial and ethnic disparities in analgesic prescription among a national sample of emergency department patients.

Methods. We analyzed Black, Latino, and White patients in the 1997–1999 National Hospital Ambulatory Medical Care Surveys to compare prescription of any analgesics and opioid analgesics by race/ethnicity.

Results. For any analgesic, no association was found between race and prescription; opioids, however, were less likely to be prescribed to Blacks than to Whites with migraines and back pain, though race was not significant for patients with long bone fracture. Differences in opioid use between Latinos and Whites with the same conditions were less and nonsignificant.

Conclusions. Physicians were less likely to prescribe opioids to Blacks; this disparity appears greatest for conditions with fewer objective findings (e.g., migraine). (Am J Public Health. 2003;93:2067–2073)

as long bone fracture; intermediate for conditions with few objective findings, such as back pain or strain; and greatest for conditions with almost no objective findings, such as a migraine headache. Similarly, if physicians have greater social distance from minorities, then these physicians may view their patients' reports of pain with less credibility. If so, disparities in the act of prescribing should be greatest for opioid analgesics, which require some level of trust that the patient's complaint is valid and that the medication will not be misused. Opioids generally offer several advantages to the clinician: the ability to deliver medication intravenously, potency, quick action, easily reversed action, and easy titration relative to nonopioid analgesics.16 However, opioids may also raise physician concerns that the patient may be seeking opioids in order to satisfy an addiction or to sell them. Physicians may have more negative perceptions of minority patients 10,17 and feel they are at higher risk for abuse or sale of the opioid. 18 If physicians tend to trust minority patients less than White patients, then we hypothesized the disparities would be greater for prescription of opioid analgesics than nonopioid analgesics.

Previous studies of analgesic prescription in the ED were small^{2,3,19} and limited to single institutions. These studies may not be representative of care patterns across the United

States. The purpose of this study was to examine racial/ethnic disparities in analgesic practice at a national level. To test the hypothesis that physician–patient communication contributes to the racial/ethnic disparity, we compared the racial/ethnic disparities among 3 conditions with increasing levels of objective findings: migraine, back pain or strain, and long bone fracture.

METHODS

Study Population

We combined data from the 1997, 1998, and 1999 National Hospital Ambulatory Medical Care Survey (NHAMCS). The NHAMCS is a nationally representative sample of visits to nonfederal, short-stay hospital EDs that was conducted by the National Center for Health Statistics (NCHS).20-22 The NHAMCS used a 4-stage probability sampling procedure that selected counties (or equivalents), then hospitals, then emergency service areas. Finally, hospital staff trained by NCHS personnel prospectively selected a random sample of patient visits during a randomly assigned 4-week reporting period. A patient record form was completed by hospital staff and reviewed and validated by NCHS staff. The NHAMCS employed routine quality control measures. A NCHS field representative reviewed the log or other records

used for visit sampling to determine whether any cases were missing and also edited completed forms for missing data. Attempts were made to retrieve both missing cases and missing data on specific cases, either by consulting with the appropriate hospital staff or by reviewing the pertinent medical records. All medical and drug coding and keying operations were subject to quality control procedures. Quality control for the medical and drug coding operation, as well as straight-key items, involved a 2-way, 10% independent verification procedure. As an additional quality control, all patient record forms with coding variations or with illegible entries for the reason for a visit, diagnostic and therapeutic procedures, diagnosis, E-code (cause of injury), and medication items were reviewed and adjudicated at NCHS. The NHAMCS data can be used to produce national estimates through the weighting procedure that accounts for the sample design, nonresponse, and fixed totals.

We first examined analgesic use for all patients in the NHAMCS database. To further explore disparities in analgesic prescription, we examined 3 common conditions with increasing levels of objective physical findings: migraine (International Classification of Diseases, 9th Revision [ICD-9] 346), back pain or strain (ICD-9 724), and long bone fracture (ICD-9 812, 813, 821, and 823).

Dependent Variables

Each medication administered in the ED, prescribed at discharge, or discussed (i.e., recommended that a patient continue to take a previously prescribed medication) was abstracted from the patient record form. We identified analgesic medications using all the National Drug Code Directory codes for "analgesia" determined by NHAMCS, including medications specific for headache/migraine relief. Identification of opioid analgesic was similarly identified using the National Drug Code Directory code for "narcotic analgesia" determined by NHAMCS. From this data, we created 2 dependent variables to indicate whether the patient was prescribed, received, or instructed to continue to use (1) any analgesic and (2) any opioid analgesic.

Independent Variables

Classification of individuals into racial and ethnic groups has been a contentious issue. 23-26 For the purpose of this study, we were interested in how analgesic prescription varies according to patients' appearance, which includes physical characteristics, dress, language, and mannerisms. A classic definition of race is "any group of people who are distinguished, or consider themselves distinguished, in social relations with other peoples, by their physical characteristics."27 A more contemporary definition of race, however, emphasizes that the importance of physical variations in the human species are socially constructed.²⁸ Ethnicity can be thought of as a self-perceived group of people who hold a common set of traditions, including folk and religious beliefs and practices, language, a sense of historical continuity, and common ancestry or place of origin.²⁹ Thus, health care providers' impressions of a patient's appearance are likely to be based on an individual's race and ethnicity, and we therefore use the term race/ethnicity throughout.

The race/ethnicity recorded in NHAMCS likely reflects the hospital staff's perception of a patient's race and ethnicity rather than the classification that a patient might choose. Because the clinician determines the prescription of analgesics, it is the clinician's perception of a patient's race/ethnicity that is most relevant for this analysis. The NHAMCS classified the patient's race as White, Black, American Indian or Alaska Native, or Asian or Pacific Islander by the hospital staff, with explicit instructions from NHAMCS not to ask the patient unless it was hospital procedure to do so. The patient's ethnicity was categorized as Hispanic or non-Hispanic. Based on this, we created 5 racial/ ethnic groups that we refer to as American Indian or Alaska Native, Asian or Pacific Islander, Black, Latino, and White.25 We considered any patient recorded as Hispanic to be Latino, regardless of other racial classifications. For this analysis, we present data only for Blacks, Whites, and Latinos.

Covariates

Many factors other than race/ethnicity may influence physicians' decisions to prescribe analgesics in general and opioid analgesics in particular. We adjusted for patients' age, sex, insurance coverage (recorded as method of payment), major complaints presented (using the 3 diagnoses recorded), severity (triage assignment and pain assessment), visit characteristics (mode of arrival and discharge status), and hospital characteristics (hospital ownership, region, urban vs rural, and year of visit). Method of expected payment was categorized as private, Medicare, Medicaid, worker's compensation, self-pay/uninsured, other, unknown, or missing. The triage nurse's assessment of the time within which the patient should be examined was categorized as <15 minutes, 15 to 60 minutes, 1 to 2 hours, or \geq 2 hours by means of a predefined scheme to convert the hospital's triage system to the above scale. Pain severity was categorized as none, mild, moderate, severe, unknown, or missing, based on chart abstraction with a predefined scheme to convert terms in the medical record into the above scale. The mode of arrival was categorized as ambulance, police or social services (public service), walk-in, unknown, or missing. Discharge status was categorized as no follow-up planned, follow-up with referring physician, return to the ED if needed, triaged out of ED before being seen, follow-up with a physician, left without being seen, admitted to the hospital, admitted to the intensive care unit or cardiac care unit, transferred to another hospital, dead on arrival, referred to social services, or other. Hospital ownership was categorized as for profit, voluntary nonprofit, or government nonprofit. Region was categorized as Northeast, Midwest, South, or West.

Statistical Analyses

Chi-square tests were used to test bivariate associations, and multiple logistic regression was used to determine the independent association between race/ethnicity and analgesic prescription after adjusting for covariates. To fully adjust for possible racial/ethnic differences in patients' presenting complaints, we conducted additional analyses using fixedeffects logistic regression, 30 with the full logistic model comparing only patients who had the same 3 diagnoses. To determine whether disparities in prescription practices differed by the objectivity of clinical findings, we conducted 3 separate stratified regressions for patients with long bone fracture, back pain and strain, and migraine, by means of the same variables from the full logistic model used with the entire sample. To ensure that the results were not affected by racial/ethnic

differences in whether the target conditions were primary versus secondary complaints, we repeated the regression model with 2 different samples: only patients with a primary diagnosis of the condition (e.g., migraine) and no secondary or tertiary diagnoses, and any patient with the diagnosis. The coefficients did not differ between the 2 models,³¹ so we used the more inclusive sample to provide greater precision of the estimated association. We then approximated the relative risk from the odds ratios as described by Zhang and Yu.³² Adjusted proportions were calculated based on the relevant model using the ADJUST command in Stata Version 7.0 (Stata Corp, College Station, Tex), which sets all covariates in the model to their sample mean values. All bivariate and multivariate analyses that we used adjusted sample

weights to account for the sampling design by a method suggested by the NCHS and used previously.^{21,33} Based on findings from previous studies,³⁴ we examined possible interactions between race and gender, but the interaction was not significant in the whole population and the sample size was not sufficient to produce stable estimates within each of the 3 specific conditions. All analyses were performed with *Stata Version 7.0*.

RESULTS

Sample Characteristics

Of the 67 487 patients in our sample, 21% (15 108) were Black, 9% (7 523) were Latino, and 68% (42 926) were White (Table 1); 2% (1930) of patients were classified as being from other races (other than White, Black, or

TABLE 1—Patient Characteristics of Participants in the National Hospital Ambulatory Medical Care Survey: 1997-1999

Characteristic	Total	White	Black	Latino	P value
Sample size ^a	67 487	42 926	15 108	7 523	
Population estimate ^b	99 361 812	67 487 644	20 373 375	9 219 558	
Female (%)	53	52	55	51	<.001
Age (mean + SD)	35 ± 24	38 ± 25	32 ± 22	27 ± 21	<.001
Urgency (%)					<.001
<15 min	19	21	15	15	
15-60 min	31	32	29	32	
1-2 h	15	14	18	19	
2-24 h	9	9	11	11	
Unknown	25	25	27	23	
Pain (%)					<.001
None	12	12	12	14	
Mild	20	20	20	18	
Moderate	14	15	12	14	
Severe	4	5	3	4	
Unknown/blank	50	48	53	50	
Arrival (%)					.12
Ambulance	6	6	6	5	
Public service	10	11	10	8	
Walk-in	28	28	28	32	
Unknown/missing	56	55	56	55	
Urban Hospital (%)	76	71	86	93	<.001
Admitted (% yes)					<.001
General	13	14	10	10	
ICU/CCU	1.4	1.7	0.9	0.7	

Notes. SD = standard deviation; ICU/CCU = intensive care unit/cardiac care unit.

Latino). This sample is representative of an average of 99 million ED visits annually in the United States between 1997 and 1999. Important differences across racial/ethnic groups were noted for several patient characteristics. Mean age was highest for Whites and lowest for Latinos. Whites had more visits classified as urgent and were admitted to the hospital more often than Blacks or Latinos (Table 1). Pain severity ratings were similar for the 3 groups, although the differences were significant because of the very large sample size.

Bivariate Results

In bivariate analysis, Whites, Blacks, and Latinos in the entire sample appeared to be equally likely to receive some form of analgesic, but Whites were more likely to have received an opioid analgesic. Among the entire population, 62% of Whites did not receive any analgesic; the rates were similar for Blacks and Latinos (Table 2). Among those who received some type of analgesic, 34% of Whites, 23% of Blacks, and 23% of Latinos (P < 0.001) received an opioid.

For patients with migraines, 16% of Whites, 28% of Blacks, and 20% of Latinos received no analgesic (Table 2). Whites were not only more likely than Blacks to receive an analgesic for migraines (P<0.001) but also were more likely to receive an opioid analgesic (P < 0.001). Latinos were less likely to receive an analgesic or opioid analgesic than Whites, but these differences were not statistically significant. Among patients with back pain, 20% of Whites received no analgesic, compared with 30% of Blacks (P < 0.01) and 23% of Latinos (P=0.51) (Table 2). More than half of Whites (54%) who received an analgesic received an opioid analgesic. In contrast, only a quarter (27%) of Blacks who received any analgesic received an opioid. Latinos received opioids at similar rates to Whites. Finally, one third of patients with long bone fractures did not receive any analgesic while approximately one third received an opioid; these proportions were similar for all 3 races/ ethnicities (Table 2).

Multivariate Results

Multiple logistic regression demonstrated that race/ethnicity was not associated with whether or not a patient received any analgesic (data not shown). After adjusting for SES,

^aA total of 1930 patients were classified as being from other races (other than White, Black, or Latino).

^bThis figure represents the average annual number of visits for the US population between 1997 and 1999.

TABLE 2—Receipt of Analgesic in the Emergency Department for Patients in the National Hospital Ambulatory Medical Care Study: 1997–1999

Entire Population	White	Black	P vs White	Latino	P vs White
Sample size	42 926	15 108		7 523	
Population estimate	67 487 644	20 373 375		9 219 558	
Received analgesic (%)			< 0.01		<.01
None	62	63		62	
Nonopioid	25	28		29	
Opioid	13	8		9	
Opioid/all analgesics (%)	34	23	< 0.01	23	<.01
		Patients with migrai	ne		
Sample size	471	92		43	
Avg. annual visit (est.)	784 582	140 914		52 949	
Received analgesic (%)			< 0.01		.51
None	16	28		20	
Nonopioid	24	39		30	
Opioid	61	33		50	
Opioid/all analgesics (%)	72	45	< 0.01	63	.29
	Pa	tients with back prot	olems		
Sample size	912	273		126	
Avg. annual visit (est.)	1 441 918	396 398		156 054	
Received analgesic (%)			< 0.01		.38
None	20	30		23	
Nonopioid	37	59		42	
Opioid	43	21		35	
Opioid/all analgesics (%)	54	27	< 0.01	45	.18
	Patie	nts with long bone fr	actures		
Sample size	602	80		71	
Avg. annual visit (est.)	969 912	90 606		94 339	
Received Analgesic (%)			0.34		.57
None	33	28		37	
Nonopioid	31	42		25	
Opioid	36	30		38	
Opioid/all analgesics (%)	54	42	0.20	61	.46

demographics, severity, visit characteristics, and hospital characteristics, the adjusted relative risk for Blacks and Latinos to receive any analgesic ranged from 0.89 to 1.03, none of which was statistically significant. The adjusted relative risk for Blacks in the whole population was 1.02 (95% confidence interval [CI], 0.98–1.05) and was 1.03 (95% CI, 0.98–1.08) for Latinos, which demonstrates that race/ethnicity was not associated with whether or not a patient was prescribed any analgesic.

Although race/ethnicity was not independently associated with receiving any analgesic, race/ethnicity was associated with receiving an opioid analgesic in multivariate


analyses. Among the entire population, the conditional logistic regression, which conditioned the regression on the first 3 diagnoses listed for the visit, demonstrated that Blacks (adjusted relative risk 0.72; 95% CI, 0.66-0.79) and Latinos (adjusted relative risk 0.72; 95% CI, 0.64-0.81) were both 28% less likely than Whites to receive opioid analgesic (Table 3). The differences in opioid prescription were similar regardless of patients' insurance coverage (i.e., private, Medicaid, or self-pay). The difference in opioid use for Blacks and Whites was also seen for the subgroups of patients with migraine and back pain; however, the difference was somewhat less for patients with long bone fracture and

was not statistically significant (Table 3). The differences in opioid use between Latinos and Whites were much smaller than for Blacks, and none were statistically significant (Table 3). Nevertheless, as we hypothesized, the greatest difference was seen for the condition with the least objective findings (migraine) and the least difference was seen for the condition with the most objective findings (long bone fracture; Figure 1). The results were similar when we analyzed the likelihood of receiving an opioid among only those patients who received any form of analgesic.

DISCUSSION

Blacks, Latinos, and Whites were equally likely to receive some form of analgesic for the entire population of patients using the ED as well as the subgroups with migraine, back pain, or long bone fracture. Previously, Todd et al. found that Latinos and Blacks were less likely to receive any analgesic for long bone fracture,^{2,3} while Karpman¹⁹ found no disparity between Latinos and Whites. It is possible that the single-institution studies were not nationally representative, or that the seminal report by Todd et al. in 1993 focused the attention of ED physicians on this issue, so that any national discrepancies that existed at the time of the publication of their study had been minimized by the period 1997-1999, when the data were collected for this study.

The differences between our results and those of earlier reports could also result from differences in study methodology. We identified long bone fractures using the same ICD-9 codes. However, Todd recorded only analgesics administered in the ED. Additionally, we were unable to exclude patients being treated for complications of previously treated fractures and cases where the radiology report did not confirm a fracture. Despite these methodological differences, the rate of analgesic use among White patients with long bone fracture was similar in our study and the reports from Todd et al. In contrast, the rates of analgesic use for Blacks and Latinos were substantially higher than in previous studies.² Nationally, 67% of Whites, 72% of Blacks, and 63% of Latinos received some type of analgesic, in contrast to Todd et al.'s



Note. Whites = black bar; Blacks = dark gray bar; Latinos = light gray bar. All logistic regressions adjust for the covariates of socioeconomic status (SES)/demographics (assessed by sex, age, and method of payment), severity (assessed by triage assignment and pain assessment), visit characteristics (assessed by mode of arrival and discharge status), and hospital characteristics (assessed by hospital ownership, region, urban vs rural, and year of visit).

^aA conditional logistic or fixed effects logistic regression was done for the entire population; therefore, the estimates are based on comparisons being made only among patients with the same 3 diagnoses by ICD-9 codes.

FIGURE 1—Adjusted proportions of patients who received an opioid analgesic, for all patients in National Hospital Ambulatory Medical Care Survey combined and for those presenting with migraine, back pain, and long bone fracture.

TABLE 3—Adjusted Relative Risk^a (95% CI) of Being Prescribed an Opioid Analgesic for African Americans and Latinos Compared With Whites in the National Hospital Ambulatory Medical Care Study: 1997–1999

	African Americans	Latinos
All patients ^b	0.72 (0.66-0.79)	0.72 (0.64-0.81)
Migraine	0.63 (0.40-0.89)	0.74 (0.41-1.11)
Back pain	0.50 (0.36-0.69)	0.85 (0.56-1.18)
Long bone fracture	0.74 (0.43-1.18)	0.97 (0.56-1.49)

Note. CI = confidence interval.

^aAnalyses are presented for all patients combined and for patients presenting with migraine, back pain, and long bone fracture. All logistic regressions adjust for the covariates of socioeconomic status or demographics (assessed by sex, age, and method of payment), severity (assessed by triage assignment and pain assessment), visit characteristics (assessed by mode of arrival and discharge status), and hospital characteristics (assessed by hospital ownership, region, urban vs rural, and year of visit).

finding that 74% of Whites, ^{2,3} 57% of Blacks, ² and 45% of Latinos ³ received any type of analgesic.

Although we found no difference in overall analgesic prescription, Blacks and Latinos in the entire sample were less likely than Whites to receive an opioid analgesic. This finding is consistent with our hypothesis that disparities would be greater for opioid prescriptions than nonopioids, because prescribing an opioid re-

quires more trust of the patient by the physician. Among the subgroups, Blacks were far less likely to receive an opioid analgesic than Whites for both migraine and back pain, but there was no difference for all patients with a long bone fracture. This finding is consistent with our a priori hypothesis that racial/ethnic differences in analgesic prescription would be least for conditions with clear, objective findings (long bone fracture) and greatest for con-

ditions with less objective findings (migraine, back pain) that require more provider—patient communication to arrive at a diagnosis and a treatment plan. There were no differences in opioid use between Whites and Latinos for these 3 conditions, although the power to detect differences was limited by the small number of Latinos with these diagnoses and the need to inflate standard errors to account for the clustering of patients within hospitals.³³

Although we found no racial/ethnic differences in overall analgesic use, our finding that between a sixth and a third of patients diagnosed with 1 of 3 painful conditions did not have the prescription, administration, or recommendation of an analgesic recorded during an ED visit should raise concern. The low rate of analgesic use is consistent with a recent study that found only 44% of ED patients rated their pain control as "very good." The NHAMCS attempted to identify all "medications that were ordered, supplied, administered, or continued during this visit," including "drugs and medications that the physician ordered or provided before this visit and instructs or expects the patient to continue taking regardless of whether a 'refill' is provided at the time of visit." This apparently high proportion of patients not prescribed an analgesic could partly be due to lack of documentation when physicians tell patients to take over-the-counter analgesics or to continue with medications they have at home.² Restricting the measure to only medications administered in the ED may have reduced this misclassification bias, but the separate data were not collected by NHAMCS.

Alternative Explanations for Findings

We hypothesized that racial/ethnic differences in patient assertiveness, ⁸ physician perception of the patient, ¹⁰ and social distance ¹² may contribute to differences in physician—patient communication ¹¹ and trust that are responsible in part for the racial/ethnic disparities in analgesic prescription. The results of this study provide support for the role of patient communication and trust in the creation of these disparities, but the dimensions were not directly measured in the current study and the data support several alternative explanations. The 3 conditions studied vary substantially in their natural history, ranging from

^bA conditional logistic or fixed effects logistic regression was done for the entire population; therefore, the estimates are based on comparisons being made only among patients with the same 3 diagnoses by ICD-9 codes.

RESEARCH AND PRACTICE

acute with defined duration (fracture), to acute but slowly resolving with an uncertain prognosis (back pain), to recurrent over many years (migraine). The variation in the racial/ ethnic differences in opioid prescription found across these 3 conditions could result from this variation in natural history and course of disease rather than the presence of objective findings on physical examination. Perhaps physicians were reluctant to prescribe opioids for a lengthy condition if a patient lacked a regular source of care. Additionally, some physicians may have prescribed nonopioid analgesics to Blacks and Latinos instead of an opioid because of the difficulties faced by some minorities in acquiring opioid analgesics from local pharmacies.³⁶ The NHAMCS did not distinguish between medications administered in the ED and those prescribed at discharge. Therefore, we could not determine whether the racial/ethnic differences in opioid prescription resulted from variations in opioid administration or discharge medications. Possibly Whites were more likely to have already attempted pain control with nonopioid, over-the-counter pain relievers. When a patient presented to the ED, the physician may have been more likely to prescribe an opioid because the patient reported using a nonopioid without success. Additionally, medications that the physician offered a patient but were declined were not identified. Possibly the disparities in treatment were a result of differing patient preferences. However, previous research has demonstrated no racial/ethnic differences in ratings of pain severity^{7,37–39} or preference for analgesia.⁷

Limitations

Although our study found an association between race/ethnicity and the rate of opioid prescription, it is not possible to say whether these differences are related to race/ethnicity per se or to some unmeasured variable that is highly associated with race/ethnicity, such as SES. 10 Although the NHAMCS does not contain information on patients' SES, neither Medicaid nor self-pay status was significant in any of our models. Additionally, Blacks and Latinos are equally likely to be poor, 40 and Latinos tend to have lower educational attainment than Blacks.41 Although this study did not have adequate

power, the trend for opioid prescription to be lower among Blacks than Latinos suggests that race may be a more important correlate than SES.

Additionally, despite rigorous efforts by NHAMCS to achieve consistent data collection across sites, there may have been significant variation in the coding of some covariates (i.e., assessment of pain and urgency) or systematic bias in these measurements by race/ethnicity (e.g., triage nurses giving Whites higher urgency ratings for similar problems). However, if race/ethnicity were related to any of the severity variables, such that a White was assigned a higher severity despite being clinically identical to a Black or Latino, then our multivariate analysis would underestimate the true association between race/ethnicity and analgesic use. This would lead to an underestimate of the differences in prescription of both opioids and any analgesics and contribute to our finding of no difference in prescription of any analgesic. The large number of missing pain scores could have introduced bias if one group was more likely to have their pain assessed; however, missing pain assessment did not differ by race/ethnicity within the 3 conditions and removal of the pain variable from the multiple regression did not significantly change the coefficients for race.³¹

Future Directions

The Joint Commission on Accreditation of Hospital Organizations has recommended routine use of a pain-rating scale at the time that vital signs are taken. 42 Although this recommendation may be helpful for addressing the widespread problem of inadequate pain control, it is unclear whether it will achieve greater equity of analgesic prescription across all racial/ethnic groups. Earlier studies found differences in analgesic use for Whites and Latinos, even though physicians' average ratings of pain severity for these groups were nearly identical.^{3,6} It may be helpful for EDs to routinely monitor the relationship between analgesic prescription and patients' pain severity ratings for Whites, Blacks, and Latinos to detect inadequate pain control and to attempt to address it. In addition, more studies are needed to understand the fundamental causes of disparities in analgesic prescription. Our findings and previous research suggest that a better understanding of provider-patient communication and trust may be a fruitful focus for future research and interventions.

About the Authors

Joshua H. Tamayo-Sarver is with the Department of Epidemiology and Biostatistics, Case Western Reserve University School of Medicine, Cleveland, Ohio. Susan W. Hinze is with the Department of Sociology, Case Western Reserve University. Rita K. Cydulka is with the Department of Emergency Medicine, MetroHealth Medical Center, Case Western Reserve University School of Medicine. David W. Baker is with the Division of General Internal Medicine, Northwestern University Medical School, Chicago, Illinois.

Correspondence should be sent to Joshua H. Tamayo-Sarver, 4009 Cullen Drive, Cleveland, OH 44105 (e-mail: sarver@po.cwru.edu).

Contributors

J.H. Tamayo-Sarver conceived the study, performed the analysis, interpreted the results, and led the writing. D.W. Baker assisted in the analysis, interpretation, and manuscript revision. S. W. Hinze and R. K. Cydulka assisted in the interpretation of the data and revision of the article. All authors helped to conceptualize ideas, interpret findings, and review drafts of the article.

Acknowledgments

J.H. Tamayo-Sarver was supported by the Agency for Healthcare Research and Training (grant HS-00059-06) and Case Western Reserve University School of Medicine.

Human Participant Protection

Local institutional review board approval was not sought because this study was a secondary analysis of a publicly available dataset that contained no individual or institutional identifying information.

References

- 1. Bonham VL. Race, ethnicity, and pain treatment: striving to understand the causes and solutions to the disparities in pain treatment. J Law Med Ethics. 2001; 29:52-68.
- Todd KH, Deaton C, D'Adamo AP, Goe L. Ethnicity and analgesic practice. Ann Emerg Med. 2000;35:
- Todd KH, Samaroo N, Hoffman JR. Ethnicity as a risk factor for inadequate emergency department analgesia. JAMA. 1993;269:1537-1539.
- 4. Bernabei R, Gambassi G, Lapane K, et al. Management of pain in elderly patients with cancer. SAGE Study Group. Systematic Assessment of Geriatric Drug Use via Epidemiology. JAMA. 1998;279:1877-1882.
- Ng B, Dimsdale JE, Rollnik JD, Shapiro H. The effect of ethnicity on prescriptions for patient-controlled analgesia for post-operative pain. Pain. 1996;66:9-12.
- Todd KH, Lee T, Hoffman JR. The effect of ethnicity on physician estimates of pain severity in patients with isolated extremity trauma. JAMA. 1994; 271:925-928.

RESEARCH AND PRACTICE

- 7. Todd KH. Pain assessment and ethnicity. Ann Emerg Med. 1996;27:421-423.
- Sleath B, Roter D, Chewning B, Svarstad B. Asking questions about medication: analysis of physicianpatient interactions and physician perceptions. Med Care. 1999;37:1169-1173.
- 9. Cooper-Patrick L, Gallo JJ, Gonzales JJ, et al. Race, gender, and partnership in the patient-physician relationship. JAMA. 1999;282:583-589.
- 10. van Ryn M, Burke J. The effect of patient race and socio-economic status on physicians' perceptions of patients. Soc Sci Med. 2000;50:813-828.
- 11. LaVeist TA, Nickerson KJ, Bowie JV. Attitudes about racism, medical mistrust, and satisfaction with care among African American and white cardiac patients. Med Care Res Rev. 2000;57(Suppl 1):146-161.
- 12. Atkinson P. Medical Talk and Medical Work: The Liturgy of the Clinic. London, England: Sage, 1995.
- 13. Ross CE, Duff RS. Returning to the doctor: the effect of client characteristics, type of practice, and experiences with care. J Health Soc Behav. 1982;23: 119-131.
- 14. Robinson BJ. Africanisms and the study of folklore. In: Holloway JE, ed. Africanisms in American Culture. Bloomington: Indiana University Press, 1991.
- 15. Madsen W. Hidalgo Project on Differential Culture Change and Mental Health. Mexican-Americans of South Texas. 2d ed. New York, NY: Holt, Rinehart and Winston, 1973.
- 16. Golianu B, Krane EJ, Galloway KS, Yaster M. Pediatric acute pain management. Pediatr Clin North Am. 2000;47:559-587.
- Rathore SS, Lenert LA, Weinfurt KP, et al. The effects of patient sex and race on medical students' ratings of quality of life. Am J Med. 2000;108:561-566.
- 18. Martin ML. Ethnicity and analgesic practice: an editorial. Ann Emerg Med. 2000 Jan;35:77-79.
- Karpman RR, Del Mar N, Bay C. Analgesia for emergency centers' orthopaedic patients: does an ethnic bias exist? Clin Orthop. 1997 Jan;(334):270-275.
- 20. Nourjah P. National Hospital Ambulatory Medical Care Survey: 1997 emergency department summary. Adv Data. 1999 May 6;(304):1-24.
- 21. Stone S, Gonzales R, Maselli J, Lowenstein SR. Antibiotic prescribing for patients with colds, upper respiratory tract infections, and bronchitis: a national study of hospital-based emergency departments. Ann Emerg Med. 2000;36:320-327.
- 22. McCaig LF. National Hospital Ambulatory Medical Care Survey: 1998 emergency department summary. Adv Data. 2000 May 10;(313):1-23.
- 23. LaVeist TA. On the study of race, racism, and health: a shift from description to explanation. Int J Health Serv. 2000;30:217-219.
- 24. LaVeist TA. Beyond dummy variables and sample selection: what health services researchers ought to know about race as a variable. Health Serv Res. 1994 Apr;29:1-16.
- 25. Williams DR. Race and health: basic questions, emerging directions. Ann Epidemiol. 1997 Jul;7: 322-333.
- 26. Winant H. Race and race theory. Annu Rev Sociol 2000;26:169-185.

- 27. Cox OC. Caste, Class, & Race: A Study In Social Dynamics. 1st ed. Garden City, NY: Doubleday, 1948.
- 28. American Anthropological Association. Statement on "Race." Available at: http://www.aaanet.org/stmts/ racepp.htm. Accessed September 15, 2000.
- 29. De Vos GA. Ethnic pluralism: conflict and accommodation. In: De Vos GA, Romanucci-Ross L, eds. Ethnic Identity: Cultural Continuities and Change. Chicago: University of Chicago Press, 1982:5-41.
- 30. Diggle PJ, Liang K, Zeger SL. Analysis of Longitudinal Data. Midsomer Norton, Avon, Great Britain: Oxford University Press, 1994.
- 31. D'Agostino RB Sr, Grundy S, Sullivan LM, Wilson P. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. JAMA. 2001;286:180-187.
- 32. Zhang J, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. JAMA. 1998;280:1690-1691.
- 33. Potthoff RF, Woodbury MA, Manton KD. "Equivalent sample size" and "equivalent degrees of freedom": refinements for inference using survey weights under superpopulation models. J Am Stat Assn. 1992;87: 383-396
- 34. Schulman KA, Berlin JA, Harless W, et al. The effect of race and sex on physicians' recommendations for cardiac catheterization [published erratum appears in N Engl J Med 1999;340:1130]. N Engl J Med. 1999; 340:618-626.
- 35. Lanser P, Gesell S. The fifth vital sign. Satisfaction Monitor. March/April 2001;1-10.
- 36. Morrison RS, Wallenstein S, Natale DK, Senzel RS, Huang LL. "We don't carry that"-failure of pharmacies in predominantly nonwhite neighborhoods to stock opioid analgesics. N Engl J Med. 2000;342: 1023-1026
- 37. Zatzick DF, Dimsdale JE. Cultural variations in response to painful stimuli. Psychosom Med. 1990;52: 544-557.
- 38. Pfefferbaum B, Adams J, Aceves J. The influence of culture on pain in Anglo and Hispanic children with cancer. J Am Acad Child Adolesc Psychiatry. 1990;29: 642 - 647.
- 39. Greenwald HP. Interethnic differences in pain perception. Pain. 1991:44:157-163.
- 40. Dalaker J, Proctor BD. Poverty in the United States: 1999. US Census Bureau, Current Population Reports, Series P60-210, Washington, DC: U.S. Government Printing Office; 2000.
- 41. Newburger EC, Curry A. Educational Attainment in the United States: 1999. US Census Bureau, Current Population Reports, Series P20-528. Washington, DC: U.S. Government Printing Office; 2000.
- 42. Gorman C. Feel no pain. What you need to know about the new rules for treating pain at hospitals and other facilities. Time. January 2, 2001;157(1):76.



Early Lessons Learned in Children's Environmental Health

By Sarah Bartlett and John Petrarca

The Schools of Ground Zero is the story ■ of how public school districts in Lower Manhattan dealt with the events of Sept.11, 2001, and their aftermath. Written by the parents of two children who attend public school in lower Manhattan, this book focuses on the effect that the Sept. 11 attacks had on the physical and mental health of New York City schoolchildren and their parents.

The Schools of Ground Zero uses the example of the World Trade Center attacks to illustrate how school officials may be unprepared to cope with emergencies, and uncertain how to proceed after the event. Drawing on interviews with parents, teachers, New York Board of Education officials and environmental consultants, the authors make practical recommendations for safeguarding the health and safety of schoolchildren in times of crisis.

> ISBN 0-87553-028-1 402 pages ■ Softcover ■ 2002 Nonmember: 29.95 Member: \$21.50 Plus shipping and handling

American Public Health Association



Publication Sales Web: www.apha.org E-mail: APHA@TASCO1.com Tel: (301) 893-1894 FAX: (301) 843-0159 SG710111

This article has been cited by:

- 1. Sara E. Heins, Mark J. Sorbero, Christopher M. Jones, Andrew W. Dick, Bradley D. Stein. 2018. High-Risk Prescribing to Medicaid Enrollees Receiving Opioid Analgesics: Individual- and County-Level Factors. Substance Use & Misuse 53:10, 1591-1601. [Crossref]
- 2. Carlo J. Milani, Sean D. Rundell, Jeffrey G. Jarvik, Janna Friedly, Patrick J. Heagerty, Andy Avins, David Nerenz, Laura S. Gold, Judith A. Turner, Thiru Annaswamy, Srdjan S. Nedeljkovic, Pradeep Suri. 2018. Associations of Race and Ethnicity With Patient-Reported Outcomes and Health Care Utilization Among Older Adults Initiating a New Episode of Care for Back Pain. SPINE 43:14, 1007-1017. [Crossref]
- 3. Brad Wright, Xuan Zhang, Momotazur Rahman, Mahshid Abir, Padmaja Ayyagari, Keith E. Kocher. 2018. Evidence of Racial and Geographic Disparities in the Use of Medicare Observation Stays and Subsequent Patient Outcomes Relative to Short-Stay Hospitalizations. *Health Equity* 2:1, 45-54. [Crossref]
- 4. Amit Jain, Emmanuel Menga, Addisu Mesfin. 2018. Outcomes Following Surgical Management of Cauda Equina Syndrome: Does Race Matter?. *Journal of Racial and Ethnic Health Disparities* 5:2, 287-292. [Crossref]
- 5. Larry Charleston IV, James Francis Burke. 2018. Do racial/ethnic disparities exist in recommended migraine treatments in US ambulatory care?. *Cephalalgia* 38:5, 876-882. [Crossref]
- 6. Hsien-Chang Lin, Zhi Wang, Carol Boyd, Linda Simoni-Wastila, Anne Buu. 2018. Associations between statewide prescription drug monitoring program (PDMP) requirement and physician patterns of prescribing opioid analgesics for patients with non-cancer chronic pain. *Addictive Behaviors* 76, 348-354. [Crossref]
- 7. Samantha M. Pruden, Angela Starkweather. 2017. Prescription Opioid Abuse and Addiction. *Topics in Pain Management* 33:5, 1-7. [Crossref]
- 8. Daniel P. Marcusa, Rachel A. Mann, David C. Cron, Brooklyn R. Fillinger, Alexandra K. Rzepecki, Jeffrey H. Kozlow, Adeyiza Momoh, Michael Englesbe, Chad Brummett, Jennifer F. Waljee. 2017. Prescription Opioid Use among Opioid-Naive Women Undergoing Immediate Breast Reconstruction. *Plastic and Reconstructive Surgery* 140:6, 1081-1090. [Crossref]
- 9. Kavita Parikh, Jay Berry, Matt Hall, Grant M. Mussman, Amanda Montalbano, Joanna Thomson, Rustin Morse, Karen M. Wilson, Samir S. Shah. 2017. Racial and Ethnic Differences in Pediatric Readmissions for Common Chronic Conditions. *The Journal of Pediatrics* 186, 158-164.e1. [Crossref]
- 10. Micaela A. Finnegan, Robyn Shaffer, Austin Remington, Jereen Kwong, Catherine Curtin, Tina Hernandez-Boussard. 2017. Emergency Department Visits Following Elective Total Hip and Knee Replacement Surgery. The Journal of Bone and Joint Surgery 99:12, 1005-1012. [Crossref]
- 11. Atwood D. Gaines. Sociocultural Construction of Medical Knowledge 612-616. [Crossref]
- 12. Max O. Krucoff, Steven Cook, Owoicho Adogwa, Jessica Moreno, Siyun Yang, Jichun Xie, Alexander O. Firempong, Nandan Lad, Carlos A. Bagley. 2017. Racial, Socioeconomic, and Gender Disparities in the Presentation, Treatment, and Outcomes of Adult Chiari I Malformations. World Neurosurgery 97, 431-437. [Crossref]
- 13. Raphaël Charron-Chénier, Joshua J. Fink, Lisa A. Keister. 2017. Race and Consumption. Sociology of Race and Ethnicity 3:1, 50-67. [Crossref]
- 14. Geesje Van Woerden, Crispijn L. Van Den Brand, Cornelis F. Den Hartog, Floris J. Idenburg, Diana C. Grootendorst, M. Christien Van Der Linden. 2016. Increased analgesia administration in emergency medicine after implementation of revised guidelines. International Journal of Emergency Medicine 9:1. . [Crossref]
- 15. Jing Feng, Joseph P. Iser, Wei Yang. 2016. Medical encounters for opioid-related intoxications in Southern Nevada: sociodemographic and clinical correlates. *BMC Health Services Research* 16:1. . [Crossref]
- Y. Jenny Xiao, Géraldine Coppin, Jay J. Van Bavel. 2016. Perceiving the World Through Group-Colored Glasses: A Perceptual Model of Intergroup Relations. Psychological Inquiry 27:4, 255-274. [Crossref]
- 17. Nour S. Kteily, Jennifer A. Richeson. 2016. Perceiving the World Through Hierarchy-Shaped Glasses: On the Need to Embed Social Identity Effects on Perception Within the Broader Context of Intergroup Hierarchy. *Psychological Inquiry* 27:4, 327-334. [Crossref]
- 18. John W. Barrington, Scott T. Lovald, Kevin L. Ong, Heather N. Watson, Roger H. Emerson. 2016. How Do Demographic, Surgical, Patient, and Cultural Factors Affect Pain Control After Unicompartmental Knee Arthroplasty? A Multivariable Regression Analysis. *The Journal of Arthroplasty* 31:9, 97-101. [Crossref]
- 19. Astha Singhal, Yu-Yu Tien, Renee Y. Hsia. 2016. Racial-Ethnic Disparities in Opioid Prescriptions at Emergency Department Visits for Conditions Commonly Associated with Prescription Drug Abuse. *PLOS ONE* 11:8, e0159224. [Crossref]

- 20. Claire E. Ashton-James, Michael K. Nicholas. 2016. Appearance of trustworthiness. PAIN 157:8, 1583-1585. [Crossref]
- 21. Amir M. Mohareb, Andrea F. Dugas, Yu-Hsiang Hsieh. 2016. Changing epidemiology and management of infectious diseases in US EDs. *The American Journal of Emergency Medicine* 34:6, 1059-1065. [Crossref]
- 22. Jacob M. Vigil, Patrick Coulombe, Joe Alcock, Eric Kruger, Sarah S. Stith, Chance Strenth, Mark Parshall, Sara B. Cichowski. 2016. Patient Ethnicity Affects Triage Assessments and Patient Prioritization in U.S. Department of Veterans Affairs Emergency Departments. *Medicine* 95:14, e3191. [Crossref]
- 23. Alice Pressman, Alice Jacobson, Roderick Eguilos, Amy Gelfand, Cynthia Huynh, Luisa Hamilton, Andrew Avins, Nandini Bakshi, Kathleen Merikangas. 2016. Prevalence of migraine in a diverse community—electronic methods for migraine ascertainment in a large integrated health plan. *Cephalalgia* 36:4, 325-334. [Crossref]
- 24. John Hoberman. 2016. Why Bioethics Has a Race Problem. Hastings Center Report 46:2, 12-18. [Crossref]
- 25. John M. McLaughlin, Angela Lambing, Michelle L. Witkop, Terry L. Anderson, James Munn, Bartholomew Tortella. 2016. Racial Differences in Chronic Pain and Quality of Life among Adolescents and Young Adults with Moderate or Severe Hemophilia. *Journal of Racial and Ethnic Health Disparities* 3:1, 11-20. [Crossref]
- 26. Adel Alrwisan, Efe Eworuke. 2016. Are Discrepancies in Waiting Time for Chest Pain at Emergency Departments between African Americans and Whites Improving Over Time?. *The Journal of Emergency Medicine* **50**:2, 349-355. [Crossref]
- 27. Adil A. Shah, Cheryl K. Zogg, Syed Nabeel Zafar, Eric B. Schneider, Lisa A. Cooper, Alyssa B. Chapital, Susan M. Peterson, Joaquim M. Havens, Roland J. Thorpe, Debra L. Roter, Renan C. Castillo, Ali Salim, Adil H. Haider. 2015. Analgesic Access for Acute Abdominal Pain in the Emergency Department Among Racial/Ethnic Minority Patients. *Medical Care* 53:12, 1000-1009. [Crossref]
- 28. Robin Richardson, Thomas Charters, Nicholas King, Sam Harper. 2015. Trends in Educational Inequalities in Drug Poisoning Mortality: United States, 1994–2010. *American Journal of Public Health* 105:9, 1859-1865. [Abstract] [Full Text] [PDF] [PDF Plus] [Supplemental Material]
- 29. Ryan E. Childers, J. Lucas Williams, Amnon Sonnenberg. 2015. Practice patterns of sedation for colonoscopy. *Gastrointestinal Endoscopy* 82:3, 503-511. [Crossref]
- 30. Ashley C. Rondini. 2015. [Mis]Placing Race in Biomedical Clinical Context: Racial Categories, Medical Research, and the Reproduction of Health Disparities. *Sociology Compass* 9:8, 704-717. [Crossref]
- 31. Brian S Marcus, Arvind Venkat. 2015. Ethical pain management in the emergency department: the canary in the coal mine. *Pain Management* 5:4, 251-260. [Crossref]
- 32. Adil H. Haider, Eric B. Schneider, N. Sriram, Valerie K. Scott, Sandra M. Swoboda, Cheryl K. Zogg, Nitasha Dhiman, Elliott R. Haut, David T. Efron, Peter J. Pronovost, Julie A. Freischlag, Pamela A. Lipsett, Edward E. Cornwell, Ellen J. MacKenzie, Lisa A. Cooper. 2015. Unconscious Race and Class Biases among Registered Nurses: Vignette-Based Study Using Implicit Association Testing. *Journal of the American College of Surgeons* 220:6, 1077-1086.e3. [Crossref]
- 33. Jacob M. Vigil, Joe Alcock, Patrick Coulombe, Laurie McPherson, Mark Parshall, Allison Murata, Heather Brislen. 2015. Ethnic Disparities in Emergency Severity Index Scores among U.S. Veteran's Affairs Emergency Department Patients. PLOS ONE 10:5, e0126792. [Crossref]
- 34. Madhulika A. Gupta, Aditya K. Gupta, Branka Vujcic, Meghan Piccinin. 2015. Use of opioid analgesics in skin disorders: Results from a nationally representative US sample. *Journal of Dermatological Treatment* 26:3, 269-274. [Crossref]
- 35. Nicole A. Hollingshead, Marianne S. Matthias, Matthew J. Bair, Adam T. Hirsh. 2015. Impact of Race and Sex on Pain Management by Medical Trainees: A Mixed Methods Pilot Study of Decision Making and Awareness of Influence. *Pain Medicine* 16:2, 280-290. [Crossref]
- 36. Paul E. Hilliard. 2015. The Prescription Opioid Abuse Crisis in America. Advances in Anesthesia 33:1, 175-189. [Crossref]
- 37. Jacob P Prunuske, Catherine A St. Hill, Keri D Hager, Andrine M Lemieux, Michael T Swanoski, Grant W Anderson, M Nawal Lutfiyya. 2014. Opioid prescribing patterns for non-malignant chronic pain for rural versus non-rural US adults: a population-based study using 2010 NAMCS data. *BMC Health Services Research* 14:1. . [Crossref]
- 38. Shokoofeh Jamshidi, Fahimeh Baghaei, Zakie Doniavi, Alireza Jalalvand, Saeid Moosavi, Shahrbanoo Radi. 2014. Attitude of Dentists Towards the Administration of Analgesics for Management of Post-Endodontic Pain in Hamadan. *Avicenna Journal of Dental Research* 6:2. . [Crossref]
- 39. Tiffani J. Johnson, Brian R. Schultz, Francis X. Guyette. 2014. Characterizing Analgesic Use during Air Medical Transport of Injured Children. *Prehospital Emergency Care* 18:4, 531-538. [Crossref]

- 40. Philip Craven, Orhan Cinar, David Fosnocht, Jessica Carey, Adrienne Carey, LeGrand Rogers, Kajsa Vlasic, Troy Madsen. 2014. Prospective, 10-year evaluation of the impact of Hispanic ethnicity on pain management practices in the ED. *The American Journal of Emergency Medicine* 32:9, 1055-1059. [Crossref]
- 41. Adil H. Haider, Eric B. Schneider, N Sriram, Deborah S. Dossick, Valerie K. Scott, Sandra M. Swoboda, Lia Losonczy, Elliott R. Haut, David T. Efron, Peter J. Pronovost, Julie A. Freischlag, Pamela A. Lipsett, Edward E. Cornwell, Ellen J. MacKenzie, Lisa A. Cooper. 2014. Unconscious race and class bias. *Journal of Trauma and Acute Care Surgery* 77:3, 409-416. [Crossref]
- 42. Christopher Okunseri, Elaye Okunseri, Qun Xiang, Joshua M. Thorpe, Aniko Szabo. 2014. Prescription of opioid and nonopioid analgesics for dental care in emergency departments: Findings from the National Hospital Ambulatory Medical Care Survey. *Journal of Public Health Dentistry* 74:4, 283-292. [Crossref]
- 43. Nicholas B. King, Veronique Fraser, Constantina Boikos, Robin Richardson, Sam Harper. 2014. Determinants of Increased Opioid-Related Mortality in the United States and Canada, 1990–2013: A Systematic Review. *American Journal of Public Health* 104:8, e32-e42. [Abstract] [Full Text] [PDF] [PDF Plus] [Supplemental Material]
- 44. Allison E. Infinger, Jonathan R. Studnek. 2014. An Assessment of Pain Management Among Patients Presenting to Emergency Medical Services After Suffering a Fall. *Prehospital and Disaster Medicine* 29:04, 344-349. [Crossref]
- 45. Diana J. Burgess, Sean Phelan, Michael Workman, Emily Hagel, David B. Nelson, Steven S. Fu, Rachel Widome, Michael van Ryn. 2014. The Effect of Cognitive Load and Patient Race on Physicians' Decisions to Prescribe Opioids for Chronic Low Back Pain: A Randomized Trial: Table 1. *Pain Medicine* 15:6, 965-974. [Crossref]
- 46. Susan W. Hinze. Race and Class, Intersections of 1-6. [Crossref]
- 47. Nancy N. Maserejian, Michael A. Fischer, Felicia L. Trachtenberg, Jing Yu, Lisa D. Marceau, John B. McKinlay, Jeffrey N. Katz. 2014. Variations Among Primary Care Physicians in Exercise Advice, Imaging, and Analgesics for Musculoskeletal Pain: Results From a Factorial Experiment. *Arthritis Care & Research* 66:1, 147-156. [Crossref]
- 48. Melanie Johnson, Katrina Richardson, Suha Al-Oballi Kridli. 2014. Disparities in Pain Management in the Emergency Department: An Integrative Review. *Open Journal of Nursing* 04:08, 608-619. [Crossref]
- 49. Richard A. Epstein, William V. Bobo, Peter R. Martin, James A. Morrow, Wei Wang, Rameela Chandrasekhar, William O. Cooper. 2013. Increasing pregnancy-related use of prescribed opioid analgesics. *Annals of Epidemiology* 23:8, 498-503. [Crossref]
- 50. Megann F. Young, H. Gene Hern, Harrison J. Alter, Joseph Barger, Farnaz Vahidnia. 2013. Racial Differences in Receiving Morphine among Prehospital Patients with Blunt Trauma. *The Journal of Emergency Medicine* 45:1, 46-52. [Crossref]
- 51. Arvind Venkat, Christian Fromm, Eric Isaacs, Jordan Ibarra. 2013. An Ethical Framework for the Management of Pain in the Emergency Department. *Academic Emergency Medicine* 20:7, 716-723. [Crossref]
- 52. Catherine A. Marco, William Kanitz, Matthew Jolly. 2013. Pain Scores among Emergency Department (ED) Patients: Comparison by ED Diagnosis. *The Journal of Emergency Medicine* 44:1, 46-52. [Crossref]
- 53. Leslie R.M. Hausmann, Shasha Gao, Edward S. Lee, Kent C. Kwoh. 2013. Racial disparities in the monitoring of patients on chronic opioid therapy. *Pain* 154:1, 46-52. [Crossref]
- 54. Stefan Stremersch, Vardit Landsman, Sriram Venkataraman. 2013. The Relationship Between DTCA, Drug Requests, and Prescriptions: Uncovering Variation in Specialty and Space. *Marketing Science* 32:1, 89-110. [Crossref]
- 55. Laurie Jowers Ware, Cynthia D. Epps, Julie Clark, Ayona Chatterjee. 2012. Do Ethnic Differences Still Exist in Pain Assessment and Treatment in the Emergency Department?. *Pain Management Nursing* 13:4, 194-201. [Crossref]
- 56. Chu-Lin Tsai, Ashley F. Sullivan, James A. Gordon, Rainu Kaushal, David J. Magid, David Blumenthal, Carlos A. Camargo. 2012. Racial/ethnic differences in emergency care for joint dislocation in 53 US EDs. The American Journal of Emergency Medicine 30:9, 1970-1980. [Crossref]
- 57. Steven M. Green. 2012. There Is Oligo-Evidence for Oligoanalgesia. Annals of Emergency Medicine 60:2, 212-214. [Crossref]
- 58. Nathalia Jimenez, Gail D. Anderson, Danny D. Shen, Susan S. Nielsen, Federico M. Farin, Kristy Seidel, Anne M. Lynn. 2012. Is ethnicity associated with morphine's side effects in children? Morphine pharmacokinetics, analgesic response, and side effects in children having tonsillectomy. *Pediatric Anesthesia* 22:7, 669-675. [Crossref]
- 59. Christopher Okunseri, Elaye Okunseri, Joshua M. Thorpe, Qun Xiang, Aniko Szabo. 2012. Medications Prescribed in Emergency Departments for Nontraumatic Dental Condition Visits in the United States. *Medical Care* 50:6, 508-512. [Crossref]
- 60. Ptlene Minick, Patricia C. Clark, Jo Ann Dalton, Eva Horne, Debbie Greene, Monica Brown. 2012. Long-Bone Fracture Pain Management in the Emergency Department. *Journal of Emergency Nursing* 38:3, 211-217. [Crossref]
- 61. Timothy F. Platts-Mills, Katie M. Hunold, Andrey V. Bortsov, April C. Soward, David A. Peak, Jeffrey S. Jones, Robert A. Swor, David C. Lee, Robert M. Domeier, Phyllis L. Hendry, Niels K. Rathlev, Samuel A. McLean. 2012. More educated emergency department patients are less likely to receive opioids for acute pain. *Pain* 153:5, 967-973. [Crossref]

- 62. V. M. Banz, B. Christen, K. Paul, L. Martinolli, D. Candinas, H. Zimmermann, A. K. Exadaktylos. 2012. Gender, age and ethnic aspects of analgesia in acute abdominal pain: is analgesia even across the groups?. *Internal Medicine Journal* 42:3, 281-288. [Crossref]
- 63. Salimah H. Meghani, Eeeseung Byun, Rollin M. Gallagher. 2012. Time to Take Stock: A Meta-Analysis and Systematic Review of Analgesic Treatment Disparities for Pain in the United States. *Pain Medicine* 13:2, 150-174. [Crossref]
- 64. Consuelo M. Flores, Diane C. Zelman, Yvette Flores. 2012. "I Have Not a Want But a Hunger to Feel No Pain" Mexican Immigrant Women with Chronic Pain: Narratives and Psychotherapeutic Implications. Women & Therapy 35:1-2, 31-44. [Crossref]
- 65. Marion R. Sills, Diane L. Fairclough, Daksha Ranade, Michael S. Mitchell, Michael G. Kahn. 2011. Emergency Department Crowding Is Associated with Decreased Quality of Analgesia Delivery for Children with Pain Related to Acute, Isolated, Longbone Fractures. *Academic Emergency Medicine* 18:12, 1330-1338. [Crossref]
- 66. Molly J. Dingel, Katrina Karkazis, Barbara A. Koenig. 2011. Framing Nicotine Addiction as a "Disease of the Brain": Social and Ethical Consequences. *Social Science Quarterly* 5, n/a-n/a. [Crossref]
- 67. Diana J. Burgess. 2011. Addressing Racial Healthcare Disparities: How Can We Shift the Focus from Patients to Providers?. *Journal of General Internal Medicine* 26:8, 828-830. [Crossref]
- 68. Jana M. Mossey. 2011. Defining Racial and Ethnic Disparities in Pain Management. Clinical Orthopaedics and Related Research® 469:7, 1859-1870. [Crossref]
- 69. Atul F. Kamath, Mary I. O'Connor. 2011. Breakout Session: Gender and Ethnic Disparities in Pain Management. Clinical Orthopaedics and Related Research® 469:7, 1962-1966. [Crossref]
- 70. Kimberly T. Sibille, Lindsay L. Kindler, Toni L. Glover, Ricardo D. Gonzalez, Roland Staud, Joseph L. Riley, Roger B. Fillingim. 2011. Individual Differences in Morphine and Butorphanol Analgesia: A Laboratory Pain Study. *Pain Medicine* 12:7, 1076-1085. [Crossref]
- 71. Elisabeth K. Stephens, Phuong L. Nguyen, Aminah Jatoi. 2011. Practical Strategies to Circumvent and/or Surmount Six Barriers in Health Disparities Research: Observations from a Pilot Project Among Vietnamese Americans. *Journal of Cancer Education* 26:2, 243-247. [Crossref]
- 72. Brian B. Drwecki, Colleen F. Moore, Sandra E. Ward, Kenneth M. Prkachin. 2011. Reducing racial disparities in pain treatment: The role of empathy and perspective-taking. *Pain* 152:5, 1001-1006. [Crossref]
- 73. Andrei Barasch, Monika M. Safford, Sandre F. McNeal, Michelle Robinson, Vivian S. Grant, Gregg H. Gilbert. 2011. Patterns of postoperative pain medication prescribing after invasive dental procedures. *Special Care in Dentistry* 31:2, 53-57. [Crossref]
- 74. Constance L. Monitto, Sabine Kost-Byerly, Myron Yaster. Pain Management 418-451. [Crossref]
- 75. Atul F. Kamath, John G. Horneff, Vandy Gaffney, Craig L. Israelite, Charles L. Nelson. 2010. Ethnic and Gender Differences in the Functional Disparities after Primary Total Knee Arthroplasty. *Clinical Orthopaedics and Related Research*® 468:12, 3355-3361. [Crossref]
- 76. Lynne Neumayr, Steven Pringle, Stephen Giles, Keith C. Quirolo, Susan Paulukonis, Elliott P. Vichinsky, Marsha J. Treadwell. 2010. Chart Card: Feasibility of a Tool for Improving Emergency Department Care in Sickle Cell Disease. *Journal of the National Medical Association* 102:11, 1017-1024. [Crossref]
- 77. William T. Zempsky, Kristin A. Loiselle, Kathleen McKay, Brian H. Lee, J. Nathan Hagstrom, Neil L. Schechter. 2010. Do Children with Sickle Cell Disease Receive Disparate Care for Pain in the Emergency Department?. *The Journal of Emergency Medicine* 39:5, 691-695. [Crossref]
- 78. 2010. Optimizing the Treatment of Pain in Patients with Acute Presentations. Annals of Emergency Medicine 56:1, 77-79. [Crossref]
- 79. Alan Heins, Peter Homel, Basmah Safdar, Knox Todd. 2010. Physician Race/Ethnicity Predicts Successful Emergency Department Analgesia. *The Journal of Pain* 11:7, 692-697. [Crossref]
- 80. Steven A. Iacono. 2010. Dialysis clinician bias regarding patient non-compliance. *Dialysis & Transplantation* **39**:6, 247-253. [Crossref]
- 81. Erlinda Wheeler, Thomas Hardie, Paula Klemm, Ibironke Akanji, Erin Schonewolf, Janna Scott, Beth Sterling. 2010. Level of Pain and Waiting Time in the Emergency Department. *Pain Management Nursing* 11:2, 108-114. [Crossref]
- 82. Virginia Dzul-Church, Jenica W. Cimino, Shelley R. Adler, Piera Wong, Wendy G. Anderson. 2010. "I'm Sitting Here By Myself...": Experiences of Patients with Serious Illness at an Urban Public Hospital. *Journal of Palliative Medicine* 13:6, 695-701. [Crossref]

- 83. Timothy S. Carey, Janet K. Freburger, G. Mark Holmes, Anne Jackman, Stefanie Knauer, Andrea Wallace, Jane Darter. 2010. Race, Care Seeking, and Utilization for Chronic Back and Neck Pain: Population Perspectives. *The Journal of Pain* 11:4, 343-350. [Crossref]
- 84. Carlos J. Lavernia, Jose C. Alcerro, Mark D. Rossi. 2010. Fear in Arthroplasty Surgery: The Role of Race. *Clinical Orthopaedics and Related Research*® 468:2, 547-554. [Crossref]
- 85. James R. Miner, Paul M. Paris, Donald M. Yealy. Pain Management 2410-2428. [Crossref]
- 86. Edward Bernstein, Judith Bernstein, Thea James. Multiculturalism and Care Delivery 2540-2546. [Crossref]
- 87. Kai'En Leong, Tracey J. Weiland, Andrew W. Dent. 2010. Exploring beliefs of the four major ethnic groups in Melbourne regarding healthcare and treatment. *Australian Health Review* 34:4, 458. [Crossref]
- 88. Karen O. Anderson, Carmen R. Green, Richard Payne. 2009. Racial and Ethnic Disparities in Pain: Causes and Consequences of Unequal Care. *The Journal of Pain* 10:12, 1187-1204. [Crossref]
- 89. Megan Crowley-Matoka, Somnath Saha, Steven K. Dobscha, Diana J. Burgess. 2009. Problems of Quality and Equity in Pain Management: Exploring the Role of Biomedical Culture. *Pain Medicine* 10:7, 1312-1324. [Crossref]
- 90. Richard Brown, Jeremy Furyk. 2009. Racial disparities in health care—emergency department management of minor head injury. *International Journal of Emergency Medicine* 2:3, 161-166. [Crossref]
- 91. R. Daniulaityte, R.S. Falck, J. Wang, R.G. Carlson. 2009. Illicit use of pharmaceutical opioids among young polydrug users in Ohio. *Addictive Behaviors* 34:8, 649-653. [Crossref]
- 92. BD Heckman, KA Holroyd, G Tietjen, FJ O'Donnell, L Himawan, C Utley, R Watakakosol, M Stillman. 2009. Whites and African-Americans in Headache Specialty Clinics Respond Equally Well to Treatment. *Cephalalgia* 29:6, 650-661. [Crossref]
- 93. Alec B. O'Connor, Frank L. Zwemer, Daniel P. Hays, Changyong Feng. 2009. Outcomes After Intravenous Opioids in Emergency Patients: A Prospective Cohort Analysis. *Academic Emergency Medicine* 16:6, 477-487. [Crossref]
- 94. Adam T. Hirsh, Steven Z. George, Michael E. Robinson. 2009. Pain assessment and treatment disparities: A virtual human technology investigation. *Pain* 143:1, 106-113. [Crossref]
- 95. Basmah Safdar, Alan Heins, Peter Homel, James Miner, Martha Neighbor, Paul DeSandre, Knox H. Todd. 2009. Impact of Physician and Patient Gender on Pain Management in the Emergency Department—A Multicenter Study. *Pain Medicine* 10:2, 364-372. [Crossref]
- 96. Chu-Lin Tsai, Carlos A Camargo Jr. 2009. Racial and Ethnic Differences in Emergency Care for Acute Exacerbation of Chronic Obstructive Pulmonary Disease. *Academic Emergency Medicine* 16:2, 108-115. [Crossref]
- 97. Susan W. Hinze, Noah J. Webster, Heidi T. Chirayath, Joshua H. Tamayo-Sarver. Hurt running from police? No chance of (pain) relief: The social construction of deserving patients in emergency departments 235-261. [Crossref]
- 98. Florence T. Bourgeois, Michael W. Shannon, Anne M. Stack. 2008. "Left Without Being Seen": A National Profile of Children Who Leave the Emergency Department Before Evaluation. *Annals of Emergency Medicine* **52**:6, 599-605. [Crossref]
- 99. Adam J. Singer, Gregory Garra, Jasmine K. Chohan, Charles Dalmedo, Henry C. Thode. 2008. Triage Pain Scores and the Desire for and Use of Analgesics. *Annals of Emergency Medicine* **52**:6, 689-695. [Crossref]
- 100. Diana Jill Burgess, Megan Crowley-Matoka, Sean Phelan, John F. Dovidio, Robert Kerns, Craig Roth, Somnath Saha, Michelle van Ryn. 2008. Patient race and physicians' decisions to prescribe opioids for chronic low back pain. *Social Science & Medicine* 67:11, 1852-1860. [Crossref]
- 101. Polly Bijur, Anick Bérard, David Esses, Yvette Calderon, E. John Gallagher. 2008. Race, Ethnicity, and Management of Pain from Long-bone Fractures: A Prospective Study of Two Academic Urban Emergency Departments. *Academic Emergency Medicine* 15:7, 589-597. [Crossref]
- 102. Alex Y. Chen, Susan Wu. 2008. Dispensing Pattern of Generic and Brand-Name Drugs in Children. *Ambulatory Pediatrics* 8:3, 189-194. [Crossref]
- 103. Polly Bijur, Anick Bérard, Jordan Nestor, Yvette Calderon, Michelle Davitt, E. John Gallagher. 2008. No racial or ethnic disparity in treatment of long-bone fractures. *The American Journal of Emergency Medicine* 26:3, 270-274. [Crossref]
- 104. Cynthia D. Epps, Laurie Jowers Ware, Abbot Packard. 2008. Ethnic Wait Time Differences in Analgesic Administration in the Emergency Department. *Pain Management Nursing* **9**:1, 26-32. [Crossref]
- 105. Alan K. Geller, M. Kevin O'Connor. 2008. The Sickle Cell Crisis: A Dilemma in Pain Relief. *Mayo Clinic Proceedings* 83:3, 320-323. [Crossref]
- 106. Melanie Arthur, Jerris R. Hedges, Craig D. Newgard, Brian S. Diggs, Richard J. Mullins. 2008. Racial Disparities in Mortality Among Adults Hospitalized After Injury. *Medical Care* 46:2, 192-199. [Crossref]

- 107. A.D. Gaines. Socio-Cultural Construction of Medical Knowledge 114-119. [Crossref]
- 108. Serge Perrot, Victor Concas, Francis Allaert, Françoise Laroche. 2008. Évaluation de la prise de décision de prescription d'un antalgique en prise quotidienne unique ou multiple dans les rachialgies aiguës. *La Presse Médicale* 37:1, 14-20. [Crossref]
- 109. Jeffrey T. Berger. 2008. The Influence of Physicians' Demographic Characteristics and Their Patients' Demographic Characteristics on Physician Practice: Implications for Education and Research. *Academic Medicine* 83:1, 100-105. [Crossref]
- 110. Ruth A. Griffin, Denise F. Polit, Mary W. Byrne. 2007. Stereotyping and nurses' recommendations for treating pain in hospitalized children. *Research in Nursing & Health* 30:6, 655-666. [Crossref]
- 111. Glen E. Michael, Karl A. Sporer, Glen M. Youngblood. 2007. Women are less likely than men to receive prehospital analgesia for isolated extremity injuries. *The American Journal of Emergency Medicine* 25:8, 901-906. [Crossref]
- 112. Isabelle Decosterd, Olivier Hugli, Emmanuel Tamchès, Catherine Blanc, Elyazid Mouhsine, Jean-Claude Givel, Bertrand Yersin, Thierry Buclin. 2007. Oligoanalgesia in the Emergency Department: Short-Term Beneficial Effects of an Education Program on Acute Pain. Annals of Emergency Medicine 50:4, 462-471. [Crossref]
- 113. Jerzy P. Szaflarski, Jason M. Meckler, Magdalena Szaflarski, Lori A. Shutter, Michael D. Privitera, Stephen L. Yates. 2007. Levetiracetam use in critically ill patients. *Neurocritical Care* 7:2, 140-147. [Crossref]
- 114. Hillary Veda Kunins, Eran Bellin, Cynthia Chazotte, Evelyn Du, Julia Hope Arnsten. 2007. The Effect of Race on Provider Decisions to Test for Illicit Drug Use in The Peripartum Setting. *Journal of Women's Health* 16:2, 245-255. [Crossref]
- 115. Tamara S. Ritsema, Gabor D. Kelen, Peter J. Pronovost, Julius Cuong Pham. 2007. The National Trend in Quality of Emergency Department Pain Management for Long Bone Fractures. *Academic Emergency Medicine* 14:2, 163-169. [Crossref]
- 116. Joanne Lafleur, Qayyim Said, Carrie McAdam-Marx, Kenneth Jackson, Maysam Mortazavi. 2007. Problems in Studying the Association Between Race and Pain in Outcomes Research. *Journal of Pain & Palliative Care Pharmacotherapy* 21:3, 57-62. [Crossref]
- 117. Laurent G. Glance, Richard . Wissler, Christopher Glantz, Turner M. Osler, Dana B. Mukamel, Andrew W. Dick. 2007. Racial Differences in the Use of Epidural Analgesia for Labor. *Anesthesiology* 106:1, 19-25. [Crossref]
- 118. Gregg Husk, Saadia Akhtar. 2007. Chief Complaints, Emergency Department Clinical Documentation Systems, and the Challenge of Dealing with the Patient's Own Words. *Academic Emergency Medicine* 14:1, 69-73. [Crossref]
- 119. Alex D. Federman, Ann Litke, R. Sean Morrison. 2006. Association of age with analgesic use for back and joint disorders in outpatient settings. *The American Journal of Geriatric Pharmacotherapy* 4:4, 306-315. [Crossref]
- 120. Alexie Cintron, R. Sean Morrison. 2006. Pain and Ethnicity in the United States: A Systematic Review. *Journal of Palliative Medicine* 9:6, 1454–1473. [Crossref]
- 121. Constance Liu, Brook Watts, David Litaker. 2006. Access to and utilization of healthcare: the provider's role. Expert Review of Pharmacoeconomics & Outcomes Research 6:6, 653-660. [Crossref]
- 122. Sheryl L. Heron, Edward Stettner, Leon L. Haley. 2006. Racial and Ethnic Disparities in the Emergency Department: A Public Health Perspective. *Emergency Medicine Clinics of North America* 24:4, 905-923. [Crossref]
- 123. Bert D. Stover, Judith A. Turner, Gary Franklin, Jeremy V. Gluck, Deborah Fulton-Kehoe, Lianne Sheppard, Thomas M. Wickizer, Joel Kaufman, Kathleen Egan. 2006. Factors Associated With Early Opioid Prescription Among Workers With Low Back Injuries. *The Journal of Pain* 7:10, 718-725. [Crossref]
- 124. Marilyn G. Foreman, Sandra K. Willsie. 2006. Health Care Disparities in Critical Illness. *Clinics in Chest Medicine* 27:3, 473-486. [Crossref]
- 125. Gerda G. Fillenbaum, Joseph T. Hanlon. 2006. Racial and ethnic disparities in medication use among older adults. *The American Journal of Geriatric Pharmacotherapy* 4:2, 93-95. [Crossref]
- 126. Janet Kaye Heins, Alan Heins, Marianthe Grammas, Melissa Costello, Kun Huang, Satya Mishra. 2006. Disparities in Analgesia and Opioid Prescribing Practices for Patients With Musculoskeletal Pain in the Emergency Department. *Journal of Emergency Nursing* 32:3, 219-224. [Crossref]
- 127. Denise J. Drevdahl, Debby A. Philips, Janette Y. Taylor. 2006. Uncontested categories: the use of race and ethnicity variables in nursing research. *Nursing Inquiry* 13:1, 52-63. [Crossref]
- 128. Diana J. Burgess, Michelle Van Ryn, Megan Crowley-Matoka, Jennifer Malat. 2006. Understanding the Provider Contribution to Race/Ethnicity Disparities in Pain Treatment: Insights from Dual Process Models of Stereotyping. *Pain Medicine* 7:2, 119-134. [Crossref]
- 129. D A Paul, R Locke, K Zook, K H Leef, J L Stefano, G Colmorgen. 2006. Racial differences in prenatal care of mothers delivering very low birth weight infants. *Journal of Perinatology* 26:2, 74-78. [Crossref]

- 130. L. L. Haley. 2006. Stuck in Neutral: Continued Challenges with Healthcare Disparities. *Academic Emergency Medicine* 13:2, 191-194. [Crossref]
- 131. Hester J. Lipscomb, Dana Loomis, Mary Anne McDonald, Robin A. Argue, Steve Wing. 2006. A Conceptual Model of Work and Health Disparities in the United States. *International Journal of Health Services* 36:1, 25-50. [Crossref]
- 132. Sandra Soo-Jin Lee. 2005. Racializing Drug Design: Implications of Pharmacogenomics for Health Disparities. *American Journal of Public Health* 95:12, 2133-2138. [Abstract] [Full Text] [PDF] [PDF Plus]
- 133. James Slover, Jennifer Gibson, Tor Tosteson, Brian Smith, Kenneth Koval. 2005. Racial and Economic Disparity and the Treatment of Pediatric Fractures. *Journal of Pediatric Orthopaedics* 25:6, 717-721. [Crossref]
- 134. Susan J. Appel, Joyce Newman Giger, Ruth Elaine Davidhizar. 2005. Opportunity Cost. *The Journal of Cardiovascular Nursing* 20:5, 315-324. [Crossref]
- 135. Daniel B. Carr, David H. Reines, Jonathan Schaffer, Rosemary C. Polomano, Stephen Lande. 2005. The Impact of Technology on the Analgesic Gap and Quality of Acute Pain Management. *Regional Anesthesia and Pain Medicine* 30:3, 286-291. [Crossref]
- 136. Salimah H. Meghani. 2005. Leveling the Playing Field: Does Pain Disparity Literature Suffer from a Reporting Bias?. *Pain Medicine* 6:3, 269-270. [Crossref]
- 137. Andrew Sucov, Andrew Nathanson, Jackie McCormick, Lawrence Proano, Steven E. Reinert, Gregory Jay. 2005. Peer Review and Feedback Can Modify Pain Treatment Patterns for Emergency Department Patients With Fractures. *American Journal of Medical Quality* 20:3, 138-143. [Crossref]
- 138. Carol S. Weisse, Kemoy K. Foster, Elizabeth A. Fisher. 2005. The Influence of Experimenter Gender and Race on Pain Reporting: Does Racial or Gender Concordance Matter?. *Pain Medicine* 6:1, 80-87. [Crossref]
- 139. Louis W. Sullivan, Barry A. Eagel. 2005. Leveling the Playing Field: Recognizing and Rectifying Disparities in Management of Pain. *Pain Medicine* 6:1, 5-10. [Crossref]
- 140. Xuemei Luo, Ricardo Pietrobon, Lesley H. Curtis, Lloyd A. Hey. 2004. Prescription of Nonsteroidal Anti-inflammatory Drugs and Muscle Relaxants for Back Pain in the United States. *Spine* 29:23, E531-E537. [Crossref]
- 141. Diana D. Cardenas, Thomas N. Bryce, Kazuko Shem, J.Scott Richards, Hanaa Elhefni. 2004. Gender and minority differences in the pain experience of people with spinal cord injury11No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the authors or upon any organization with which the authors are associated. *Archives of Physical Medicine and Rehabilitation* 85:11, 1774-1781. [Crossref]
- 142. Polly Gerber Zimmermann. 2004. Tips for Managing Pain More Effectively. *Journal of Emergency Nursing* 30:5, 470-472. [Crossref]
- 143. Joshua H. Tamayo-Sarver, Neal V. Dawson, Rita K. Cydulka, Robert S. Wigton, David W. Baker. 2004. Variability in emergency physician decisionmaking about prescribing opioid analgesics. *Annals of Emergency Medicine* 43:4, 483-493. [Crossref]
- 144. Augustine J Kposowa, Glenn T Tsunokai. 2002. Searching for relief: racial differences in treatment of patients with back pain. *Race and Society* 5:2, 193-223. [Crossref]
- 145. Joshua H. Tamayo-Sarver, Neal V. Dawson, Susan W. Hinze, Rita K. Cydulka, Robert S. Wigton, David W. Baker. Rapid Clinical Decisions in Context: A Theoretical Model to Understand Physicians' Decision-Making With an Application to Racial/Ethnic Treatment Disparities 183-213. [Crossref]