

Notes on a reflective theory of sets

Lucius Gregory Meredith¹

Managing Partner, RTech Unchained 9336 California Ave SW, Seattle, WA 98103, USA
lgreg.meredith@gmail.com

Abstract. We describe a universe of set theories enjoying atoms without enduring infinite risk.

In the early 2000's Pitts and Gabbay's use of Fraenkl-Mostowski set theory (FM set theory), a theory of sets with atoms, to model nominal phenomena, sparked a vibrant line of research. [3] [2]. As Chaitin points out, the axioms and assumptions of a theory comprise its risk [1]. This observation raises an interesting point: is it possible to have a set theory with atoms without taking on *infinite* risk? It turns out it is.

We describe a theory of sets that enjoys both the cleanliness of **ZF**, that is arising only from the empty set, and yet with an infinite supply of atoms.

1 Intuitions

We imagine two copies of FM set theory, say a **red one** and a black one. The **red theory** has red comprehensions, $\{\dots\}$ and a red element-of relation, \in , while the black theory, correspondingly has black comprehensions, $\{\dots\}$ and a black element-of relation, \in . The red element-of relation, \in , can “see into” red sets, $\{\dots\}$, but not black ones. Meanwhile, the black element-of relation, \in , can “see into” black sets $\{\dots\}$, but not red ones. Thus, the black sets can serve as atoms for the red sets, while the red sets can serve as atoms for the black sets.

So, the two theories are mutually recursively defined. This mutual recursion is closely linked to game semantics in the sense of Abramsky, Hyland, and Ong, et al. Without loss of generality, we can think of **red** as player and black as opponent. A well-formed red set, S , is a winning strategy for player, while a well-formed black set, S , is a winning strategy for opponent.

2 Formal theory

The judgment $S[X]; \Gamma \vdash S$ is pronounced “ $S[X]$ thinks that S is well-formed, given dependencies, Γ .” Similarly, $S[X]; \Gamma \vdash x$ is pronounced “ $S[X]$ thinks that x is an admissible atom, given dependencies, Γ .”

A rule of the form

$$\frac{H_1, \dots, H_n}{C} R$$

is pronounced “ R concludes that C given H_1, \dots, H_n ”.

We use e, f, \dots to range over sets and atoms.

The rules for judging when an atom is admissible or set is well formed are as follows

$$\frac{}{\mathsf{S}[X]; () \vdash \emptyset} \textit{Foundation}$$

$$\frac{x \in X}{\mathsf{S}[X]; () \vdash \{x\}} \textit{Atomicity}$$

$$\frac{\mathsf{S}[X]; \Gamma_1 \vdash S_1 \quad \mathsf{S}[X]; \Gamma_2 \vdash S_2}{\mathsf{S}[X]; \Gamma_1, \Gamma_2 \vdash S_1 \cup S_2} \textit{Union}$$

$$\frac{\mathsf{S}[X]; \Gamma_1 \vdash S_1 \quad \mathsf{S}[X]; \Gamma_2 \vdash S_2}{\mathsf{S}[X]; \Gamma_1, \Gamma_2 \vdash S_1 \cap S_2} \textit{Intersection}$$

$$\frac{\mathsf{S}[X]; \Gamma_1 \vdash S_1 \quad \mathsf{S}[X]; \Gamma_2 \vdash S_2}{\mathsf{S}[X]; \Gamma_1, \Gamma_2 \vdash S_1 \setminus S_2} \textit{Subtraction}$$

$$\frac{\mathsf{S}[X]; \Gamma \vdash S \quad \mathsf{S}[X]; \Gamma_1 \vdash S_1 \quad \dots \quad \mathsf{S}[X]; \Gamma_n \vdash S_n \quad \mathsf{S}[X]; \Delta \vdash f : S_1 \times \dots \times S_n \rightarrow S}{\mathsf{S}[X]; \Gamma, \Gamma_1, \dots, \Gamma_n, \Delta \vdash \{f(s_1, \dots, s_n) : s_1 \in S_1, \dots, s_n \in S_n\}} \textit{Comprehension}$$

2.1 Equations

The syntactic theory is too fine grained. It makes syntactic distinctions that do not correspond to distinct sets. We erase these syntactic distinctions with a set of equations on set expressions.

$$\frac{\mathsf{S}[X]; \Gamma \vdash e}{\mathsf{S}[X]; \Gamma \vdash e = e} \textit{Identity}$$

$$\frac{\mathsf{S}[X]; \Gamma_1 \vdash_1 \quad \mathsf{S}[X]; \Gamma_2 \vdash e_2 \quad \mathsf{S}[X]; \Gamma_3 \vdash e_1 = e_2}{\mathsf{S}[X]; \Gamma_3 \vdash e_1 = e_2} \textit{Symmetry}$$

$$\frac{\mathsf{S}[X]; \Gamma_1 \vdash e_1 \quad \mathsf{S}[X]; \Gamma_2 \vdash e_2 \quad \mathsf{S}[X]; \Gamma_3 \vdash e_3 \quad \mathsf{S}[X]; \Gamma_4 \vdash e_1 = e_2 \quad \mathsf{S}[X]; \Gamma_5 \vdash e_2 = e_3}{\mathsf{S}[X]; \Gamma_4, \Gamma_5 \vdash e_1 = e_3} \textit{Transitivity}$$

$$\frac{\mathsf{S}[X]; \Gamma \vdash S}{\mathsf{S}[X]; \Gamma \vdash S \cup S = S} \textit{Idempotence}_{union}$$

$$\frac{\mathsf{S}[X]; \Gamma \vdash S}{\mathsf{S}[X]; \Gamma \vdash S \cap S = S} \textit{Idempotence}_{intersection}$$

3 Conclusions and future work

We have presented a formal theory of sets that reconciles set theory with atoms with a set theory built only from the empty set.

Acknowledgments. The author wishes to thank Jamie Gabbay for stimulating discussions that led the author to devise this theory.

References

1. G.J. Chaitin. *The Unknowable*. Discrete Mathematics and Theoretical Computer Science. Springer Singapore, 1999.
2. Ranald Clouston. Nominal lawvere theories: A category theoretic account of equational theories with names. *J. Comput. Syst. Sci.*, 80(6):1067–1086, 2014.
3. Murdoch Gabbay and Andrew M. Pitts. A new approach to abstract syntax with variable binding. *Formal Asp. Comput.*, 13(3-5):341–363, 2002.