



Contents lists available at ScienceDirect

Geochimica et Cosmochimica Acta

journal homepage: www.elsevier.com/locate/gca

Corrigendum

Corrigendum to “Isotopic fractionation accompanying CO₂ hydroxylation and carbonate precipitation from high pH waters at the Cedars, California, USA” [Geochim. Cosmochim. Acta 301 (2021) 91–115]

John N. Christensen

Earth and Environmental Science Area, Energy Geosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States

James M. Watkins

Department of Earth Sciences, University of Oregon, Eugene, OR 97403, United States

Laurent S. Devriendt

Department of Earth Sciences, University of Oregon, Eugene, OR 97403, United States

Department of Ocean System Sciences, Royal Netherlands Institute for Sea Research (NIOZ), and Utrecht University, Texel, the Netherlands

Donald J. DePaolo

Earth and Environmental Science Area, Energy Geosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States

Department of Earth and Planetary Science, University of California, Berkeley, CA 94720, United States

Mark E. Conrad, Marco Voltolini

Earth and Environmental Science Area, Energy Geosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States

Wenbo Yang

Department of Integrative Biology, University of California, Berkeley, CA 94720, United States

Wenming Dong

Earth and Environmental Science Area, Energy Geosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States

The authors regret that in the original article, the kinetic fractionation factors (KFFs) related to the CO₂ hydroxylation reaction (CO_{2(aq)} + OH[−] → HCO₃[−]) were calculated using atmospheric δ¹⁸O_{CO₂} values of 0.0 ± 0.5 ‰ (VPDB) while the correct values are 10.25 ± 0.5 ‰ (VPDB). We erroneously employed the atmospheric δ¹⁸O_{CO₂} values from the NOAA CO₂ global network (Troler et al., 1996) by not converting atmospheric δ¹⁸O_{CO₂} from the VPDB-CO₂ scale to the VPDB scale through the following expression (e.g. Srivastava and Verkouteren, 2018):

$$\delta^{18}\text{O}_{\text{VPDB}-\text{CO}_2} = (\delta^{18}\text{O}_{\text{VPDB}} - 10.25)/1.01025. \quad (1)$$

This correction affects the reported oxygen isotope KFFs for CO₂ hydroxylation to form HCO₃[−] prior to CaCO₃ precipitation. We have corrected Table 7 and Fig. 13 accordingly (see below). The composition of atmospheric CO₂ was also incorrectly plotted on the VPDB scale in Figs. 1, 8 and 15 of our original manuscript based on previous works (Fig. 8 in Clark and Fontes, 1990; Fig. 2 in Clark et al., 1992).

Importantly, the modelling outputs and main conclusions of the original article are not affected by these corrections. Nevertheless, the updated KFFs resolve an apparent discrepancy between the laboratory and field studies in our original article. Moreover, the oxygen isotope KFF for CO₂ hydroxylation now displays a positive temperature dependence over the 4–28 °C range, which was not statistically significant in the original article (Fig. 13c, r² = 0.86, n = 10, p-value <0.01):

DOI of original article: <https://doi.org/10.1016/j.gca.2021.01.003>

<https://doi.org/10.1016/j.gca.2022.09.022>

0016-7037/© 2022 Elsevier Ltd. All rights reserved.




Fig. 13. Compilation of kinetic isotope fractionation factors (KFFs) related to the CO_2 hydroxylation reaction determined from laboratory and field studies. (A) Carbon KFF estimates showing a consistent cluster of field data at around $\sim -17.0 \pm 0.3$ ‰ and more scattered KFF estimates from laboratory studies. (B) Oxygen KFF estimates expressed relative to the weighted sum of reactants ' $\text{CO}_2 + \text{OH}^-$ '. (C) Oxygen KFF* estimates expressed relative to the weighted sum of ' $\text{CO}_2 + \text{H}_2\text{O}$ '.

$$\text{KFF}^* (\text{‰}) = (0.14 \pm 0.02)T - 17.4 \pm 0.6, \quad (2)$$

where T is in $^\circ\text{C}$, the uncertainty represents one standard error, and KFF^* is the oxygen isotope kinetic fractionation factor between the HCO_3^- product and ' $\text{CO}_{2(\text{aq})} + \text{H}_2\text{O}$ ' expressed as the per mil deviation from unity:

$$\text{KFF}^* = (\alpha_{\text{HCO}_3^-} / \alpha_{\text{CO}_{2(\text{aq})} + \text{H}_2\text{O}} - 1) \cdot 10^3. \quad (3)$$

Here and in the original article, the KFF is expressed relative to the weighted sum ' $\text{CO}_{2(\text{aq})} + \text{H}_2\text{O}$ ' rather than ' $\text{CO}_{2(\text{aq})} + \text{OH}^-$ ' (i.e. the reactants) to avoid the uncertainty on the OH^- - H_2O fractionation factor (Zeebe, 2020; Bajnai and Herwartz, 2021).

$\delta^{18}\text{O}$ and oxygen KFF values in bold have been corrected by converting atmospheric $\delta^{18}\text{O}_{\text{CO}_2}$ values from the VPDB- CO_2 scale to the VPDB scale using Eq. (1).

The authors apologize for any inconvenience caused.

Table 7
Compilation of CO₂ hydroxylation KFFs.

Reference	Study type	Mineral	T (°C)	±	pH	δ ¹³ C (‰ VPDB)				Carbon KFF		δ ¹⁸ O (‰ VSMOW)						Oxygen bulk KFF			
						CO _{2(g)}	±	Mineral	±	vs CO _{2(aq)} ^h	±	H ₂ O	±	CO _{2(g)}	±	Mineral	±	vs 'CO ₂ (aq) + OH ^{-k}	±	vs 'CO _{2(aq)} + H ₂ O' (KFF*)	±
Craig (1953)	Lab	Witherite	20	3	?	-9.1	0.1	-23.1	0.1	-13.7 ⁱ	0.1	-	-	-	-	-	-	-	-	-	-
Usdowski and Hoefs (1986)	Lab	Witherite	18	1	10.0	-7.7 ^a	0.3	-25.6	0.4	-17.0	0.5	-	-	-	-	-	-	-	-	-	-
Clark et al. (1992)	Lab	Witherite	22	2	> 11.5	-10.5	0.4	-25.0	1.0	-13.9 ⁱ	1.1	-6.7	0.1	36.6	0.3	6.8	1.0	-8.5	1.4	-15.0	0.2
Clark et al. (1992)	Lab	Witherite	22	2	12.8	-45.2	0.1	-61.3	0.1	-16.3 ⁱ	0.1	-11.9	0.1	11.1	0.1	-11.2	0.1	-7.9	0.8	-14.5	0.1
Dietzel et al. (2009)	Lab	Calcite	5	1	10.5	-	-	-	-	-	-	-9.6	0.1	35.7 ^j	0.1	3.4	0.1	-10.0	0.9	-16.8	0.1
Böttcher (2018)	Lab	Witherite	4	1	12.4	-8.6 ^b	0.5	-21.4	0.1	-11.8	0.5	-7.0	0.1	41.5^b	1.0	8.3	0.1	-9.7	1.5	-16.6	0.7
Böttcher (2018)	Lab	Witherite	21	1	12.4	-8.6 ^b	0.5	-25.3	0.1	-15.8	0.5	-7.0	0.1	41.5^b	1.0	10.0	0.1	-8.4	1.4	-14.9	0.7
Clark et al. (1992)	Field	Calcite	28	6	11.5	-7.5	1.1	-25.4 ^c	0.2	-17.0	1.1	-0.5	0.1	41.7^b	0.5	14.3 ^c	0.9	-6.5	1.4	-13.0	0.4
Clark et al. (1992)	Field	Calcite	28	6	11.5	-7.5	1.1	-25.3 ^d	0.7	-16.9	1.3	-0.9	0.1	41.7^b	0.5	13.5 ^d	0.1	-7.2	1.0	-13.6	0.4
Mervine et al. (2014)	Field	Calcite	28	6	11.0	-8.6 ^b	0.2	-26.5 ^e	0.3	-17.0	0.3	-0.5	1.5	41.7^b	0.5	14.4 ^e	0.4	-6.4	1.1	-12.9	0.8
Falk et al. (2016)	Field	Arag. & calcite	27	5	11.7	-8.6 ^b	0.2	-26.8 ^f	0.6	-17.2	0.6	0.1	1.0	41.7^b	0.5	14.0 ^f	0.4	-6.9	1.1	-13.4	0.7
This study	Field	Arag. & calcite	17	1	11.0	-8.6 ^b	0.4	-26.6 ^g	0.7	-17.1	0.8	-5.7	1.2	41.5^b	1.0	11.4 ^g	0.4	-7.3	1.5	-14.0	1.1

Note: KFFs are expressed as $\epsilon = (\alpha - 1)10^3$.

^a Keeling et al. (2001), station La Jolla (CA, USA).

^b NOAA CO₂ global network station closest to study site (incl. 2017 Ochsenkopf, Germany; 2007–2012 Ketura, Israel; 2007–2012 Kaashidhoo, Maldives; 2013–2018 Trinidad Head, USA).

^c Average modern crust 'NJ' samples.

^d Average modern crust 'clinic' samples.

^e Average of two lowest 'crust' samples.

^f Average of 'WHOI surface film' samples.

^g Average 'BSC floes' samples.

^h CO_{2(aq)}-CO_{2(g)} carbon fractionation of 1.1‰ from Vogel et al. (1970).

ⁱ Corrected for distillation effect.

^j CO₂-H₂O isotopic equilibrium assumed.

^k Calculated with OH⁻-H₂O EFF values from Zeebe (2020).

Acknowledgements

We thank Ellen Olsen for discovering the error and bringing it to our attention.

References

- Bajnai, D., Herwartz, D., 2021. Kinetic oxygen isotope fractionation between water and aqueous OH⁻ during hydroxylation of CO₂. *ACS Earth Space Chem.* 5 (12), 3375–3384.
- Clark, I.D., Fontes, J.C., 1990. Palaeoclimatic reconstruction in northern Oman based on carbonates from hyperalkaline groundwaters. *Quat. Res.* 33, 320–336.
- Clark, I.D., Fontes, J., Fritz, P., 1992. Stable isotope disequilibria in travertine from high pH waters: Laboratory investigations and field observations from Oman. *Geochim. Cosmochim. Acta* 56, 2041–2050.
- Keeling, C. D., Piper, S. C., Bacastow, R. B., Wahlen, M., Whorf, T. P., Heimann, M. and Meijer, H. A. (2001) Exchanges of atmospheric CO₂ and ¹³CO₂ with the terrestrial biosphere and oceans from 1978 to 2000. I. Global aspects, SIO Reference Series, No. 01-06, Scripps Institution of Oceanography, San Diego, pp. 88.
- Srivastava, A., Verkouteren, R.M., 2018. Metrology for stable isotope reference materials: ¹³C/¹²C and ¹⁸O/¹⁶O isotope ratio value assignment of pure carbon dioxide gas samples on the Vienna PeeDee Belemnite-CO₂ scale using dual-inlet mass spectrometry. *Anal. Bioanal. Chem.* 410 (17), 4153–4163.
- Trolier, M., White, J.W.C., Tans, P.P., Masarie, K.A., Gemery, P.A., 1996. Monitoring the isotopic composition of atmospheric CO₂: Measurements from the NOAA Global Air Sampling Network. *J. Geophys. Res.* 101, 25,897–25,916.
- Vogel, J., Grootes, P., Mook, W., 1970. Isotopic fractionation between gaseous and dissolved carbon dioxide. *Zeitschrift für Physik* 230, 225–238.
- Zeebe, R.E., 2020. Oxygen isotope fractionation between water and the aqueous hydroxide ion. *Geochim. Cosmochim. Acta* 289, 182–195.

Article history:
Available online xxx