

# Finding Treatments for COVID-19: A Trial of Antiviral Pharmacodynamics in Early Symptomatic COVID-19 (PLATCOV): Statistical Analysis Plan

September 21, 2022

Registered at clinicaltrials.gov number **NCT05041907**

**Version 2.2**; refers to Master Protocol version V0.1

**Written by:**

Dr James Watson (Statistician)

Signature:




Date: **21/09/2022**

**Reviewed by and approved by:**

Prof Sir Nicholas White (Co-PI)

Signature:



Date: **21/09/2022**

Dr William Schilling (Co-PI)

Signature:

Date:

# Contents

<b>1</b>	<b>Summary of changes since version 1</b>	<b>3</b>
<b>2</b>	<b>Trial Overview</b>	<b>3</b>
2.1	Main research questions . . . . .	3
<b>3</b>	<b>Study Methods</b>	<b>4</b>
3.1	Trial design . . . . .	4
3.2	Randomisation . . . . .	4
3.3	Sample size . . . . .	5
3.4	Framework . . . . .	5
<b>4</b>	<b>Statistical interim analyses and stopping rules</b>	<b>5</b>
4.1	Interim analysis decision making . . . . .	6
4.2	Timing of final analysis . . . . .	6
4.3	Timing of outcome assessments . . . . .	7
<b>5</b>	<b>Statistical Principles</b>	<b>7</b>
5.1	Posterior estimates . . . . .	7
5.2	Adherence and protocol deviations . . . . .	7
5.3	Analysis populations . . . . .	7
<b>6</b>	<b>Trial population</b>	<b>7</b>
6.1	Screening data . . . . .	8
6.2	Eligibility . . . . .	8
6.3	Withdrawal and follow-up . . . . .	8
6.4	Baseline patient characteristics . . . . .	9
<b>7</b>	<b>Analysis</b>	<b>9</b>
7.1	Outcome definitions . . . . .	9
7.1.1	Viral load quantification . . . . .	9
7.2	Analysis Methods . . . . .	9
7.2.1	Statistical model for the analysis of the serial viral load data . . . . .	9
7.2.2	Covariate adjustment . . . . .	10
7.2.3	Resolution of fever and symptoms . . . . .	11
7.3	Pharmacokinetic analyses . . . . .	11
7.4	Subgroup analyses . . . . .	11
7.5	Prior distributions . . . . .	12
7.6	Sensitivity analyses . . . . .	12
7.7	Missing data . . . . .	13
7.8	Harms . . . . .	13
7.9	Statistical software and analysis implementation . . . . .	13

# 1 Summary of changes since version 1

Version 2.0: The first interim analysis was triggered on the 2nd of March 2022 when qPCR data from the first 50 patients was received. Following the first interim analysis, the Trial Steering Committee and the Data Safety and Monitoring Board recommended changing the minimum efficacy threshold from a 5% increase in viral clearance to a 12.5% increase. This would result in a lower number of patients required to make a futility decision and a larger number of patients required to make a success decision for treatments with no antiviral effects and antiviral effects similar to those estimated from published data on molnupiravir, respectively.

We also changed the primary analysis from modelling the viral load on the CT scale with batch effects to modelling the viral load on the log copies per mL scale, with standard curve transformation done using control samples from each batch. The interim analysis suggested this was more computationally stable and easier to interpret.

Version 2.1: We have added a section regarding the primary analysis of the pharmacokinetic data gathered from all patients.

Version 2.2: We have added a section regarding the analysis of the time to resolution of fever, time to resolution of symptoms, and proportion with normal lymphocyte counts. We updated the current status of the trial (interventions being randomised and the current positive control).

## 2 Trial Overview

An effective, well tolerated, safe, affordable and generally available treatment of COVID-19 that prevented progression to severe disease would be of enormous global health benefit. There are many potential antiviral therapeutics for COVID-19, mainly consisting of repurposed small molecule drugs. At time of writing, however, there is strong evidence for clinical benefit only for the monoclonal antibodies (notably the monoclonal casirivimab/imdevimab in pre-Omicron SARS-CoV-2 variants [1]) and the small molecule drugs remdesivir [2], molnupiravir [3], and nirmatrelvir [4]. The clinical benefits of these interventions were determined from large phase 3 studies in high-risk individuals enrolled shortly after the onset of a symptomatic SARS-CoV-2 infection. The primary endpoint was most commonly either hospitalisation or death from COVID-19 (usually less than a 10% event rate). As vaccine availability has become widespread in addition to high levels of natural immunity from infection, conducting these types of clinical trials is more and more difficult as the event rates for the primary endpoints become increasingly low. This calls for improved study designs which allow for the assessment of antiviral interventions in an efficient manner (hundreds rather than thousands of patients). Currently there is no consensus pharmacometric methodology to determine which antivirals should be prioritised for large phase 3 and 4 evaluation, and how to compare current candidates [5].

PLATCOV is a multi-country platform adaptive trial which will develop and validate a methodology for the quantitative assessment of antiviral effects in low-risk patients with high viral burdens and uncomplicated COVID-19. We choose patients with a low risk of progression as this justifies a no treatment arm; we target patients with high viral loads as this is a subgroup in whom antiviral effects can be detected more easily.

This statistical analysis plan (SAP) covers analyses for both the interim and the final reports. We developed the SAP following the Guidelines for the Content of Statistical Analysis Plans in Clinical Trials [6]. It includes pre-specified decision rules for continuing or stopping individual trial arms based on effectiveness or futility.

### 2.1 Main research questions

Each clinical site in this platform trial will test multiple interventions simultaneously (depending on local regulations and drug availability). The control arm consists of no intervention other than antipyretics and will be compared to intervention arms of three distinct types:

- Newly available or repurposed antiviral drugs and monoclonal antibodies (currently randomizing patients to favipiravir, nitazoxanide, fluoxetine, tixagevimab/cilgavimab, molnupiravir)
- Positive control (currently nirmatrelvir/ritonavir)
- Novel small molecule drugs (none currently)

There are distinct primary objectives for each intervention type. For the newly available antivirals and repurposed drugs, we want to characterise their antiviral activity by comparing viral clearance dynamics with no treatment. Many of these drugs are already used and recommended in some countries. Showing that they do not have clinically significant antiviral activity is as important as showing that they do. For interventions that do have a clear measurable antiviral activity, we will then compare their activity to the current gold-standard antiviral treatment (since August 2022 this is nirmatrelvir/ritonavir).

For monoclonal antibodies included on the platform, we are primarily interested in tracking their performance over time with respect to new SARS-CoV-2 variants (comparison with the no treatment group).

For any new small molecule drugs in development that pass phase 1 testing, we want to rapidly demonstrate antiviral activity (they will have a stronger rationale, i.e. a priori are more likely to be effective) compared to available drugs. In this case, showing superiority or non-inferiority compared to a gold-standard becomes the key objective.

The secondary objectives of this study are:

- To characterise the determinants of viral clearance in early symptomatic SARS-CoV-2 (e.g. estimate the contribution of age, baseline serology, virus genotype, and prior vaccination);
- To determine optimal dosing regimens for drugs shown to have considerable antiviral activity (for interventions of type A and C that have  $>0.9$  probability of increasing viral clearance rate  $>12.5\%$ : we will conduct a pharmacokinetic-pharmacodynamic sub-study);
- To compare the antiviral activity of active interventions against current standard of care monoclonal antibodies.
- To compare fever clearance time and time to symptom resolution with respect to no treatment for interventions that are shown to have a measurable antiviral effect.

The tertiary objective of this study is to characterise the relationship between viral clearance and the risk of subsequent hospitalisation or death by day 28. However, the event rate in the enrolled population is likely to be extremely low (we hope 0%) so it is very unlikely that we will be able to demonstrate any link between viral clearance and progression to severe COVID-19.

## 3 Study Methods

### 3.1 Trial design

This is a multi-centre, multi-country, open label, randomised, controlled, adaptive platform trial of antiviral interventions in early symptomatic SARS-CoV-2. The control arm consists of no study drug other than antipyretics. Interventions currently included in the platform are: favipiravir, nitazoxanide, fluoxetine, tixagevimab/cilgavimab, molnupiravir, and nirmatrelvir/ritonavir. We stopped randomisation to ivermectin (futility decision) in April 2022 [7], and remdesivir (success decision) in June 2022. Casirivimab/imdevimab will be stopped when the all drug supplies are used up. Nirmatrelvir became the positive control in August 2022 when it met the success stopping rule. At each site there is equal allocation to all interventions but with a minimum of 20% of patients randomised to the negative control arm.

### 3.2 Randomisation

Randomisation is performed via a centralised web-app designed by MORU software engineers using RShiny, hosted on a MORU server. Each study nurse responsible for randomising patients has unique login credentials provided by email from the main study statistician (James Watson). Randomisation sheets are pre-generated for each study site separately using blocks of size  $K$  (where  $K$  is equal to 3 times the number of interventions available at the site) with an additional  $100/K\%$  fuzziness (randomly interchanging one of the allocations per block to avoid the study nurse knowing exactly what the  $K$ th allocation will be).

Randomisation sheets and randomisation event logs are stored on a secure Dropbox folder (professional version that has full version control) which is accessed directly by the RShiny app. Only the study statistician James Watson and the MORU IT manager have read/write privileges

to this Dropbox folder. For cross checking purposes, the randomisation app also records the patient age and sex. Each randomisation event is logged with the corresponding time-stamp and the identity of the nurse who performed the randomisation. Each time the set of interventions available at a given study site changes, a new randomisation sheet is generated, overwriting the previous one.

### 3.3 Sample size

The sample size is adaptive (there is no fixed sample size). The required sample size depends on how stringent the thresholds are that define futility or success. We used a simulation approach to determine futility and success thresholds that result in control of the type 1 error at approximately 10% and control of the type 2 error at approximately 20% (see study protocol for exact details). The simulations assumed that the decline in viral loads was log-linear (linear on the cycle threshold (CT) scale), using measurement error estimated from an open access database of prospectively followed individuals with frequent viral load measurement [8, 5]. We calibrated plausible effect sizes using preliminary data from the casirivimab/imdevimab phase 2 studies (this suggested increases of approximately 20% in the slope of the viral clearance on the CT scale) [9]. For these futility and success thresholds, the simulations suggested that the number of patients randomised per arm in order to reach a decision is:

- Approximately 50 patients (average; median is 30) for each efficacious intervention (assuming an effect size of 10% increase in viral clearance slope);
- Approximately 40 patients (average; median is 25) for each inactive intervention (assuming an effect size of 0%).

Therefore, supposing there are 5 interventions of type A in the platform, and assuming that only 1 of the 5 are, in fact, effective, we would need on average a total of  $50 + 4 \times 40 = 210$  patients randomised to interventions of type A (approximately 66% of the total sample size: a minimum of 20% of samples are randomised to the negative control and thus 13% of patients are randomised to each intervention arm and the positive control arm). This implies a total of 320 patients for the first set of interventions identified (this does not include subsequent patients randomised to the intensive PK sub-studies).

### 3.4 Framework

All primary comparisons will be made with respect to the negative control arm (no “across intervention” arm comparisons in the primary analysis). In the case of a site not randomising patients to the negative control (because of local objection), that site will only provide indirect data to support the treatment estimates (under the hierarchical model structure). All decisions concerning efficacy will be based on super-superiority relative to the negative control arm whereby the probability of an effect is defined under the model as the posterior probability that the increase in viral clearance relative to the control arm is greater than 12.5%.

For interventions which show a measurable antiviral effect (i.e. meet the stopping rule for success), we will then compare their antiviral effect to the positive control arm (currently nirmatrelvir/ritonavir).

All primary analyses and stopping decisions will be made using **contemporaneous controls only**. As this is an adaptive trial, temporal drift (due to mutations in the virus, changes in the patient characteristics etc) can confound comparisons with non-contemporaneous controls. However, we will also fit models to the full data set (using all controls), with additional covariate factors for the different trial epochs.

## 4 Statistical interim analyses and stopping rules

We plan frequent interim analyses (Figure 1). This is to allow for near-real time monitoring of accrued data in order to detect possible issues in patient recruitment, viral load swabbing, or with the PCR assays at the different sites. We allow for a slightly inflated type 1 error (10%) as this is a phase 2 study (and so the type 1 vs type 2 error trade-off is different from typical phase 3 studies). In addition, meeting the success threshold will trigger a subsequent intensive




Figure 1: Planned interim analyses and decision rules for stopping an arm or triggering an intensive PK-PD sub-study.

PK-PD sub-study to determine dose-response effects thus gathering additional efficacy data for that intervention. If there is strong *a priori* evidence of antiviral efficacy (e.g. molnupiravir, nirmatrelvir) additional dense blood samples for population PK-PD modelling will be obtained from enrolled patients willing to have daily blood sampling (all patients in the study have blood samples taken at baseline, day 3 and day 7 trough levels, and day 14).

For the type A interventions, we are interested in making decisions regarding futility (highly unlikely that the viral clearance is increased by  $> 12.5\%$ ) and regarding success (highly likely that the viral clearance is increased by  $> 12.5\%$ ). For type B interventions (positive control) we are interested in estimating their antiviral effect (in order to interpret and calibrate the observed effect sizes for the type A/C interventions). For monoclonal antibodies we are interested in detecting temporal changes related to virus genotype (variant escape).

#### 4.1 Interim analysis decision making

Only the study statistician (James Watson) will be unblinded to all interim analyses. Each interim analysis report will sent to the Data Safety and Monitoring Board (DSMB) along with a summary of whether the futility or success thresholds for any of the interventions have been met. The DSMB will then make recommendations for continuing or stopping the intervention arms. Concurrent data from other trials (e.g. safety data) may be used in making these decisions. If an arm is stopped for futility or an intensive PK-PD study is triggered, then all the data pertaining to the negative control arm and the intervention arm will be presented to the Trial Steering Committee (TSC), along with the DSMB recommendations. The interim analysis of the first 50 patients with available qPCR data will be presented unblinded to the TSC in order to determine any necessary final changes to the statistical analysis plan.

#### 4.2 Timing of final analysis

As this is a platform trial, no overall final analysis is planned. For each intervention studied, we define the “final analysis” as:

- For an intervention of type A or C that meets the futility criteria ( $< 0.1$  probability that viral clearance is increased by at least  $12.5\%$ ): final analysis will be timed from when all patients randomised to that arm have passed their day 28 follow-up.

- For an intervention of type A or C that meets the success criteria ( $>0.9$  probability that viral clearance is increased by at least 12.5%): final analysis will be timed from when the last patient is enrolled in the intensive PK-PD sub-study and has passed their day 28 follow-up.
- For an intervention of type B: if a monoclonal antibody then the decision to drop the arm is made based on loss of efficacy due to variant escape; if a small molecule drug final analysis will be timed from when the last patient randomized to that arm passes their day 28 follow-up.

### 4.3 Timing of outcome assessments

For the primary and secondary endpoints we will use all viral load measurements taken up until day 7. Time will be defined as time since randomisation (units of days, including all timepoints  $< 8$  days since randomisation as there will be some variation in exact clock times of the follow-up swabs).

For fever and symptom resolution, time to resolution will be calculated from daily measurements taken over the first week with right censoring for unobserved events (withdrawal or no resolution by 7 days).

The tertiary endpoint (all cause hospitalisation) will be taken up until day 28.

## 5 Statistical Principles

### 5.1 Posterior estimates

Treatment effects will be estimated under a Bayesian framework using a hierarchical model with weakly informative prior distributions. For the hyperparameters, the prior distributions are chosen for computational reasons to aid model convergence. For the key population level parameters, we will use priors based on analysis of open access data [5]. The prior on the treatment effect is driven by plausible maximum effect sizes given published data on viral decreases in patients randomised to casirivimab/imdevimab or placebo [1].

### 5.2 Adherence and protocol deviations

This study is open label and thus it is possible that some patients who are randomized to the no study drug arm, or to study interventions perceived to be ineffective, may be given alternative rescue treatment if their symptoms persist and the treating physician is worried that they will deteriorate clinically. This can introduce confounding between the treatment allocation and outcome. Notably, if multiple patients randomized to the negative control were in fact given the positive control before day 7, this could impact the viral loads during follow-up and thus bias the estimates of the rate of viral clearance in the negative control arm. For all protocol deviations regarding treatment (either stopping treatment for safety reasons or switching treatment arms), we will consider all viral load measurements taken after the deviation as censored, i.e. we will estimate viral load clearance rates only using the data during the period of protocol compliance. We will summarise the number of treatment protocol deviations by site.

### 5.3 Analysis populations

The primary analysis will be in the intention-to-treat (ITT) population, including follow-up data only from the period of treatment adherence (modified intention-to-treat, mITT). A minimum of 3 days of follow-up data are necessary in order to be included in the mITT population (e.g. patients who discontinue before day 2 will not be included).

A secondary analysis will be performed in the per-protocol population, defined as patients who had 100% adherence to the allocated intervention arm. Specific reasons for treatment discontinuation will be outlined and recorded in order to assess confounding bias. The safety population will include all patients who have received at least one dose of the intervention.

## 6 Trial population

Figure 2 shows an example CONSORT diagram summarising the number of patients screened; the reasons for exclusion; the number of enrolled patients randomized to each arm; and the number of




Figure 2: Example CONSORT diagram summarising screening, reasons for excluding patients and randomisation.

patients who complete treatment and follow-up and who are included in the modified intention to treat analysis (mITT).

## 6.1 Screening data

We will summarise the reasons for exclusion for the screened patients not eligible for enrolment. This will not be done for the interim analysis, only for the final analysis once an arm is stopped for futility or success.

## 6.2 Eligibility

The key trial eligibility criteria are as follows:

- Previously healthy adults, aged 18 to 50 years (low risk of developing severe COVID-19);
- SARS-CoV-2 positive by lateral flow antigen test OR a positive PCR test for SARS-CoV-2 within the last 24 hours with a CT value of less than 25 (all viral targets);
- Symptoms of COVID-19 (including fever, or history of fever) for less than 4 days (96 hours);
- Oxygen saturation  $\geq 96\%$  measured by pulse-oximetry at time of screening.

## 6.3 Withdrawal and follow-up

Level of withdrawal will be tabulated and cover the following aspects:

- Discontinuation from any of the study interventions
- Withdrawal from study follow-up
- Withdrawal from entire study and requests that data is not used.

Data will be tabulated for the timing of withdrawal from follow up or lost to follow up. The number of withdrawals and reason will be recorded. A participant may withdraw from the intervention, or be withdrawn for the following reason.

- Withdrawal of consent by participant



- Alteration of participants circumstances or condition which gives justification for discontinuation as decided by investigator.

A table will summarise the loss to follow up and withdrawals during the study with the corresponding reasons.

## 6.4 Baseline patient characteristics

The following key baseline characteristics will be summarised, stratified by site:

- Age and sex;
- Baseline viral load (in copies per mL: the mean value of the 4 swabs taken at randomisation);
- Result of SARS-CoV-2 rapid antibody test;
- Baseline quantitative serum antibody titres (when available);
- Vaccination status (vaccine type, number of doses: expressed as not vaccinated, partially vaccinated and fully vaccinated);
- Days since onset of symptoms.

These will be summarised by their mean or median and standard deviation/range for the continuous variables and by their mean for the binary variables.

# 7 Analysis

## 7.1 Outcome definitions

The primary outcome is the inferred rate of viral clearance expressed as a slope coefficient (the mean posterior estimate and the 95% credible interval) for the daily change in the log viral load (on the  $\log_{10}$  copies per mL scale). This will be inferred under a Bayesian mixed effects (hierarchical) model using the  $\log_{10}$  viral load measurements up until day 7. The observed CT values will be transformed to copies per mL using the set of control samples from each batch.

### 7.1.1 Viral load quantification

All viral load quantification will be done using ThermoFisher TaqCheck TM SARS-CoV-2 Fast PCR assay. Each PCR assay is run on a 96 well plate, composed of all the aliquots from the transport media of the swabs from 4 patients (10 timepoints each done in duplicate, i.e. 20 samples per patient) and one control set composed of 6 spiked control samples of known viral densities over the range  $10^7$  to  $10^2$  viral copies per mL, each done in duplicate (12 in total). For each plate (i.e. batch) we can thus estimate a standard curve (which defines the conversion from the CT value to a number of viral copies per mL). At each interim analysis we will perform a quality control check by comparing the estimated standard curves from each plate, both within sites and across sites (each country will run the PCR assays separately). In addition, we will use the standard curve data to estimate heteroscedasticity in the viral load estimation. We expect the PCR measurement error to increase as the viral load decreases. Heteroscedasticity can be shown visually by plotting the true viral load densities in the control duplicated samples against their observed difference in CT value. We can then estimate the variance in the differences as a function of the true viral density.

To help adjust for variation in the human cell content of the swabbed sample the qPCR method assesses viral densities and also human cell densities using RNaseP as a human cell marker.

## 7.2 Analysis Methods

### 7.2.1 Statistical model for the analysis of the serial viral load data

The primary analysis will consist of fitting Bayesian hierarchical (mixed effects) linear models to the  $\log_{10}$  viral load data up until day 7. The treatment effect is defined as the proportional change (expressed as a multiplicative term) in the population slope of the daily change in log viral load. We will model the data on the copies per mL scale, after conversion from CT values using controls

from each 96 well plate. The standard curve transformation will be done by fitting a linear mixed effects (random slope and random intercept for each plate) model to the control data: regressing the CT values on the known log viral densities. This borrows information across plates but allows for batch effects.

The general model likelihood takes the following form:

$$y_{i,t,k} \sim \text{Student}(\lambda, \alpha_0 + \alpha_i + \alpha_k + \alpha_{cov} + \gamma x_{i,t} + \beta_0 e^{\beta_k + \beta_i + \beta_{cov} + \beta_{T(i)} t}, \sigma^2)$$

where:

- $y_{i,t,k}$  is the log viral load ( $\log_{10}$  copies per mL) for patient  $i$  at time  $t$  (in days since randomisation) from site  $k$ .
- $T(i)$  is the randomized treatment allocation for individual  $i$ .
- $\sigma^2$  is the variance of the measurement error.
- $\lambda$  is the degrees of freedom of the student-t error model.
- The slope of viral clearance decomposes into 5 multiplicative terms: the population mean slope  $\beta_0$ ; the site random effect  $\beta_k$  ( $e^{\beta_k}$  is thus the proportional change in slope); the individual random effect  $\beta_i$ ; the sum of the covariate effects  $\beta_{cov}$ ; and the treatment effect (fixed across sites and individuals)  $\beta_{T(i)}$ . All comparisons are made relative to the no antiviral treatment control arm, so we set  $\beta_{control} = 0$ .
- The intercept term (baseline log viral load) decomposes into 4 terms: the population intercept  $\alpha_0$ ; the site random effect  $\alpha_k$ ; the individual random effect  $\alpha_i$ ; and the sum of any covariate effects  $\alpha_{cov}$ .
- $x_{i,t}$  is the relative human RNaseP quantification (RNaseP  $\Delta$ CT value: this is proportional to the log number of human cells) for patient  $i$  at time  $t$ . The parameter  $\gamma$  thus provides an adjustment for human cell content in each swab. We scale  $x_{i,t}$  so that it has mean 0.

The analysis will sequentially fit four nested models. The base model (model 0: ‘vanilla’) only includes individual and site random effect terms for the slope and intercept. Model 1 adds the parameter  $\gamma$  for the adjustment for human RNaseP. Model 2 then incorporates the covariate adjustment terms  $\alpha_{cov}$  and  $\beta_{cov}$  (see below for their definition). If analysis only has data from one site, then the site random effect term  $\alpha_k$  will be dropped. The final model (model 3) is a non-linear version of model 2 (see below). Efficacy estimation is made based on model 2 (log-linear model with covariate adjustment).

We have chosen a student- $t$  distribution for the model likelihood (with the number of degrees of freedom  $\lambda$  estimated from the data) as this is robust against departures from normal (Gaussian) error, and against model mis-specification, notably concerning the assumption of log-linear decline in viral loads. Prior analyses carried out in order to prepare this statistical analysis plan suggest that a considerable number (e.g. up to 5%) of viral load densities can depart substantially from the expected distribution under a simple log-linear model with Gaussian error [5]. This can partially be explained by patterns of bi-exponential decay in viral loads. It is unclear if the slope of the second phase of elimination in a bi-exponential decline is affected by interventions. We therefore choose not to fit bi-exponential models as this makes the interpretation of the treatment effect more difficult and prior modelling suggested no major impact in terms of treatment effect estimation [5].

Goodness of fits will be assessed using leave-one-out validation [10] and Bayesian  $R^2$  approximation [11].

### 7.2.2 Covariate adjustment

In addition to the human RNaseP adjustment, the primary analyses will adjust for the following covariates (through the parameters  $\alpha_{cov}, \beta_{cov}$ ):

- The serological status on admission as a binary variable corresponding to the result of the screening rapid antibody test (+/-);
- Age (in years - scaled to have mean zero);

- A quantitative measure of serological status using a pre-specified antibody (not yet decided). These data will only be available after August 2022.
- Previous SARS-CoV-2 vaccination (as an ordinal variable: 0: no doses; 1: partially vaccinated with only one dose; 2: fully vaccinated with 2 doses; 3: 3 or more doses);
- The SARS-CoV-2 variant of concern (following WHO labels, with the reference strain chosen to be the major Delta lineage which was the dominant variant of concern when the trial started in Bangkok in September 2021; for sub-lineages, e.g. BA.2, data analysis will be exploratory and not part of the primary analysis of treatment effects).
- The trial “epoch”: this is defined as the discrete intervals of time at which new interventions enter or exit the study. This is to adjust for temporal effects which could confound across arm comparisons made for non-contemporaneously recruiting arms.

If any of the binary covariates show no variation (e.g. all patients are vaccinated) then we will drop the term from the model. We will fit both the covariate adjusted model and the non-adjusted model in all instances as to check for sensitivity to the covariate model. A sensitivity analysis regarding temporal drift will be done by analysing each arm individually along with contemporaneously enrolled controls.

### 7.2.3 Resolution of fever and symptoms

Time to resolution of fever and time to resolution of symptoms will be analysed by comparing Kaplan-Meier curves (right censored data). Resolution of fever is defined as an axillary temperature since  $\leq 37.0^\circ\text{C}$  (for both measurements taken over one day). Resolution of symptoms is defined as no symptoms. An additional analysis will look at the proportion of patients with lymphocyte count within the normal ranges at D3 and D7. Differences in time to resolution for both endpoints will be tested using the log-rank test. Differences will be described by the change in the median time to resolution.

## 7.3 Pharmacokinetic analyses

All patients will have 4 plasma samples taken at: randomisation, day 3, day 7 (all three are pre-dose), and finally day 14. Drug concentrations will be quantitated from these samples in order to explore dose-response relationships between drug (or metabolite) exposure and outcome (slope of viral clearance). There are two objectives for the pharmacokinetic data:

- Firstly, to verify that plasma concentrations are within the expected range (using published pharmacokinetic data from equivalent patient groups and dosing);
- Secondly, to explore whether there is any correlation between plasma concentrations and outcome (rate of viral clearance).

The first objective will use published pharmacokinetic data to assess whether drug levels are within the expected range. For the second objective, the primary analysis will be performed using the day 3 level (a trough level). We will fit a linear model between the day 3 level (log transformed) and the mean viral clearance slope estimate from the main analysis in patients allocated to the intervention arm. Significance is defined as a two-sided test at the 5% level.

## 7.4 Subgroup analyses

The primary pre-specified subgroup analysis will look at the relationship between the efficacy of the monoclonal antibodies (currently the Regeneron antibody cocktail) and the virus genotype expressed as the major WHO lineages. This will be done by fitting an interaction term  $\delta x_{var}(i)e^{\beta_{T(i)}}$  for the relevant  $T(i)$ ; where  $x_{var}(i)$  is the virus variant infecting individual  $i$ .

Secondary subgroup analyses will be carried out only for treatment arms that show an antiviral effect (meet the success criteria). We will look at subgroup effects for (listed in order of priority):

- SARS-CoV-2 variant of concern;
- Serological status on admission (using quantitative serum antibody);

- Vaccination status;
- Age and sex;

## 7.5 Prior distributions

In all the following, for the normal distributions, the second term corresponds to the scale (standard deviation not the variance). The viral load values are on the  $\log_{10}$  scale (i.e. for the intercept term 6 would correspond to 1,000,000 copies per ml).

### Population level parameters

$$\begin{aligned}\alpha_0 &\sim \text{Normal}(6, 2) \quad (\text{population intercept}) \\ \beta_0 &\sim \text{Normal}(-0.5, 1) \quad (\text{population slope}) \\ \sigma &\sim \text{Normal}(1.5, 3) \quad (\text{standard deviation of measurement error}) \\ \gamma &\sim \text{Normal}(0, 1) \quad (\text{human RNaseP adjustment}) \\ \beta_{T(i)} &\sim \text{Normal}(0, 0.5) \quad \text{log treatment effect (subgroups also)}\end{aligned}$$

### Covariate effects

$$\begin{aligned}\beta_{cov} &\sim \text{Normal}(0, 1) \quad (\text{covariate effect on slope}) \\ \alpha_{cov} &\sim \text{Normal}(0, 1) \quad (\text{covariate effect on intercept})\end{aligned}$$

### Random effects and hyperparameters

$$\begin{aligned}\Omega &\sim \text{Cholesky}(2) \quad (\text{correlation matrix for individual random effects}) \\ \Sigma &\sim \text{Exponential}(1) \quad (\text{standard deviation for individual random effects}) \\ \lambda &\sim \text{Exponential}(1) \quad (\text{t-distribution degrees of freedom})\end{aligned}$$

## 7.6 Sensitivity analyses

Sensitivity to the prior specification will be assessed by multiplying all prior standard deviation terms by 5. Sensitivity to the assumption of log-linear decline will be assessed by fitting an ‘up-down’ model of viral dynamics [12] (where “up” refers to viral multiplication and “down” the subsequent clearance). Under this model the likelihood is (dropping batch, site, and covariate terms for notational simplicity, these are added analogous to the previous log-linear model) given by:

$$y_{i,t} \sim \text{Student} \left( \lambda, \alpha_0 + \alpha_i + \log \left\{ \frac{\beta_1^i + \beta_2^i}{\beta_2^i e^{-\beta_1^i [t - t_{\max}^i]} + \beta_1^i e^{\beta_2^i [t - t_{\max}^i]}} \right\}, \sigma^2 \right)$$

Under this model, the term  $\alpha_0 + \alpha_i$  is the individual peak log viral load which occurs at time  $t_{\max}^i$  (this is the time relative to randomisation, if negative it is therefore unobserved). The parameters  $\beta_1^i, \beta_2^i$  are the growth and clearance rates for individual  $i$ , respectively.

If some patients are enrolled very early in the course of their illness, it is possible that their viral loads increase over the first days of follow-up. This would mean that the clearance rate estimated under a simple log-linear decline model would be biased towards a shallower slope. Randomisation protects against confounding bias as the sample size increases, however for small sample sizes unequal allocation of ‘pre-peak’ versus ‘post-peak’ individuals could influence treatment effect estimates. Treatment effects are inferred in the same way as for the log-linear model (proportional change in the population slope coefficients), by acting on  $\beta_2$  (note that in reality, the treatment effect most likely would change all three parameters  $\alpha_0, \beta_1, \beta_2$ , however this would mean inferring three separate treatment effects as the model is non-mechanistic).

## 7.7 Missing data

Missing viral load data will not be imputed. The model will be fit to all available viral load densities up until day 7 or day of treatment deviation. For all analyses, missing data on the key baseline variables (vaccination, virus genotype, antibody rapid test) will be imputed at random using the observed site-specific covariate distribution multiple times (5 times, the model will be refit). We do not expect much missing data (eg <1%). If delays in genotyping result in missing variant data, we will impute the variant data using genetic surveillance data.

## 7.8 Harms

Safety analyses will include all patients who have received at least one dose of the intervention. The safety and tolerability data will be pooled from all the sites that receive the same study intervention. The safety and tolerability will be assessed by comparing the frequency of adverse events and serious adverse events when compared to the control intervention.

All adverse event summaries will refer to treatment emergent adverse events, i.e. adverse events that newly started or increased in intensity after study drug administration (or from hour 0 in the control group receiving no intervention). Adverse events will be graded according to CTCAE V5.0. Adverse Events (AEs) of grade 3 and above will be recorded, the grading is as follows - 1 = mild, 2 = moderate, 3 = severe, 4 = life-threatening, 5 = fatal. All Serious Adverse Events (SAEs) will be recorded, a serious adverse event was any untoward medical occurrence that resulted in death, was life threatening, required inpatient hospitalization or prolongation of existing hospitalization, resulted in persistent or significant disability/incapacity or consisted of a congenital anomaly or birth defect.

We will tabulate AE of grade 3 and higher, SAEs, and deaths, by treatment arm with comparisons made relative to the no study drug arm.

None of the drugs evaluated will be pre-registration or pre-approval, so adverse effect profiles for each are already well known.

## 7.9 Statistical software and analysis implementation

All analyses will be done in R. Interim analyses and the final analyses will be done using RMarkdown scripts, pre-coded to ensure full transparency. All code for the statistical analysis is available on the following github repository: <https://github.com/jwatowatson/PLATCOV-SAP>. This ensures full version control by tracking all changes made to the analysis code.

The statistical models for the analysis of the serial viral load data are written in *stan* and fitted via the *rstan* interface [13]. These are available on the github repository. For each model, we will run 6 parallel chains for 10,000 iterations, discarding half for burn-in and thinning every 40 (thus 1,000 posterior draws).

Convergence of Bayesian fits will be assessed visually by examining the traceplots of the key model parameters and assessing the Rhat values for each parameter (Gelman-Rubin statistic).

## References

- [1] Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGEN-COV Antibody Combination and Outcomes in Outpatients with Covid-19. *New England Journal of Medicine*. 2021;385(23):e81. doi:10.1056/NEJMoa2108163.
- [2] Gottlieb RL, Vaca CE, Paredes R, Mera J, Webb BJ, Perez G, et al. Early Remdesivir to Prevent Progression to Severe Covid-19 in Outpatients. *New England Journal of Medicine*. 2022;386(4):305–315. doi:10.1056/NEJMoa2116846.
- [3] Jayk Bernal A, Gomes da Silva MM, Musungaie DB, Kovalchuk E, Gonzalez A, Delos Reyes V, et al. Molnupiravir for oral treatment of Covid-19 in nonhospitalized patients. *New England Journal of Medicine*. 2021;.
- [4] Hammond J, Leister-Tebbe H, Gardner A, Abreu P, Bao W, Wisemandle W, et al. Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19. *New England Journal of Medicine*. 0;0(0):null. doi:10.1056/NEJMoa2118542.

- [5] Watson JA, Kissler SM, Day NPJ, Grad YH, White NJ. Characterizing SARS-CoV-2 Viral Clearance Kinetics to Improve the Design of Antiviral Pharmacometric Studies. *Antimicrobial Agents and Chemotherapy*. 2022;66(7):e00192–22. doi:10.1128/aac.00192-22.
- [6] Gamble C, Krishan A, Stocken D, Lewis S, Juszczak E, Doré C, et al. Guidelines for the Content of Statistical Analysis Plans in Clinical Trials. *JAMA*. 2017;318(23):2337–2343. doi:10.1001/jama.2017.18556.
- [7] Schilling WH, Jittamala P, Watson JA, Ekkapongpisit M, Siripoon T, Ngamprasertchai T, et al. Pharmacometric assessment of the in vivo antiviral activity of ivermectin in early symptomatic COVID-19. *medRxiv*. 2022;doi:10.1101/2022.07.15.22277570.
- [8] Kissler SM, Fauver JR, Mack C, Olesen SW, Tai C, Shiue KY, et al. Viral dynamics of acute SARS-CoV-2 infection and applications to diagnostic and public health strategies. *PLoS biology*. 2021;19(7):e3001333.
- [9] Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with COVID-19. *New England Journal of Medicine*. 2020;doi:10.1056/NEJMoa2035002.
- [10] Vehtari A, Gelman A, Gabry J. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and computing*. 2017;27(5):1413–1432.
- [11] Gelman A, Goodrich B, Gabry J, Vehtari A. R-squared for Bayesian Regression Models. *The American Statistician*. 2019;73(3):307–309. doi:10.1080/00031305.2018.1549100.
- [12] Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, et al. Community transmission and viral load kinetics of the SARS-CoV-2 delta (B. 1.617. 2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *The Lancet Infectious Diseases*. 2021;.
- [13] Stan Development Team. RStan: the R interface to Stan; 2020. Available from: <http://mc-stan.org/>.