

# BirdNET-Pi Weekly Bird Report

BirdWeather / PUC

```
knitr::opts_chunk$set(echo = FALSE, message = FALSE, warning = FALSE)
```

```
library(readr)
library(dplyr)
```

```
##
## Attaching package: 'dplyr'
```

```
## The following objects are masked from 'package:stats':
##
##   filter, lag
```

```
## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union
```

```
library(lubridate)
library(ggplot2)
library(forcats)
```

```
tz_local    <- "Europe/Amsterdam"    # adjust if needed
```

```
no_data <- FALSE
reason  <- NULL
```

```
# --- Robust path lookup for master_detections.csv ---
```

```
# We try both from repo root and from scripts/ (one level below)
```

```
candidate_paths <- c(
  "data/master_detections.csv",
  "../data/master_detections.csv"
)
```

```
path_master <- NULL
for (p in candidate_paths) {
  if (file.exists(p) && file.info(p)$size > 0) {
    path_master <- p
  }
}
```

```

    break
  }
}

if (is.null(path_master)) {
  no_data <- TRUE
  reason <- paste(
    "Master file missing or empty. Tried paths:",
    paste(candidate_paths, collapse = ", ")
  )
} else {
  df_raw <- read_csv(path_master, show_col_types = FALSE)

  # auto-detect species + time columns
  if (!all(c("timestamp", "species.commonName") %in% names(df_raw))) {
    no_data <- TRUE
    reason <- paste(
      "Expected columns 'timestamp' and 'species.commonName' not found. Columns present:",
      paste(names(df_raw), collapse = ", ")
    )
  } else {
    df <- df_raw %>%
      mutate(
        species = .data[["species.commonName"]],
        ts_utc = ymd_hms(.data[["timestamp"]], quiet = TRUE),
        ts_local = with_tz(ts_utc, tz_local),
        date = as_date(ts_local),
        hour = hour(ts_local)
      ) %>%
      filter(!is.na(species), !is.na(ts_local))

    if (nrow(df) == 0) {
      no_data <- TRUE
      reason <- "No valid detections after parsing timestamps and species."
    } else {
      # define 7-day window based on latest date in data
      last_date <- max(df$date, na.rm = TRUE)
      start_date <- last_date - days(6)
      end_date <- last_date

      df7 <- df %>% filter(date >= start_date & date <= end_date)

      if (nrow(df7) == 0) {
        no_data <- TRUE
        reason <- paste(
          "No detections in the last 7 days of data (",
          as.character(start_date), "-", as.character(end_date), ")."
        )
      }
    }
  }
}

```

```

)
} else {
  # global weekly metrics
  total_detections <- nrow(df7)
  unique_species <- n_distinct(df7$species)

  most_common <- df7 %>%
    count(species, sort = TRUE) %>%
    slice(1)

  # confidence if present
  conf_col <- intersect(c("confidence", "score", "probability"), names(df7))[1]
  avg_conf <- if (!is.na(conf_col)) {
    mean(df7[[conf_col]], na.rm = TRUE)
  } else NA_real_

  # top 10 species
  top10 <- df7 %>%
    count(species, sort = TRUE) %>%
    slice_head(n = 10)

  # BirdNET-style palette
  birdnet_colors <- c(
    "#4e79a7", "#59a14f", "#f28e2c", "#e15759", "#76b7b2",
    "#edc949", "#af7aa1", "#ff9da7", "#9c755f", "#bab0ab"
  )

  # order species for plotting
  top10 <- top10 %>%
    arrange(n) %>%
    mutate(species = factor(species, levels = species))

  # time-of-day categories
  df7_periods <- df7 %>%
    mutate(period = case_when(
      hour >= 0 & hour <= 4 ~ "Night (0-4)",
      hour >= 5 & hour <= 8 ~ "Dawn (5-8)",
      hour >= 9 & hour <= 11 ~ "Morning (9-11)",
      hour >= 12 & hour <= 16 ~ "Afternoon (12-16)",
      hour >= 17 & hour <= 19 ~ "Evening (17-19)",
      hour >= 20 & hour <= 23 ~ "Night (20-23)"
    ))

  tod <- df7_periods %>%
    count(period) %>%
    mutate(period = factor(period, levels = c(
      "Night (0-4)", "Dawn (5-8)", "Morning (9-11)",

```

```

      "Afternoon (12-16)", "Evening (17-19)", "Night (20-23)"
    )))

  # daily detection trends
  daily_trend <- df7 %>%
    count(date) %>%
    arrange(date)

  # new species in this 7-day window
  first_all <- df %>%
    group_by(species) %>%
    summarise(first_date = min(date), .groups = "drop")

  new_this_week <- first_all %>%
    filter(first_date >= start_date & first_date <= end_date) %>%
    left_join(
      df7 %>% count(species, name = "detections_this_week"),
      by = "species"
    ) %>%
    arrange(first_date)
}
}
}
}
if (no_data) {
  cat("No data available for this weekly report.\n\nReason:\n", reason)
} else {
  summary_tbl <- tibble::tibble(
    Metric = c(
      "Total detections",
      "Unique species",
      "Most common species",
      "Detections of most common",
      "Average confidence"
    ),
    Value = c(
      total_detections,
      unique_species,
      most_common$species,
      most_common$n,
      if (!is.na(avg_conf)) sprintf("%.1f%%", avg_conf * 100) else "n/a"
    )
  )

  cat(
    "Period: ",

```


```
as.character(start_date), " - ", as.character(end_date), "\n\n",
sep = ""
)

knitr::kable(summary_tbl, align = c("l","r"))
}
```


## Period: 2026-02-13 - 2026-02-19

Metric	Value
Total detections	3177
Unique species	63
Most common species	Eurasian Blue Tit
Detections of most common	854
Average confidence	82.9%


```
if (!no_data) {
  ggplot(top10, aes(x = species, y = n, fill = species)) +
  geom_col() +
  scale_fill_manual(values = birdnet_colors) +
  coord_flip() +
  labs(
    x = "Species",
    y = "Detections"
  ) +
  theme_minimal(base_size = 11) +
  theme(legend.position = "none")
}
```



```
if (!no_data) {
  ggplot(tod, aes(x = period, y = n, fill = period)) +
  geom_col() +
  scale_fill_manual(values = birdnet_colors) +
  labs(
    x = "Time period",
    y = "Detections"
  ) +
  theme_minimal(base_size = 11) +
  theme(legend.position = "none")
}
```



```
if (!no_data) {  
  ggplot(daily_trend, aes(x = date, y = n)) +  
    geom_line(color = "#4e79a7", linewidth = 1) +  
    geom_point(color = "#4e79a7", size = 2) +  
    labs(  
      x = "Date",  
      y = "Detections per day"  
    ) +  
    theme_minimal(base_size = 11)  
}
```



```

if (no_data) {
  cat("No data available.")
} else {
  if (nrow(new_this_week) == 0) {
    cat("No new species detected in this 7-day period.")
  } else {
    knitr::kable(
      new_this_week %>%
      select(species, first_date, detections_this_week),
      col.names = c("Species", "First detection", "Detections this week"),
      align = c("l", "c", "r")
    )
  }
}

```

Species	First detection	Detections this week
Common Redshank	2026-02-14	1
Egyptian Goose	2026-02-18	1