

# TOPICS IN PERMUTATION GROUPS

## A SHORT SEMINAR SERIES

JOSHUA WISCONS

### CONTENTS

1. A first example	2
2. Permutation groups	4
2.1. The alternating group	4
2.2. One group, many actions	5
2.3. Orbits and stabilizers	8
3. Degrees of symmetry	10
3.1. Affine and projective linear groups	12
3.2. Mathieu groups	15
4. Relational complexity	17
4.1. First examples	18
References	19

These (in progress!) notes are being prepared for a six-part seminar series on permutation groups in Spring 2022 at California State University, Sacramento. The seminar is meant to be casual, example driven, hopefully fun, and accessible to anyone having completed a first course in linear algebra, though exposure to groups will help.

---


*Date:* June 6, 2022.

The author was partially supported by the National Science Foundation under Grant No. [DMS-1954127](#).

## 1. A FIRST EXAMPLE

Let's start by exploring symmetries of a familiar object. We'll also introduce notation as we go.

**Example 1.** Consider a regular tetrahedron, like the [Pyraminx](#) (pictured below). One way to represent the various symmetries is to consider how each symmetry permutes the vertices, so let's label them as follows



Let  $\text{TETRA}$  denote the collection of *all* symmetries of this object and  $\text{TETRA}^+$  denote just the *rotational* symmetries. We will view each element of  $\text{TETRA}$  (and  $\text{TETRA}^+$ ) as a permutation of the set  $\{1, 2, 3, 4\}$ . The collection of *all* permutations of  $\{1, 2, 3, 4\}$  will be denoted  $\text{Sym}(4)$ , so with this notation we have  $\text{TETRA}^+ \subseteq \text{TETRA} \subseteq \text{Sym}(4)$ . Let's pause to define and introduce notation for  $\text{Sym}(X)$  generally.

**Definition.** Let  $X$  be a set. The **symmetric group** on  $X$ , denoted  $\text{Sym}(X)$ , is the set of all permutations of  $X$ . When  $X = \{1, 2, \dots, n\}$ , we write  $\text{Sym}(n)$  in place of  $\text{Sym}(X)$ . We also adopt the following notation.

- The identity permutation will be denoted  $\text{id}_X$  (or just  $\text{id}$ ), so  $\text{id}_X(x) = x$  for all  $x \in X$ .
- For  $\sigma, \tau \in \text{Sym}(X)$ ,  $\sigma\tau$  denotes composition of  $\sigma$  and  $\tau$ , so  $\sigma\tau(x) = \sigma(\tau(x))$  for all  $x \in X$ .
- For  $\sigma \in \text{Sym}(X)$  and  $n \in \mathbb{N}$ ,  $\sigma^n$  denotes the composition of  $\sigma$  with itself  $n$ -times, and  $\sigma^{-n}$  denotes the composition of  $\sigma^{-1}$  with itself  $n$ -times.

We may refer to  $\sigma\tau$  as the *product* of  $\sigma$  and  $\tau$ , but in  $\text{Sym}(X)$  this means composition.

Let's resume our investigation of the tetrahedron, and list some elements of  $\text{TETRA}^+$ . As we do, we'll introduce additional notation for writing permutations.

If we hold the top vertex 1, we can rotate the remaining ones by  $120^\circ$  to obtain the permutation  $\gamma$  defined via  $\gamma(1) = 1$ ,  $\gamma(2) = 3$ ,  $\gamma(3) = 4$ , and  $\gamma(4) = 2$ . We occasionally write permutations in [two-line notation](#) as below; the top row lists the inputs while the bottom lists the corresponding outputs.

$$\gamma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 4 & 2 \end{pmatrix}$$

More often, we will write permutations in so-called (disjoint) [cycle notation](#), which will be "defined" via various examples. Writing  $\gamma$  using cycle notation yields  $\gamma = (234)$ . This is read left to right and indicates that a given number is mapped by  $\gamma$  to the number immediately to the right, cycling back to the beginning when the rightmost parenthesis is encountered. We can picture this as  $2 \rightarrow 3 \rightarrow 4 \rightarrow 2$ . Any number not appearing (like 1

in this case) is left fixed by the permutation. Because of the cyclical nature of this notation, there are various ways to write  $\gamma$  in this notation:  $\gamma = (234) = (342) = (423)$ .

Let's look for more symmetries. If we continue to hold the top vertex and rotate  $120^\circ$  again (for a total of  $240^\circ$ ), we have the permutation  $\gamma^2 = \gamma\gamma$ . In two line notation,

$$\gamma^2 = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 2 & 3 \end{pmatrix},$$

and in cycle notation,  $\gamma^2 = (243)$ . Yet another rotation by  $120^\circ$ , returns the shape to the starting configuration, so  $\gamma^3 = \text{id}$ . Also note, that we could have initially rotated by  $-120^\circ$ , which is the inverse of  $\gamma$  (i.e.  $\gamma^{-1}$ ), but this does not yield a new permutation since  $\gamma^{-1} = (432) = (243) = \gamma^2$ .

Now, this time, let's hold vertex 2, and rotate by  $120^\circ$ . Depending on your point of view, there are two options, but one of them is  $\delta = (134)$ . And rotating in the same direction by another  $120^\circ$ , yields  $\delta^2 = (143)$ . We can further consider holding the vertex 3 or 4 while rotating. So far, we've found the following rotational symmetries of the tetrahedron:

$$\{\text{id}, (234), (243), (134), (143), (124), (142), (123), (132)\} \subseteq \text{TETRA}^+.$$

There are more. Imagine the line  $\ell$  connecting the midpoint of the edge  $\{1, 4\}$  (from vertex 1 to 4) with the midpoint of the edge  $\{2, 3\}$  (from vertex 2 to 3); then let  $\alpha$  be the rotation of the tetrahedron by  $180^\circ$  about the line  $\ell$ . In two-line notation, we find that

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix},$$

and in cycle notation, we have  $\alpha = (14)(23)$ . Performing  $180^\circ$  rotations about the other two lines that connect pairs of "opposite edges", we find that  $\text{TETRA}^+$  also contains  $(12)(34)$  and  $(13)(24)$ . So now we have that

$$\left\{ \begin{array}{l} \text{id}, (12)(34), (13)(24), (14)(23), \\ (234), (243), (134), (143), (124), (142), (123), (132) \end{array} \right\} \subseteq \text{TETRA}^+.$$

What else could be in  $\text{TETRA}^+$ ? Well,  $\text{Sym}(4)$  contains 12 more permutations that we have not encountered yet. Six of the remaining permutations move only two vertices while fixing the others, e.g.  $\tau = (12)$ ; these permutations are called **transpositions** (since they transpose two numbers and leave the remaining unaltered). Now,  $\tau$  does represent a symmetry of the tetrahedron, namely *reflection* through the plane that contains the edge  $\{3, 4\}$  and is orthogonal to the edge  $\{1, 2\}$ . However,  $\tau$  is not a *rotational* symmetry because a nontrivial rotational symmetry can only fix a vertex that is on the axis of rotation, and due to the structure of the tetrahedron, at most one vertex can lie on an axis of rotation that produces a symmetry. Similarly, we find that the other transpositions are in  $\text{TETRA}$  but *not*  $\text{TETRA}^+$ , so

$$\{(12), (13), (14), (23), (24), (34)\} \subseteq \text{TETRA} - \text{TETRA}^+.$$

The only permutations in  $\text{Sym}(4)$  we have not yet considered are the so-called 4-cycles, such as  $\sigma = (1234)$ . Notice that  $\sigma \in \text{TETRA}$  since  $\sigma = (12)(23)(34)$ , so it can be realized by performing three reflections in a row. However,  $\sigma$  is not a rotational symmetry. To see this, note that any rotation that moves vertex 2 to 3 and vertex 3 to 4 must have an axis of rotation orthogonal to *both* of the edges  $\{2, 3\}$  and  $\{3, 4\}$ , implying that the axis of rotation runs through vertex 1, so 1 would have to remain fixed. Thus,  $\sigma$  is not a rotational symmetry.

Similarly, we see that each 4-cycle is in  $\text{TETRA}$  but *not*  $\text{TETRA}^+$ . In conclusion, we've found

$$\begin{aligned}\text{TETRA} &= \text{Sym}(4) \\ \text{TETRA}^+ &= \left\{ \begin{array}{l} \text{id}, (12)(34), (13)(24), (14)(23), \\ (234), (243), (134), (143), (124), (142), (123), (132) \end{array} \right\}\end{aligned}$$

Incidentally, although we already knew that the composition of any two symmetries of the tetrahedron was another symmetry, we can now directly verify that the composition of any two *rotational* symmetries is another *rotational* symmetry (e.g.  $\gamma\delta = (234)(134) = (14)(23) = \alpha$ ).

## 2. PERMUTATION GROUPS

In this section, we'll formally introduce permutation groups and the related notion of group actions. We'll also see more examples (in addition to  $\text{TETRA}$  and  $\text{TETRA}^+$ ); this includes the important family of alternating groups as well as the symmetry group of a cube. The section concludes with a couple of fundamental concepts in the study of permutation groups: orbits and stabilizers.

**Definition.** Let  $X$  be a set. A subset  $G \subseteq \text{Sym}(X)$  is called a **subgroup** of  $\text{Sym}(X)$  if all of the following hold:

- [CLOSURE UNDER THE IDENTITY]  $\text{id} \in G$ ;
- [CLOSURE UNDER COMPOSITION] if  $\alpha, \beta \in G$ , then  $\alpha\beta \in G$ ;
- [CLOSURE UNDER INVERSES] if  $\alpha \in G$ , then  $\alpha^{-1} \in G$ .

In this case, we say  $G$  is a **permutation group** on  $X$ .

**Example 2.** As can be seen from Example 1,  $\text{TETRA}$  and  $\text{TETRA}^+$  can both be viewed as permutation groups on  $\{1, 2, 3, 4\}$ .

**2.1. The alternating group.** Recall that an element  $\tau \in \text{Sym}(X)$  of the form  $\tau = (ij)$  is called a transposition (for any  $i \neq j \in X$ ). Let's explore how to decompose a given permutation into a product (i.e. composition) of transpositions.

**Example 3.** Consider the permutation  $\alpha = (12345)$ . After computing various products of transpositions, one might notice that  $\alpha$  can be written as  $\alpha = (12)(23)(34)(45)$  or  $\alpha = (15)(14)(13)(12)$  or something else. Since  $(ij)(ij) = \text{id}$  for each transposition  $(ij)$ , one sees that there are a lot of ways to write  $\alpha$  as a product of transpositions; for example,  $\alpha = (12)(23)(34)(45)(12)(12)$  or  $\alpha = (67)(12)(23)(34)(45)(67)$  (assuming in the latter case that we are working in  $\text{Sym}(n)$  for  $n \geq 7$ ).

What about the permutation  $\beta = (12345)(6789)$ ? We can decompose  $\beta$  similar to before by treating each of the cycles  $(12345)$  and  $(6789)$  separately:  $\beta = (12)(23)(34)(45)(67)(78)(89)$ . And again, there are many ways to decompose  $\beta$  into a product of transpositions.

Abstracting our technique in the previous example, we get the following result; the proof is left as an exercise. In group-theoretic parlance, it says that each finite symmetric group is *generated* by its transpositions.

**Lemma.** *Every permutation in  $\text{Sym}(n)$  can be written as a product of transpositions.*

**Definition.** Let  $\sigma \in \text{Sym}(n)$ . We call  $\sigma$  an **odd** permutation if it can be written as a product of an odd number of transpositions. Similarly, we call  $\sigma$  an **even** permutation if it can be written as a product of an even number of transpositions.

Looking back to Example 3, the permutation  $\alpha$  was even; while  $\beta$  was odd. Notice that although there were several ways to write  $\alpha$  as a product of transpositions, *every* way we came up with used an even number. It turns out that this was no coincidence.

**Lemma.** *Every permutation in  $\text{Sym}(n)$  is either even or odd, but not both.*

*Proof idea.* One first observation to make is that it suffices to prove  $\text{id}$  is (even but) not odd. Indeed, suppose we know that  $\text{id}$  is not odd. Towards a contradiction, assume an arbitrary  $\sigma$  is both even and odd. Then we can write  $\sigma = \tau_1 \cdots \tau_k = \rho_1 \cdots \rho_\ell$  where  $k$  is even,  $\ell$  is odd, and each  $\tau_i$  and each  $\rho_j$  is a transposition. This implies  $\text{id} = \sigma\sigma^{-1} = \tau_1 \cdots \tau_k \rho_\ell^{-1} \cdots \rho_1^{-1}$ . As the inverse of a transposition is itself,  $\text{id} = \tau_1 \cdots \tau_k \rho_\ell \cdots \rho_1$ , implying that  $\text{id}$  is odd (since  $k$  is even and  $\ell$  is odd), a contradiction.

So, it remains to show  $\text{id}$  is not odd. This is where most of the effort is, and we will simply refer out—sorry! A quick search of the internet will turn up various proofs of this lemma, but one nice (and detailed) outline of the proof can be found in Dana Ernst’s abstract algebra book, see [Ern21, Theorem 4.103].  $\square$

Notice that composing two even permutations results in another even permutation. Moreover, the previous “proof” discussed that  $\text{id}$  is even, and it also indicated that a permutation has the same parity as its inverse. Combining this, we arrive at the following fundamental theorem-definition.

**Theorem.** *Let  $\text{Alt}(n)$  denote the collection of all even permutations in  $\text{Sym}(n)$ . Then  $\text{Alt}(n)$  is a subgroup of  $\text{Sym}(n)$ ; it is called the **alternating group**.*

**Example 4.** A concise summary of Example 1 is that if we view the elements of  $\text{TETRA}$  as permutations of  $\{1, 2, 3, 4\}$ , then  $\text{TETRA} = \text{Sym}(4)$  and  $\text{TETRA}^+ = \text{Alt}(4)$ .

Note that, technically, the elements of  $\text{TETRA}$  operate on the entire tetrahedron, not just the vertices. Thus  $\text{TETRA}$  is not literally equal to  $\text{Sym}(4)$ , so it is more precise to use the language of **isomorphisms** and write  $\text{TETRA} \cong \text{Sym}(4)$  and  $\text{TETRA}^+ \cong \text{Alt}(4)$ . But in general, we won’t much worry about this.

**2.2. One group, many actions.** When analyzing a permutation group  $G$  on some set  $X$ , it is important to keep in mind that  $G$  may very well also permute other sets, and simultaneously studying the various “actions” of  $G$  can be quite advantageous. For example,  $\text{TETRA}$  is first and foremost acting on the points of  $\mathbb{R}^3$  (in a way that preserves the tetrahedron); however, the way we studied  $\text{TETRA}$  was via its action on the *vertices* of the tetrahedron. However,  $\text{TETRA}$  also permutes the *edges* of the tetrahedron as well as the *faces* and various other related sets of objects.

Let’s formalize the notion of an “action”, and then see how to leverage this idea with an example.

**Remark.** In what follows, we will occasionally use a phrase like “let  $G$  be a group.” This can be taken to mean that  $G$  is a permutation group on some set  $X$ , but we don’t care what  $X$  is. The point is that sometimes we only want to focus on how the elements of the group  $G$  interact with each other as opposed to how they permute the elements of  $X$ . If you’ve encountered the definition of an abstract group, then you could alternatively interpret “let  $G$  be a group” as “let  $G$  be an abstract group.”

**Definition.** Let  $G$  be a group. We say  $G$  **acts** on a set  $Y$  if each  $g \in G$  defines a function  $Y \rightarrow Y : y \mapsto g \cdot y$  in such a way that

- (1) for all  $g, h \in G$  and all  $y \in Y$ ,  $g \cdot (h \cdot y) = (gh) \cdot y$ , and
- (2) for all  $y \in Y$ ,  $\text{id} \cdot y = y$ .

We say the action is **faithful** if the only element of  $G$  satisfying  $g \cdot y = y$  for all  $y \in Y$  is  $\text{id}$ .

In reading the definition, one should think of the notation  $g \cdot y$  as  $g(y)$ . In what follows, we will often use the notation  $g \cdot y$  in place of  $g(y)$  for permutation groups too.

**Remark.** If  $G$  acts on a set  $Y$ , then the definition implies that each element of  $G$  can be viewed as a *permutation* of  $Y$ . However, it is possible for different elements of  $G$  to represent the same permutation of  $Y$ . (We'll see this in a later example.)

Now, if the action is faithful, then it can be shown that different elements of  $G$  always act as different permutations of  $Y$ , so in this case, we can view  $G$  as a permutation group on  $Y$  (even though  $G$  may also be a permutation group on some other set  $X$ ). Technical note: the sentence “we can view  $G$  as a permutation group on  $Y$ ” really means that “ $G$  is isomorphic to a permutation group on  $Y$ ”.

**Example 5.** Let's work out the symmetries of a cube like [Rubik's Revenge](#) (pictured below). Let  $\text{CUBE}$  denote the collection of all symmetries of the cube, and let  $\text{CUBE}^+$  denote the rotational symmetries. As in Example 1,  $\text{CUBE}$  acts *faithfully* on the 8 vertices of the cube, so after labeling the vertices (as below), we may view  $\text{CUBE}$  as a subgroup of  $\text{Sym}(8)$ .



Note that  $\text{CUBE}$  also acts faithfully on the 6 faces as well as on the 12 edges, so we could have instead chosen to label the faces or edges to view  $\text{CUBE}$  as a subgroup of  $\text{Sym}(6)$  or  $\text{Sym}(12)$ . However, for now, let's think in terms of the action on the vertices.

Rotating clockwise by  $90^\circ$  around the axis running through the centers of faces  $\{1, 2, 3, 4\}$  and  $\{5, 6, 7, 8\}$ , we get the permutation  $\alpha = (1234)(5678)$ . Performing  $\alpha$  twice (i.e. rotation about the same axis by  $180^\circ$ ) yields the permutation  $\alpha^2 = (13)(24)(57)(68)$ . A third application of  $\alpha$  results in  $\alpha^3 = (1432)(5876)$ , and  $\alpha^4 = \text{id}$ . Performing similar rotations around the axes running through the center of the other two pairs of opposite faces, yields 6 more rotational symmetries, bringing our total thus far to 10:  $\text{id}$ , 3 of “shape”  $(13)(24)(57)(68)$ , and 6 of “shape”  $(1234)(5678)$ .

And there are more. Consider holding a pair of antipodal vertices, like 1 and 5. Let  $D_1$  denote the diagonal line (running through the interior of the cube) connecting 1 and 5. Rotating the cube by  $120^\circ$  clockwise around  $D_1$  gives the permutation  $\gamma = (247)(683)$ , and rotating by  $240^\circ$  (or  $-120^\circ$ ), yields  $\gamma^2 = (274)(638)$ . Considering the remaining three diagonals as well, we add 8 permutations of shape  $(247)(683)$  to our list of rotational symmetries, bringing our total thus far to 18.

And there are still more: we can rotate by  $180^\circ$  around the axis connecting the midpoint of opposite edges of the cube. For example, rotating about the line connecting the midpoints of edges  $\{1, 4\}$  and  $\{5, 8\}$  yields the permutation  $\beta = (14)(58)(26)(37)$ . This produces 6 more

rotational symmetries (including  $\beta$ ) of shape  $(14)(58)(26)(37)$ , and our total is now 24. Have we found all rotational symmetries? We could indeed construct a direct argument that we have found them all, but just having a complete list of the symmetries is not the end of the story. We typically also want to know more about the “structure” of the group itself.

To better understand  $\text{CUBE}^+$ , let’s change our point of view slightly. Perhaps we’ve noticed something about how the symmetries treat antipodal (i.e. opposite) vertices. For example, once we know where a given symmetry sends the vertex 1, we automatically know where 5 must be sent. In short, any symmetry of the cube (rotational or not), must move a pair of antipodal vertices to a (perhaps same) pair of antipodal vertices. This can be used to show that  $\text{CUBE}$  acts on the 4 diagonals of the cube (pictured below).



Let’s use the notation  $D_i$  for the diagonal line connecting vertex  $i$  to  $i + 4$  (so  $D_i$  connects vertices that are equal modulo 4). Our comment before is that  $\text{CUBE}$  acts on the set  $\mathcal{D} = \{D_1, D_2, D_3, D_4\}$ . However, the action of  $\text{CUBE}$  is *not* faithful on  $\mathcal{D}$ . Indeed, there is a very special reflection that takes each vertex to the antipodal vertex; let’s call this antipodal reflection  $\rho$ . Then  $\rho$  fixes every diagonal, so  $\rho$  and  $\text{id}$  have the same action on  $\mathcal{D}$ . However,  $\text{CUBE}^+$  does act faithfully on  $\mathcal{D}$ : one can argue geometrically that the only rotational symmetry fixing all four diagonals is the identity. Thus, we may view  $\text{CUBE}^+$  as a subgroup of  $\text{Sym}(\{D_1, D_2, D_3, D_4\})$ . To simplify notation, let’s forget about the  $D$ , and just label the diagonals 1, 2, 3, 4. In doing so, we may now view  $\text{CUBE}^+$  as a subgroup of  $\text{Sym}(4)$ . Let’s revisit some of the symmetries from before and see how they permute the diagonals.

Symmetry	Permuting the vertices	Permuting the diagonals
$\alpha$	$(1234)(5678)$	$(1234)$
$\alpha^2$	$(13)(24)(57)(68)$	$(13)(24)$
$\gamma$	$(247)(683)$	$(243)$
$\beta$	$(14)(58)(26)(37)$	$(14)$
$\rho \notin \text{CUBE}^+$	$(15)(26)(37)(48)$	$\text{id}$

Now, remember that we found 24 rotational symmetries. If we view  $\text{CUBE}^+ \subseteq \text{Sym}(4)$  (permuting the diagonals), then as  $|\text{Sym}(4)| = 24$ , it must be that  $\text{CUBE}^+ = \text{Sym}(4)$ . And this gives a nice description of  $\text{CUBE}^+$  as realizing all possible permutations of the 4 diagonals (in a way that different elements of  $\text{CUBE}^+$  represent different permutations of the diagonals).

And what about  $\text{CUBE}$ ? It turns out that if  $\tau \in \text{CUBE}$ , then either  $\tau \in \text{CUBE}^+$  or  $\tau = \sigma\rho$  for some  $\sigma \in \text{CUBE}^+$  (with  $\rho$  the antipodal reflection). In other words, every symmetry of the cube is either a rotational symmetry (which we understand in terms of how it permutes the diagonals) or it is a composition of a rotational symmetry with the antipodal reflection. Moreover, for each  $\sigma \in \text{CUBE}^+$ ,  $\sigma\rho = \rho\sigma$ , which (in group theoretic notation) implies that  $\text{CUBE} = \text{CUBE}^+ \times \{\text{id}, \rho\} \cong \text{Sym}(4) \times \mathbb{Z}_2$ .



### 2.3. Orbits and stabilizers.

**Example 6.** Let's look at the symmetries of a regular icosahedron. (The Rubik-esque version of this is the [Dogic](#) pictured below.) Let  $\text{Icosa}^+$  denote the set of rotational symmetries. Notice that  $\text{Icosa}^+$  acts faithfully on the 12 vertices of the icosahedron, so after labeling the vertices (as below), we can view  $\text{Icosa}^+$  as a subgroup of  $\text{Sym}(12)$ .



This time, we'll just try to count the number of symmetries in  $\text{Icosa}^+$  and leave the determination of which permutations are in  $\text{Icosa}^+$  (and  $\text{Icosa}$ ) as an exercise. Let  $\sigma \in \text{Icosa}^+$  denote an arbitrary rotational symmetry. In two-line notation,  $\sigma$  has the form

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & \cdots & 12 \\ \sigma(1) & \sigma(2) & \sigma(3) & \cdots & \sigma(12) \end{pmatrix}$$

To count  $\text{Icosa}^+$ , let's count the possible ways we can fill in the bottom row of the two-line notation for  $\sigma$ . Notice that there are rotations taking vertex 1 to any other vertex (including 1), so there are 12 possibilities for  $\sigma(1)$ .

Now assume that we have chosen  $\sigma(1)$ ; how many possibilities remain for  $\sigma(2)$ ? Notice that when counting the possibilities for  $\sigma(2)$ , the actual value of  $\sigma(1)$  does not matter, so without loss of generality, we may assume that  $\sigma(1) = 1$ , i.e.  $\sigma$  fixes 1. Since 2 is connected to 1 by an edge and  $\sigma$  fixes 1,  $\sigma(2)$  must also be connected to 1 by an edge, and we see that there are 5 possibilities for  $\sigma(2)$ , which correspond to rotations of the icosahedron about the line connecting 1 to its antipodal vertex 7.

Thus far, we found that there are 12 possibilities for  $\sigma(1)$ , and that once given  $\sigma(1)$ , there are 5 possibilities for  $\sigma(2)$ . Now assume that we know  $\sigma(1)$  and  $\sigma(2)$ . What about  $\sigma(3)$ ? Let's keep assuming that  $\sigma(1) = 1$ . Also, when counting possibilities for  $\sigma(3)$ , it does not matter which of the 5 vertices we choose for  $\sigma(2)$ , so let's assume that  $\sigma(2) = 2$  (again without any loss of generality). But if  $\sigma$  fixes the vertices 1 and 2, then the axis of rotation for  $\sigma$  passes through the edge  $\{1, 2\}$ . However, the only rotation fixing an edge of the icosahedron is the identity, meaning that the only choice for  $\sigma(3) = 3$  and in fact that  $\sigma(k) = k$  for all  $k \geq 3$ .

In summary, there are 12 possibilities for  $\sigma(1)$ , from which 5 possibilities remain for  $\sigma(2)$ , and then only one choice remains for each  $\sigma(k)$  with  $k \geq 3$ . Thus,  $|\text{Icosa}^+| = 60$ .

In Example 6, we encountered the need to discuss all possible vertices to which  $\text{Icosa}^+$  can take vertex 1. This is an important concept for all permutation groups, and so we give it a name: the *orbit* of 1 under  $\text{Icosa}^+$ . We also had need to consider all symmetries in



$\text{Icosa}^+$  that fixed the vertex 1. This is an equally important concept that we also name: the *stabilizer* of 1 in  $\text{Icosa}^+$ . Here are the proper definitions.

**Definition 2.1.** Let  $G$  act on  $X$ , and let  $y \in X$ .

- The **orbit of  $y$  under  $G$**  is defined by  $\text{Orb}_G(y) = \{x \in X \mid x = g \cdot y \text{ for some } g \in G\}$ .
- The **stabilizer of  $y$  in  $G$**  is defined by  $G_y = \{g \in G \mid g \cdot y = y\}$ .

When considering stabilizers of stabilizers, we use notation  $G_{y_1, y_2}$  in place of  $(G_{y_1})_{y_2}$ .

**Example 7.** Let's revisit Example 6 and record our findings using the language of orbits and stabilizers. Set  $G = \text{Icosa}^+$ .

We first found that  $\text{Orb}_G(1) = \{1, 2, \dots, 12\}$ . In fact, no matter which vertex  $k$  we choose,  $\text{Orb}_G(k) = \{1, 2, \dots, 12\}$ . Thus,  $G$  has only one orbit on the vertices, consisting of them all.

Next, instead of studying orbits with respect to  $G$ , we changed to considering the stabilizer  $G_1$ . Here we found that  $\text{Orb}_{G_1}(2) = \{2, 3, 4, 5, 6\}$ . We could also ask where  $G_1$  could take vertices other than 2. Since  $G_1$  fixes 1,  $\text{Orb}_{G_1}(1) = \{1\}$ . As  $G_1$  also fixes the antipodal vertex 7,  $\text{Orb}_{G_1}(7) = \{7\}$ . There are various other vertices to consider, and, for example,  $\text{Orb}_{G_1}(8) = \{8, 9, 10, 11, 12\}$ . Here is the complete list of orbits of  $G_1$ :

$$\{1\}, \{7\}, \{2, 3, 4, 5, 6\}, \{8, 9, 10, 11, 12\}.$$

Finally, we considered the stabilizer  $G_{1,2}$ . This time we found that  $G_{1,2}$  must fix every vertex, so for each vertex  $k$ ,  $\text{Orb}_{G_{1,2}}(k) = \{k\}$ .

One thing to notice in Example 7 is how the orbits of  $G_1$  *partitioned* the set  $\{1, 2, \dots, 12\}$  (into non-overlapping pieces). Of course, the same is trivially true of the orbits of  $G$  since there was only one orbit. This observation about the orbits partitioning the set being acted upon is true in general.

**Lemma.** Let  $G$  act on  $X$ . Then each  $x \in X$  is contained in one and only one orbit of  $G$ .

*Proof.* Each  $x \in X$  is contained in the orbit  $\text{Orb}_G(x)$  since  $x = \text{id} \cdot x$  for  $\text{id}$  the identity permutation. Next, suppose some  $z \in X$  is contained in both  $\text{Orb}_G(x)$  and  $\text{Orb}_G(y)$  for some  $x, y \in X$ . To prove the lemma, we want to show that this implies that the two orbits are actually the same, i.e. that  $\text{Orb}_G(x) = \text{Orb}_G(y)$ . Since  $z \in \text{Orb}_G(x) \cap \text{Orb}_G(y)$ , there exist  $h_1, h_2 \in G$  such that  $h_1 \cdot x = z = h_2 \cdot y$ . Pictorially,

$$x \xrightarrow{h_1} z \xleftarrow{h_2} y.$$

Let's prove  $\text{Orb}_G(x) \subseteq \text{Orb}_G(y)$ . Choose  $a \in \text{Orb}_G(x)$ . By definition of  $\text{Orb}_G(x)$ , there exists  $h_3 \in G$  such that  $h_3 \cdot x = a$ , so our picture becomes

$$a \xleftarrow{h_3} x \xrightarrow{h_1} z \xleftarrow{h_2} y.$$

To show  $a \in \text{Orb}_G(y)$ , we need to find some  $g \in G$  such that  $g \cdot y = a$  (pictorially:  $a \xleftarrow{g} y$ ). The key thing to remember is that  $G$  represents a permutation group, so  $G$  is closed under inverses and compositions. Thus,  $h_1^{-1} \in G$ , so also  $h_3 h_1^{-1} h_2 \in G$ . Let  $g = h_3 h_1^{-1} h_2$ . Then

$$g \cdot y = h_3 h_1^{-1} h_2 \cdot y = h_3 h_1^{-1} \cdot z = h_3 \cdot x = a,$$

so  $a \in \text{Orb}_G(y)$ . The proof that  $\text{Orb}_G(y) \subseteq \text{Orb}_G(x)$  is similar.  $\square$

Returning to Example 6, note that not only did we consider orbits and stabilizers, but we also combined information about the two in order to count the size of the set being acted upon. The next theorem formalizes that process. In the statement of the theorem we use the notation  $|A|$  to denote the cardinality (i.e. the “size”) of the set  $A$ .

**Theorem.** *Let  $G$  act on  $X$ , and let  $x \in X$ . Then  $|G| = |\text{Orb}_G(x)| \cdot |G_x|$ .*

*Proof.* We define a function  $\phi : G \rightarrow \text{Orb}_G(x)$  via  $\phi(g) = g \cdot x$ , so  $g \cdot x = y \iff \phi(g) = y$ . By definition of  $\text{Orb}_G(x)$ ,  $\phi$  maps *onto*  $\text{Orb}_G(x)$ , but  $\phi$  is not likely one-to-one. For example, every element of  $G_x$  maps to  $x$ . Here is a picture.



For a given  $g \in G$ , let's count the number of  $h \in G$  such that  $\phi(g) = \phi(h)$ . Notice that

$$\phi(g) = \phi(h) \iff g \cdot x = h \cdot x \iff x = g^{-1}h \cdot x \iff g^{-1}h \in G_x \iff h \in gG_x,$$

where we use the notation  $gG_x$  to denote the set  $\{ga \mid a \in G_x\}$ . So, counting the number of  $h$  such that  $\phi(g) = \phi(h)$  is the same as determining the cardinality of  $gG_x$ . Now notice that for all  $a, b \in G_x$ ,

$$ga = gb \iff a = g^{-1}gb \iff a = b,$$

which tells us that counting the number of  $ga \in gG_x$  is the same as counting the number of  $a \in G_x$ . Thus,  $|gG_x| = |G_x|$ , and crucially, this does *not* depend on the choice of  $g$ . So each “fiber” in the picture above has the same size, which is  $|G_x|$ .

We conclude that for every  $y \in \text{Orb}_G(x)$ , there are precisely  $|G_x|$ -many elements of  $G$  that map to  $y$ . Thus,  $|G| = |\text{Orb}_G(x)| \cdot |G_x|$ .  $\square$

**Example 8.** Let's revisit Example 6 yet again. Set  $G = \text{Icosa}^+$ .

The previous theorem says that  $|G| = |\text{Orb}_G(1)| \cdot |G_1|$ . Since we know that  $\text{Orb}_G(1) = \{1, 2, \dots, 12\}$ , we get that  $|G| = 12 \cdot |G_1|$ .

We can now apply the same theorem to  $G_1$  to get that  $|G_1| = |\text{Orb}_{G_1}(2)| \cdot |G_{1,2}|$ , so as  $\text{Orb}_{G_1}(2) = \{2, 3, 4, 5, 6\}$ ,  $|G_1| = 5 \cdot |G_{1,2}|$ . Combining with what we've already learned, we get that  $|G| = 12 \cdot 5 \cdot |G_{1,2}|$ .

And finally, we also know that  $G_{1,2}$  must fix every vertex, but the only element of  $G$  fixing all vertices is the identity. Thus,  $G_{1,2} = \{\text{id}\}$ , so  $|G_{1,2}| = 1$ . Thus,  $|G| = 12 \cdot 5 \cdot 1 = 60$ .

### 3. DEGREES OF SYMMETRY

Which would you say is “more symmetric”: a regular tetrahedron or a cube? Why? A first thought may be that an object is more symmetric if it has more symmetries. With that point of view, a cube would be more symmetric than a tetrahedron since our work in Examples 1 and 5 implies that  $|\text{CUBE}| = 48 > 24 = |\text{TETRA}|$ . However, this approach seems to privilege the cube simply because it has more vertices.

Let's work towards another approach to measuring symmetry. We saw that there is a symmetry taking any vertex of the tetrahedron to any other, and the same is true for a cube. But now, what if we ask about pairs of vertices. Notice that any two vertices of a tetrahedron can be simultaneously moved to any other two vertices. However, the same is *not* true for the cube: a pair of antipodal vertices can never be simultaneously moved to a pair of non-antipodal vertices. And this difference suggests that we might view the tetrahedron as being more symmetric than the cube. Let's formalize this idea.

**Notation.** For  $X$  a set,  $X^n$  denotes the set of all  $n$ -tuples with entries from  $X$ , and  $X^{(n)}$  denotes the subset of  $X^n$  consisting of tuples with distinct entries, i.e.

$$X^{(n)} = \{(x_1, \dots, x_n) \in X^n \mid x_i \neq x_j \text{ for all } i \neq j\}.$$

For example,  $(1, 2, 3, 2) \notin \mathbb{Z}^{(4)}$  because of the repeated 2, but  $(1, 2, 3, 7) \in \mathbb{Z}^{(4)}$  as there is no repetition in the coordinates.

**Remark.** If  $G$  acts on  $X$ , then  $G$  acts coordinatewise on both  $X^n$  and  $X^{(n)}$  via the rule

$$g \cdot (x_1, \dots, x_n) = (g \cdot x_1, \dots, g \cdot x_n).$$

The main reason for looking at the action of  $G$  on  $X^n$  or  $X^{(n)}$  is to study how a group element  $g$  *simultaneously* acts on  $x_1, \dots, x_n$  instead of simply studying how  $g$  acts on single elements of  $X$ . This better captures relationships between  $x_1, \dots, x_n$  (similar to how pairs of vertices on a cube might be antipodal or not).

**Definition.** We say that an action of  $G$  on  $X$  is **transitive** if  $G$  has only one orbit on  $X$ . More generally, we say the action is  **$k$ -transitive** if  $G$  has only one orbit on  $X^{(k)}$  (with respect to the coordinatewise action). Thus, the action is  $k$ -transitive if for all  $(x_1, \dots, x_k), (y_1, \dots, y_k) \in X^{(k)}$  there exists some  $g \in G$  such that  $g \cdot x_i = y_i$  for all  $1 \leq i \leq k$ .

**Example 9.** Let's revisit Examples 1 and 5 through the lens of  $k$ -transitivity.

The comment above that any two (distinct) vertices of a tetrahedron can be moved to any other two (distinct) vertices is stating that the action of TETRA on the vertices of the tetrahedron is 2-transitive. In fact, such pairs of vertices can be moved using a rotation, so  $\text{TETRA}^+$  also acts 2-transitively on the vertices. And more is true: TETRA in fact acts 4-transitively on the vertices.

As for the cube, CUBE and  $\text{CUBE}^+$  act transitively on the vertices, but neither act 2-transitively. However,  $\text{CUBE}^+$  does act 4-transitively on the four diagonals of the cube.

Actions that are  $k$ -transitive for "large" values of  $k$  can be viewed as encoding a "large" degree of symmetry. The rest of this section is devoted to hunting for highly symmetric structures by looking for  $k$ -transitive actions with  $k$  as large as possible. The next example shows that actions of arbitrarily large transitivity degree do exist, but we should at the same time think of these as rather boring examples. Ignoring them, our quest for highly transitive actions becomes much harder (and much more interesting).

**Example 10.** If  $X = \{1, \dots, n\}$ , then it is fairly straightforward to verify that  $\text{Sym}(n)$  acts  $n$ -transitively on  $X$ , and it is ever so slightly less-straightforward to check that the action of  $\text{Alt}(n)$  is  $(n - 2)$ -transitive but not  $(n - 1)$ -transitive.

**3.1. Affine and projective linear groups.** Let's meet some new permutation groups as we continue to investigate  $k$ -transitivity.


**Example 11.** Let's take a look at a particular group of permutations on  $\mathbb{R}$ . Specifically, we will consider the set of invertible (affine) linear functions:

$$\text{AGL}_1(\mathbb{R}) = \{f \in \text{Sym}(\mathbb{R}) \mid f(x) = mx + b \text{ for some } m, b \in \mathbb{R} \text{ with } m \neq 0\}.$$

It can be checked that  $\text{AGL}_1(\mathbb{R})$  satisfies the necessary closure axioms to be a permutation group on  $\mathbb{R}$ . Let's determine how transitive this action is.

Is the action transitive? Yes it is. For all  $r, s \in \mathbb{R}$ , the translation  $f(x) = x + (s - r)$  is in  $\text{AGL}_1(\mathbb{R})$  and  $f(r) = s$ , so  $\text{AGL}_1(\mathbb{R})$  has a single orbit on  $\mathbb{R}$ .

What about 2-transitivity? Let's consider  $(r_1, r_2), (s_1, s_2) \in \mathbb{R}^{(2)}$ . We need to determine if there exists  $f \in \text{AGL}_1(\mathbb{R})$  such that  $f(r_1, r_2) = (s_1, s_2)$ , which would mean  $f(r_1) = s_1$  and  $f(r_2) = s_2$ . If we graph the points  $(r_1, s_1)$  and  $(r_2, s_2)$  (remembering that  $r_1 \neq r_2$  and  $s_1 \neq s_2$  by definition of  $\mathbb{R}^{(2)}$ ), we have something like the following:



The function representing the line connecting the two points will take  $r_1$  to  $s_1$  and  $r_2$  to  $s_2$ , which is what we want, so the action is indeed 2-transitive.

Could it be 3-transitive? If it were, there would have to be some  $f \in \text{AGL}_1(\mathbb{R})$  such that  $f(0, 1, 2) = (0, 1, 3)$ , meaning that  $f(0) = 0$ ,  $f(1) = 1$ , and  $f(2) = 3$ . Writing  $f(x) = mx + b$ ,  $f(0) = 0$  implies  $b = 0$ , and then  $f(1) = 1$  implies  $a = 1$ . Thus  $f$  is the identity function  $f(x) = x$ , but then there is no way that  $f(2) = 3$ . So no, the action is not 3-transitive.

In Example 11, we considered the group of invertible linear functions from  $\mathbb{R}$  to  $\mathbb{R}$ , but we used very little specific knowledge of  $\mathbb{R}$ . For example, we would have reached the same conclusion for invertible linear functions from  $\mathbb{Q}$  to  $\mathbb{Q}$ . As such, we will define  $\text{AGL}_1(F)$  where  $F$  is any **field**; roughly, a **field** is a mathematical structure with “well-behaved” notions of addition and multiplication as well as additive and multiplicative inverses. Examples of fields include  $\mathbb{Q}$ ,  $\mathbb{R}$ , and  $\mathbb{C}$ , but not  $\mathbb{Z}$  because of the lack of multiplicative inverses for most elements. There are also examples of finite fields like  $\mathbb{Z}_p$ ; here  $\mathbb{Z}_p$  represents (congruence classes of) the integers with arithmetic performed modulo a prime number  $p$ .

**Definition.** For  $F$  a field, we define

$$\text{AGL}_1(F) = \{f \in \text{Sym}(F) \mid f(x) = mx + b \text{ for some } m, b \in F \text{ with } m \neq 0\}.$$

**Remark.** Essentially the same proof we gave in Example 11 shows that, for any field  $F$ ,  $\text{AGL}_1(F)$  acts 2-transitively on  $F$ . Moreover, if  $|F| \geq 4$ , then the action is *not* 3-transitive.

The group  $\text{AGL}_1(F)$  can be generalized in at least a couple of ways. One way is to directly adapt the definition to give functions from the vector space  $F^n$  to itself by considering functions  $f(x) = mx + b$  where  $m$  is an  $n \times n$  matrix and  $b$  an  $n$ -vector.

**Definition.** For  $F$  a field, let  $\text{GL}_n(F)$  denote the set of all *invertible*  $n \times n$  matrices with entries from  $F$ , and define


$$\text{AGL}_n(F) = \{f \in \text{Sym}(F^n) \mid f(\bar{x}) = A\bar{x} + \bar{b} \text{ for some } A \in \text{GL}_n(F) \text{ and some } \bar{b} \in F^n\}.$$

We refer to  $\text{AGL}_n(F)$  as the **affine general linear group** of degree  $n$ .

**Remark.** We won't explore  $\text{AGL}_n(F)$  here, but it's worth mentioning that—just like  $\text{AGL}_1(F)$ —each permutation group in this family (for all choices of  $n$  and  $F$ ) acts 2-transitively on the vector space  $F^n$ , and provided  $|F| \geq 4$ , the action is not 3-transitive.

Let's now take a look at another way to generalize the construction of  $\text{AGL}_1(F)$ . This time we will encounter a family of permutation groups that are typically 3-transitive but not 4-transitive.

**Example 12.** In Example 11, we looked at certain functions on the line  $\mathbb{R}$ ; this time we will work with the projective line  $\mathbb{P}^1(\mathbb{R}) = \mathbb{R} \cup \{\infty\}$ . Right now,  $\infty$  is just another symbol we are adding to the set  $\mathbb{R}$ , but we can think of  $\mathbb{P}^1(\mathbb{R})$  as a circle where  $\infty$  connects the positive and negative “ends” of the real line together.



We now define the so-called **fractional linear transformations** of the projective line:

$$\text{PGL}_2(\mathbb{R}) = \left\{ f \in \text{Sym}(\mathbb{P}^1(\mathbb{R})) \mid f(x) = \frac{ax+b}{cx+d} \text{ for } a, b, c, d \in \mathbb{R} \text{ with } ad - bc \neq 0 \right\}.$$


In order for a function of the form  $f(x) = \frac{ax+b}{cx+d}$  to be fully defined on  $\mathbb{P}^1(\mathbb{R})$ , we need to make sense of a couple values; set

$$f(-d/c) = \infty$$

$$f(\infty) = \begin{cases} \frac{a}{c} & \text{if } c \neq 0 \\ \infty & \text{if } c = 0. \end{cases}$$

The input  $-d/c$  results in the denominator of  $f(x)$  being 0, so  $f(-d/c) = \infty$  can be thought of via the arithmetic  $r/0 = \infty$  for each nonzero  $r \in \mathbb{R}$ . One way to make sense of  $f(\infty)$  is to think of  $\lim_{x \rightarrow \infty} f(x)$ ; this leads to  $f(\infty) = a/c$ , which is reasonably interpreted as  $\infty$  in the case when  $c = 0$ .

Note that  $\text{AGL}_1(\mathbb{R}) \subset \text{PGL}_2(\mathbb{R})$  by using  $c = 0$  and  $d = 1$  in the definition above. Moreover, every element of  $\text{AGL}_1(\mathbb{R})$  fixes  $\infty$ , and in fact,  $\text{AGL}_1(\mathbb{R})$  is *equal* to the stabilizer of  $\infty$  in  $\text{PGL}_2(\mathbb{R})$ . Also note that the “inversion” function  $i(x) = \frac{1}{x}$  is in  $\text{PGL}_2(\mathbb{R})$ . If we scale the circle above so that  $-1$  and  $1$  are the endpoints of the horizontal diameter, then the function  $i(x)$  can be viewed as a reflection over that diameter (with  $i(0) = \infty$  and  $i(\infty) = 0$ ).



We now work to show that  $\text{PGL}_2(\mathbb{R})$  acts 3-transitively on  $P^1(\mathbb{R})$ . Set  $G = \text{PGL}_2(\mathbb{R})$  and  $X = P^1(\mathbb{R})$ .

Let's first prove that the action is transitive. We'll do this by showing that  $r \in \text{Orb}_G(\infty)$  for every  $r \in X$  (implying that  $G$  has only one orbit on  $X$ ). If  $r = \infty$ , then  $r \in \text{Orb}_G(\infty)$  since  $\infty = \text{id}(\infty)$  (where  $\text{id}(x) = x$ ). Suppose now that  $r \neq \infty$ ; then  $r \in \mathbb{R}$ . Since  $\text{AGL}_1(\mathbb{R})$  is transitive on  $\mathbb{R}$ , there exists an  $f \in \text{AGL}_1(\mathbb{R}) \subset G$  such that  $f(r) = 0$ . And then, with  $i(x)$  the inversion function defined above, we find that  $i \circ f(r) = i(0) = \infty$ , which shows that  $r \in \text{Orb}_G(\infty)$ .

To show  $G$  acts 3-transitively, we will show that  $(r_1, r_2, r_3) \in \text{Orb}_G(0, 1, \infty)$  for every  $(r_1, r_2, r_3) \in X^{(3)}$ . Since we have shown  $G$  acts transitively on  $X$ , there exists  $g \in G$  such that  $g(r_3) = \infty$ ; thus  $g(r_1, r_2, r_3) = (s_1, s_2, \infty)$  for some  $s_1, s_2 \in X$ . Because  $r_1, r_2, r_3$  are all distinct and  $g$  is a one-to-one,  $s_1, s_2, \infty$  are all distinct. In particular,  $s_1, s_2 \in \mathbb{R}^{(2)}$ , so by 2-transitivity of  $\text{AGL}_1(\mathbb{R})$ , there exists  $f \in \text{AGL}_1(\mathbb{R})$  such that  $f(s_1, s_2) = (0, 1)$ . As mentioned before, the elements of  $\text{AGL}_1(\mathbb{R})$  fix  $\infty$ , so  $f(\infty) = \infty$ . Pictorially,

$$\begin{array}{ccccc} & g & & f & \\ r_1 & \rightarrow & s_1 & \rightarrow & 0 \\ r_2 & \rightarrow & s_2 & \rightarrow & 1 \\ r_3 & \rightarrow & \infty & \rightarrow & \infty \end{array}$$

Thus,  $f \circ g(r_1, r_2, r_3) = (0, 1, \infty)$ , so  $(r_1, r_2, r_3) \in \text{Orb}_G(0, 1, \infty)$  as desired.

Might the action be 4-transitive? If so, there would be an  $g \in G$  such that  $g(0, 1, \infty, 2) = (0, 1, \infty, 3)$ . Since  $g(\infty) = \infty$ , we find that  $g \in \text{AGL}_1(\mathbb{R})$ , but then (as we saw in Example 11),  $g(0) = 0$  and  $g(1) = 1$  imply that  $g$  is the identity function. Thus, there is no chance that  $g(2) = 3$ . So the action is not 4-transitive.

As with our exploration of  $\text{AGL}_1(\mathbb{R})$ , our work with  $\text{PGL}_2(\mathbb{R})$  used very few specific properties of  $\mathbb{R}$ , and we are again able to define  $\text{PGL}_2(F)$  for  $F$  any field.

**Definition.** For  $F$  a field, we define  $P^1(F) = F \cup \{\infty\}$  and

$$\text{PGL}_2(F) = \left\{ f \in \text{Sym}(P^1(F)) \mid f(x) = \frac{ax + b}{cx + d} \text{ for } a, b, c, d \in F \text{ with } ad - bc \neq 0 \right\},$$

where

$$f(-d/c) = \infty \text{ and } f(\infty) = \begin{cases} \frac{a}{c} & \text{if } c \neq 0 \\ \infty & \text{if } c = 0. \end{cases}$$

**Remark.** Our work in Example 12 easily generalizes to show that, for any field  $F$ ,  $\text{PGL}_2(F)$  acts 3-transitively on  $P^1(F)$ ; the action is *not* 4-transitive provided  $|P^1(F)| \geq 5$  (or equivalently, if  $|F| \geq 4$ ).

As you might expect from the notation,  $\text{PGL}_2(F)$  can be generalized to obtain the **projective linear groups**  $\text{PGL}_n(F)$ . Where  $\text{PGL}_2(F)$  acts naturally on the projective line  $P^1(F)$  (obtained from  $F$  by adding a point at infinity),  $\text{PGL}_n(F)$  acts on projective  $n$ -space  $P^n(F)$ ,

which is obtained from  $F^n$  by adding an entire  $(n - 1)$ -dimensional space  $F^{n-1}$  at infinity. We will not properly define  $\text{PGL}_n(F)$  here, but it's worth mentioning that the action of  $\text{PGL}_n(F)$  on  $\mathbb{P}^n(F)$  is *not* 3-transitive when  $n \geq 3$  (but it is 2-transitive).

We have now seen (families of) examples of 2- and 3-transitive groups, so what about 4-transitive actions? The rest of our work in this section will focus on precisely that question.


**Question.** Are there examples of 4-transitive actions other than those naturally associated to  $\text{Sym}(n)$  and  $\text{Alt}(n)$  (as in Example 10)?

It turns out that the answer is yes, but if we only consider finite groups, then the number of such examples can be counted on just one hand! Let's meet some of them.

**3.2. Mathieu groups.** The **Mathieu groups**, denoted  $M_{11}$ ,  $M_{12}$ ,  $M_{22}$ ,  $M_{23}$ , and  $M_{24}$ , are a special collection of permutation groups that act on, respectively, sets of size 11, 12, 22, 23, and 24. These groups can be described many different ways. We'll focus on a construction of  $M_{12}$  given by Conway and Sloane [CS99] and further concretized by examples using GAP in a [blogpost by Nasmith](#). From that, we will also be able to describe  $M_{11}$ , but we won't say much about the other three.

**Example 13** ([CS99, Section 18 of Chapter 11] and [Nas]). We return to the icosahedron. This time, instead of looking at its symmetries, we'll describe a type of game to scramble the vertices via certain allowable moves. The moves we will describe are similar in spirit to those you could perform on a [Dogic](#) (pictured in Example 6), but they cannot actually be done on one.

Holding the icosahedron so that vertex  $i$  is pointed up, let  $\tau_i$  denote a clockwise “twist” of the five vertices connected to vertex  $i$ ;  $\tau_i$  does not move any vertices that are not connected to vertex  $i$ . The twists  $\tau_1 = (2, 3, 4, 5, 6)$  and  $\tau_5 = (1, 4, 8, 9, 6)$  are illustrated below using our labeling of the vertices from before. Notice that we are using commas in our cycle notation; this is to avoid confusion when using vertices larger than 9 later on.



One game we could play is to see all of the ways in which we can scramble the vertices by repeatedly using any of the twelve twists. The collection of all permutations of the vertices arising from this process will be denoted  $T$  for the *twist group*. For example,  $\tau_5\tau_1\tau_1\tau_4 \in T$ , and we can work out that

$$\begin{aligned} \tau_5\tau_1\tau_1\tau_4 &= (1, 4, 8, 9, 6)(2, 3, 4, 5, 6)(2, 3, 4, 5, 6)(2, 3, 4, 5, 6)(1, 3, 12, 8, 5) \\ &= (1, 2)(3, 6)(4, 5, 12, 8, 9) \end{aligned}$$



In the language of group theory,  $T$  is the group **generated** by the twists; this just means  $T$  is the smallest permutation group on the vertices that contains the twists. We write this in symbols as  $T = \langle \tau_i \mid 1 \leq i \leq 12 \rangle$ .

One question whose answer may not be so clear is: can we generate *all* possible permutations of the vertices using these twists? In other words, is it true that  $T = \text{Sym}(12)$ ? Try to work it out a bit... but the answer is no. However, the twists can generate a lot of the permutations (in fact half of them), and it turns out that  $T = \text{Alt}(12)$ .

Now let's slightly change the game. This time, every time we perform a clockwise twist  $\tau_i$  it must be followed immediately by some counter-clockwise "untwist"  $\tau_j^{-1}$ . The set of permutations realized using this process will be denoted  $TU$  for the *twist-untwist* group. The group  $TU$  is generated by permutations of the form  $\tau_j^{-1}\tau_i$ ; or symbolically,  $TU = \langle \tau_j^{-1}\tau_i \mid 1 \leq i, j \leq 12 \rangle$ .

What permutations can be generated from our twist-untwist game? We can easily write down some like  $\tau_5^{-1}\tau_1 = (1, 2, 3, 4)(5, 6, 9, 8)$  and  $\tau_2^{-1}\tau_1\tau_5^{-1}\tau_7\tau_1^{-1}\tau_7 = (3, 9, 11, 8)(5, 12, 10, 6)$ , but, for example, how many permutations are in  $TU$ ? As mentioned above, the twist group  $T = \text{Alt}(12)$  contains half of the permutations of the vertices. In contrast,  $TU$  contains less than 1/5000 of the permutations. It so happens that twist-untwist group is the same as the so-called Mathieu group  $M_{12}$ , although  $M_{12}$  was not initially thought of this way. Thus,

$$M_{12} = TU = \langle \tau_j^{-1}\tau_i \mid 1 \leq i \leq 12 \rangle.$$

Additionally, the Mathieu group  $M_{11}$  is just the stabilizer of the vertex 1 in  $M_{12}$ .

Let's investigate the degree of transitivity of the action of  $M_{12}$  on the vertices. Is the action transitive? Is every vertex in the orbit of 1 under the action of  $M_{12}$ ? Notice that with the exception of vertex 7, every other vertex shares a common neighboring vertex with 1. For example, 1 and 9 have a common neighbor of 5. Thus, we can move 1 to 9 with a series of twists around 5:  $\tau_5^3(1) = 9$ . Of course,  $\tau_5^3$  is *not* an element of  $M_{12}$ , so let's adjust things. Notice how the twist around the antipodal vertex 11 does not interact with the twist around 5, and it can be checked that  $(\tau_{11}^{-1}\tau_5)^3$  also takes 1 to 9. In a similar way, we see how to take 1 to any vertex other than 7 by an element of  $M_{12}$ . But it's not hard to also find a permutation taking 1 to 7, e.g.  $\tau_{10}^{-1}\tau_{12}\tau_1^{-1}\tau_5$ . So yes, the action of  $M_{12}$  is transitive.

Is the action 2-transitive? Can we simultaneously take vertices 1 and 2 to any other pair of distinct vertices  $x$  and  $y$  under the action of  $M_{12}$ ? Using inverses, this is the same as asking if we can take  $x$  and  $y$  to 1 and 2. We know that we can take  $x$  to 1, so there is some  $\alpha \in M_{12}$  such that  $\alpha(x, y) = (1, y')$ . Thus, if we can find  $\beta \in M_{12}$  such that  $\beta(1, y') = (1, 2)$ , we will be done. In other words, we are asking if the stabilizer of 1 in  $M_{12}$  (which is  $M_{11}$ ) can take an arbitrary  $y'$  that is different from 1 to the vertex 2. So, can this be done? Can we find a sequence of twist-untwist permutations that do not move 1, but take an arbitrary  $y' \neq 1$  to 2? Try! You'll should be successful, and we can then conclude that  $M_{12}$  acts 2-transitively on the the vertices.

In fact,  $M_{12}$  acts 5-transitively on the vertices! Moreover, the stabilizer in  $M_{12}$  of five distinct vertices is just the identity permutation, which implies that  $M_{12}$  is *not* 6-transitive. Also, that fact that  $M_{12}$  acts 5-transitively can be used to quickly see that the stabilizer of the vertex 1 acts 4-transitively, so  $M_{11}$  is an example of a 4-transitive group (as is  $M_{12}$ ).

We are now in a position to address our earlier question: are there examples of 4-transitive actions other than those naturally associated to  $\text{Sym}(n)$  and  $\text{Alt}(n)$ ? Yes, there are, but more importantly, in the case of finite permutation groups, we know them all.

**Theorem** (see for example [Cam99, Theorem 4.11]). *The only finite permutation groups with a 4-transitive action are the symmetric groups, the alternating groups, and the Mathieu groups  $M_{11}$ ,  $M_{12}$ ,  $M_{23}$ , and  $M_{24}$ . Consequently, the only finite permutation groups with a 6-transitive action are the symmetric and alternating groups.*

The proof of this theorem is quite deep, utilizing the [classification of the finite “simple” groups](#). A further consequence of this classification project is that all finite permutation groups with a 2-transitive action are known. This list is quite a bit longer than for 4-transitive actions, including, for example, the affine and projective groups  $\text{AGL}_1(F)$  and  $\text{PGL}_2(F)$  (for  $F$  a finite field) as well as their generalizations to  $\text{AGL}_n(F)$  and  $\text{PGL}_n(F)$ .


#### 4. RELATIONAL COMPLEXITY

Let’s now take a look at a relatively new topic about permutation groups, known as *relational complexity*, which has its roots in an area of mathematical logic known as [model theory](#). Although quite a bit of research has occurred recently around relational complexity, there are many questions that remain. The definition can be challenging to wrap one’s head around at first, and we’ll spend some time unpacking what it means. However, the (very important) question of why one would consider this definition in the first place is best answered in model-theoretic terms, but we will simply refer out for that: see for example [Che00, Section 2] or [Wis16, Section 2.2]. Sorry!


**Definition.** Let  $G$  be a permutation group acting on  $X$ .

- (1) Tuples  $\bar{x}, \bar{y} \in X^k$  are said to be  **$r$ -equivalent**, denoted  $\bar{x} \sim_r \bar{y}$ , if every  $r$  elements from  $\bar{x}$  are in the same orbit as the *corresponding*  $r$  elements from  $\bar{y}$ .
- (2) The **relational complexity** of the action, denoted  $\text{RC}(G, X)$ , is the *smallest*  $r$  such that  $\bar{x} \sim_r \bar{y}$  implies  $\bar{x} \sim_k \bar{y}$  for all  $k \geq r$  and all  $\bar{x}, \bar{y} \in X^k$ .

Let’s first just think about the difference between 2- and 3-equivalence. Suppose we have tuples  $\bar{x} = (x_1, x_2, x_3)$  and  $\bar{y} = (y_1, y_2, y_3)$ . If  $\bar{x} \sim_2 \bar{y}$ , then every 2 elements from  $\bar{x}$  are in the same orbit as the corresponding 2 elements from  $\bar{y}$ . The two-element subtuples from  $\bar{x}$  are  $(x_1, x_2)$ ,  $(x_1, x_3)$ , and  $(x_2, x_3)$ , so there must exist three (potentially different) group elements  $g_1, g_2, g_3 \in G$  such that  $g \cdot (x_1, x_2) = (y_1, y_2)$ ,  $g \cdot (x_1, x_3) = (y_1, y_3)$ , and  $g \cdot (x_2, x_3) = (y_2, y_3)$ . We might picture this as follows.



However, if the tuples are 3-equivalent, then there exists a single  $g \in G$  that simultaneously maps all three elements of  $\bar{x}$  to the corresponding elements of  $\bar{y}$ , i.e.  $g \cdot (x_1, x_2, x_3) = (y_1, y_2, y_3)$ .



Notice, in particular, how  $\bar{x} \sim_3 \bar{y}$  implies  $\bar{x} \sim_2 \bar{y}$  since we could use the same  $g$  for each of  $g_1, g_2, g_3$ , but  $\bar{x} \sim_2 \bar{y}$  does not necessarily imply  $\bar{x} \sim_3 \bar{y}$ .

**Lemma.** *Let  $G$  be a permutation group acting on  $X$ . If  $|G| \geq 2$ , then  $\text{RC}(G, X) \geq 2$ .*

*Proof.* To establish a lower bound on the relational complexity (as we desire to do here), we typically aim produce an example showing the complexity can be no lower. In this case, we will find tuples  $(x_1, x_2), (y_1, y_2) \in X^2$  such that  $(x_1, x_2) \sim_1 (y_1, y_2)$  but  $(x_1, x_2) \not\sim_2 (y_1, y_2)$ ; since 1-equivalence does not imply 2-equivalence, the complexity must be larger than 1.

As  $|G| \geq 2$ , we may choose  $g \in G$  such that  $g$  is not the identity. Then there exists some  $a \in X$  such that  $g \cdot a \neq a$ . Set  $b = g \cdot a$ , and consider the tuples  $(a, a)$  and  $(a, b)$ . These are 1-equivalent because the identity maps the first coordinate of  $(a, a)$  to the first coordinate of  $(a, b)$ , and  $g$  does the same for the second coordinates. We now explain why  $(a, a) \not\sim_2 (a, b)$ . Suppose to the contrary that  $(a, a) \sim_2 (a, b)$ ; then  $h \cdot (a, a) = (a, b)$  for some  $h \in G$ . This means that  $h \cdot a = a$  and  $h \cdot a = b$ . Since  $h \cdot a$  must have a single value and  $a \neq b$ , we arrive at a contradiction. Thus,  $(a, a) \not\sim_2 (a, b)$ .  $\square$

One (likely unclear) consequence of the previous lemma is that when analyzing the relational complexity of an action, we can focus on tuples with *distinct* entries (i.e. tuples from  $X^{(k)}$ ), just like we did when analyzing  $k$ -transitivity. Let's formalize this comment.

**Lemma.** *Let  $G$  be a permutation group acting on  $X$  with  $|G| \geq 2$ . If  $\bar{x} \sim_r \bar{y}$  implies  $\bar{x} \sim_k \bar{y}$  for all  $k \geq r$  and all  $\bar{x}, \bar{y} \in X^{(k)}$ , then  $\text{RC}(G, X) \geq r$ .*

*Proof.* To show  $\text{RC}(G, X) \geq r$ , we must show  $\bar{x} \sim_r \bar{y}$  implies  $\bar{x} \sim_k \bar{y}$  for all  $k \geq r$  and all  $\bar{x}, \bar{y} \in X^k$ . Let  $\bar{x}, \bar{y} \in X^k$ , and assume that  $\bar{x} \sim_r \bar{y}$ . By Lemma 4, we know  $\text{RC}(G, X) \geq 2$ , so if  $r$  is 1 or 2, the conclusion holds immediately. Thus, we may assume  $r \geq 3$ , although all we really will need is that  $r \geq 2$ .

Suppose that  $\bar{x}$  has repeated entries, say  $x_i = x_j$ . Then we may use that  $\bar{x} \sim_r \bar{y}$  with  $r \geq 2$  to see that there is some  $g \in G$  such that  $g \cdot x_i = y_i$  and  $g \cdot x_j = y_j$ , so  $y_i = y_j$  as well. Thus, repeated entries in  $\bar{x}$  correspond to repeated entries in  $\bar{y}$  and vice versa. Removing all repeated entries from  $\bar{x}$  and the corresponding repeated entries from  $\bar{y}$  yields new tuples  $\bar{x}', \bar{y}' \in X^{(\ell)}$  for some  $\ell \leq k$  (where  $k - \ell$  counts the number of entries removed). Note that we still have  $\bar{x}' \sim_r \bar{y}'$ , so we can apply the assumption of the lemma to see that  $\bar{x}' \sim_\ell \bar{y}'$ . Thus, there exists some  $h \in G$  such that  $h \cdot \bar{x}' = \bar{y}'$ . Adding back in the repeated entries, we have  $h \cdot \bar{x} = \bar{y}$ , so  $\bar{x} \sim_k \bar{y}$ , as desired.  $\square$

#### 4.1. First examples.

**Example 14.** Set  $X = \{1, \dots, n\}$ , and let's work out  $\text{RC}(\text{Sym}(n), X)$  and  $\text{RC}(\text{Alt}(n), X)$ .

In the first case, we find that  $\text{RC}(\text{Sym}(n), X) = 2$ . To see this, remember that we already know that  $\text{RC}(\text{Sym}(n), X) \geq 2$

## REFERENCES

- [Cam99] Peter J. Cameron. *Permutation groups*, volume 45 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1999.
- [Che00] Gregory Cherlin. Sporadic homogeneous structures. In *The Gelfand Mathematical Seminars, 1996–1999*, Gelfand Math. Sem., pages 15–48. Birkhäuser Boston, Boston, MA, 2000.
- [CS99] J. H. Conway and N. J. A. Sloane. *Sphere packings, lattices and groups*, volume 290 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, third edition, 1999. With additional contributions by E. Bannai, R. E. Borcherds, J. Leech, S. P. Norton, A. M. Odlyzko, R. A. Parker, L. Queen and B. B. Venkov.
- [Ern21] Dana C. Ernst. An inquiry-based approach to abstract algebra. Available at <https://github.com/dcernst/IBL-AbstractAlgebra>, 2021.
- [Nas] Ben Nasmith. The rubik’s icosahedron and the mathieu group m12. <https://www.cantorsparadise.com/the-rubiks-icosahedron-and-the-mathieu-group-m12-444ae64d1ac8>. Accessed: 2022-04-11.
- [Wis16] Joshua Wiscons. A reduction theorem for primitive binary permutation groups. *Bull. Lond. Math. Soc.*, 48(2):291–299, 2016.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE UNIVERSITY, SACRAMENTO, SACRAMENTO, CA 95819, USA

*Email address:* `joshua.wiscons@csus.edu`