

Discrete Probability Distributions

Math 122

Random Variables

- A variable x whose value is determined by the outcome of an experiment.
- $P(x)$ = probability of a particular value of x
- Mean/Expected Value: $\mu = \sum xP(x)$
 - If the experiment is repeated many times and the values of x are averaged, the average should be near μ .
- Standard Deviation: σ

Range Rule of Thumb

- Usual values are between

$$\mu - 2\sigma \text{ and } \mu + 2\sigma$$

5% Rule

- If $P(x \leq N) \leq 5\%$, then N is unusually low
- If $P(x \geq N) \leq 5\%$, then N is unusually low

Special Distributions

- We want a few specific, common distributions so that we know what to do when we encounter them.
- Discrete
 - Binomial (counting successes in trials)
 - Poisson (counting events in an interval)
- Continuous
 - Uniform (simple)
 - Normal (pervasive bell curve)
 - t, F, χ^2

Binomial Distribution

Binomial Distribution

- A fixed number of trials is repeated.
- The trials are independent.
- Each trial ends in success or failure.
- The probability of success is the same for each trial.
- The value of x is the number of successes.

Examples of Binomial Distributions

- The number of female children out of 10 randomly selected children
- The number of green peas in sets of 5 offspring peas.
- The number of correct responses when you guess on a multiple choice test.
- The number of Republicans among sets of 1000 random voters.

Binomial Distribution Pre-Example

What is the probability of getting exactly 3 T/F questions correct when you guess on 4?

Binomial Distribution Notation

- n = number of trials
- p = probability of success
- q = probability of failure = $1-p$
- $P(x)$ = probability of getting exactly x successes in n trials.

A formula we won't use

$$P(x) = \frac{n!}{(n-x)!x!} p^x q^{n-x}$$

Functions we will use

- $P(x = N) = \text{binompdf}(n, p, N)$
- $P(x \leq N) = \text{binomcdf}(n, p, N)$
- “c” is for “cumulative”

What is the probability that a family with 5 children has exactly 3 boys?

What is the probability that a family with 5 children has 3 or fewer boys?

What is the probability that a family with 5 children has at least 3 boys?

Lesser/Greater

What is the probability that x is...

- equal to N ? $\text{pdf}(N)$
- less than or equal to N ? $\text{cdf}(N)$
- less than N ? $\text{cdf}(N - 1)$
- greater than or equal to N ? $1 - \text{cdf}(N - 1)$
- greater than N ? $1 - \text{cdf}(N)$

Lesser/Greater

- **at most means less than or equal**
- **at least means greater than or equal**
- **no more than means less than or equal**
- **no less than means greater than or equal**
- **up to means less than or equal**

Guessing

- A true/false test has 100 questions. Each question has 2 options, of which one is correct. You guess on every question.
 - What is the probability that you get at least half correct?
 - What is the probability you get no more than 30 correct?
 - What is the probability that you get 60 or more correct?
 - What is the probability that you get exactly 50 correct?
 - What is the probability that you get 100 correct?

Another 5% Rule

- In sampling without replacement, the individuals are not independent, but...
- If the sample size is no more than 5% of the population, then we can treat the individual as independent.

On a college campus of 10,000 students, $\frac{2}{3}$ of the students are female. If 10 students are chosen at random, what is the probability that no more than half of them are female?

Binomial Mean and Standard Deviation

- $\mu = np$
- $\sigma = \sqrt{npq}$

Gregor Mendel

- Gregor Mendel Estimated that the probability that a pea pod with green/yellow genes turns out to be green is 0.75.
- To test his claim, he bred 580 pea pods. Of these, 428 were green.
- If Mendel was correct, would this be unusual (according to the Range Rule of Thumb)?

Gregor Mendel

- Gregor Mendel Estimated that the probability that a pea pod with green/yellow genes turns out to be green is 0.75.
- To test his claim, he bred 580 pea pods. Of these, 428 were green.
- If Mendel was correct, would this be unusual (according to the 5% Rule)?

Racial Discrimination

- 79.1% of the population of Hidalgo County, TX, is of hispanic descent.
- Of 870 people selected for jury duty for a case of burglary against Rodrigo Partida, 339 or 39% were hispanic.
- After conviction, Partida was granted a new trial because of the discrepancy of 39% compared to 79%.
- Statistically, would 339 of 870 be an unusually low number in this case?

Lottery

- In the Illinois Pick 3 Lottery game, players select three digits. There is one winning sequence among the 1000 possibilities.
- If you play this game every day for a year, what is the probability that you win at least once?

Poisson Distribution

Poisson Distribution

- The Poisson distribution is a discrete probability distribution that applies to occurrences of some event over a specified interval.
- The interval can be time, distance, area, volume, or some similar unit.
- The random variable x is the number of occurrences of the event in an interval.

Example Poisson Distributions

- The number of major earthquakes during a year.
- The number of births at a hospital in a year.
- The number of emails received in an hour.
- The number of automobile accidents on a given mile of road.
- The number of bug pieces in a tablespoon of peanut butter.
- The number of dandelions on a square foot of dirt.

More Formulas We Won't Use

For a Poisson Distribution with mean μ

$$\sigma = \sqrt{\mu}$$

$$P(x) = \frac{\mu^x \cdot e^{-\mu}}{x!}$$

Where $e \approx 2.718281828459045$

Functions we will use

Functions we will use

- $P(x = N) = \text{poissonpdf}(n, p, N)$
- $P(x \leq N) = \text{poissoncdf}(n, p, N)$
- “c” is for “cumulative”

Earthquakes

- According to the USGS, there have been 16,500 earthquakes at or above magnitude 6 since 1900.
- What is the probability that there are 125 or fewer earthquakes at or above magnitude 6 in a given year?

Earthquakes

- What would be the usual range for the number of earthquakes at or above magnitude 6 during one year (according to the Range Rule of Thumb)?

Births

- 120 children are born each year at Seward Memorial Hospital.
- SMH has 2 “birthing rooms.”
- What is the probability on any given day that this is adequate?

Bugs in Peanut Butter

- The USDA allows a maximum of 30 “insect parts” in 100g or 3.53oz of peanut butter (twice that for chocolate).
- Suppose that a jar of peanut butter has the maximum allowable number of bug parts.
- If a sandwich is made with one ounce of this peanut butter, then what is the probability that the peanut butter in the sandwich does not contain any bug parts?

Bugs in Peanut Butter

- The USDA allows a maximum of 30 “insect parts” in 100g or 3.53oz of peanut butter (twice that for chocolate).
- Suppose that a jar of peanut butter has the maximum allowable number of bug parts.
- If a sandwich is made with one ounce of this peanut butter, then what is the probability it contains at least 5 bug parts?

Tornadoes

- There were 7236 tornadoes in Texas in a recent span of 54 years (the most of any state in the USA).
- What is the probability that there are 110 or fewer tornadoes in one year in Texas?
- What is the probability that there are more than 150 tornadoes in one year in Texas?

