

Problem III.4.7 Let X be a subscheme of \mathbb{P}_k^2 defined by a single homogeneous equation $f(x_0, x_1, x_2) = 0$ of degree d . (Do not assume f is irreducible.) Assume that $(1, 0, 0)$ is not on X . Then show that X can be covered by the two open affine subsets $U = X \cap \{x_1 \neq 0\}$ and $V = X \cap \{x_2 \neq 0\}$. Now calculate the Čech complex

$$\Gamma(U, \mathcal{O}_X) \oplus \Gamma(V, \mathcal{O}_X) \rightarrow \Gamma(U \cap V, \mathcal{O}_X)$$

explicitly, and thus show that

$$\dim H^0(X, \mathcal{O}_X) = 1, \quad \dim H^1(X, \mathcal{O}_X) = \frac{(d-1)(d-2)}{2}$$

III.5 The Cohomology of Projective Space

Theorem 5.1 Let A be a noetherian ring, let $S = A[x_0, \dots, x_r]$, and let $X = \text{Proj } S$ be the projective space \mathbb{P}_A^r over A . Let $\mathcal{O}_X(1)$ be the twisting sheaf of Serre (II, §5). For any sheaf of \mathcal{O}_X -modules \mathcal{F} , we denote by $\Gamma_*(\mathcal{F})$ the graded S -module $\bigoplus_{n \in \mathbb{Z}} \Gamma(X, \mathcal{F}(n))$ (see II, §5). Then

- (a) the natural map $S \rightarrow \Gamma_*(\mathcal{O}_X) = \bigoplus_{n \in \mathbb{Z}} H^0(X, \mathcal{O}_X(n))$ is an isomorphism of graded S -modules
- (b) $H^i(X, \mathcal{O}_X(n)) = 0$ for $0 < i < r$ and all $n \in \mathbb{Z}$
- (c) $H^r(X, \mathcal{O}_X(-r-1)) \simeq A$
- (d) the natural map

$$H^0(X, \mathcal{O}_X(n)) \times H^r(X, \mathcal{O}_X(-n-r-1)) \rightarrow H^r(X, \mathcal{O}_X(-r-1)) \simeq A$$

is a perfect pairing of finitely generated free A -modules, for each $n \in \mathbb{Z}$.

Problem III.5.3 Let X be a projective scheme of dimension r over a field k . We define the arithmetic genus p_a of X by

$$p_a(X) = (-1)^r (\chi(\mathcal{O}_X) - 1)$$

Note that it depends only on X , not on any projective embedding.

- (a) If X is integral, and k algebraically closed, show that $H^0(X, \mathcal{O}_X) \simeq k$, so that

$$p_a(X) = \sum_{i=0}^{r-1} (-1)^i \dim_k H^{r-i}(X, \mathcal{O}_X)$$

In particular, if X is a curve, we have

$$p_a(X) = \dim_k H^1(X, \mathcal{O}_X)$$

Hint: Use (I, 3.4)

- (b) If X is a closed subvariety of \mathbb{P}_k^r , show that this $p_a(X)$ coincides with the one defined in (I, Ex. 7.2), which apparently depended on the projective embedding.
- (c) If X is a nonsingular projective curve over an algebraically closed field k , show that $p_a(X)$ is in fact a birational invariant. Conclude that a nonsingular plane curve of degree $d \geq 3$ is not rational.