

Vv286 Honors Mathematics IV

Ordinary Differential Equations

Assignment 2

Date Due: 10:00 AM, Thursday, the 29th of September 2016



JOINT INSTITUTE
交大密西根学院

Exercise 2.1. Show that the Riccati differential equation

$$y' + g(x)y + h(x)y^2 = k(x) \quad \text{on an open interval } I \subset \mathbb{R}$$

with $g, h \in C(I)$, $h \in C^1(I)$, $h \neq 0$ on I , can be transformed into the linear differential equation of second order,

$$u'' + \left(g - \frac{h'}{h}\right)u' - khu = 0,$$

using the transformation

$$u(x) = e^{\int h(x)y(x) dx}.$$

(2 Marks)

Exercise 2.2. Show that the equation

$$h(x, y)y' + g(x, y) = 0,$$

if the orthogonal vector field is not a potential field, has an integrating factor of the form $M(x, y) = M(x \cdot y)$ if and only if

$$\frac{h_x - g_y}{xg - hy}$$

is a function of $x \cdot y$ only. Check that this is the case for

$$\left(\frac{x^2}{y} + 3\frac{y}{x}\right)y' + \left(3x + \frac{6}{y}\right) = 0.$$

and then solve the equation.

(2 + 2 Marks)

Exercise 2.3. Find an integrating factor (Euler multiplier) for the equation

$$a_1(x)y' + a_0(x)y = f(x),$$

where a_0, a_1, f are continuous functions on an interval $I \subset \mathbb{R}$. (Try guessing whether the factor might depend only on x or only on y , for example.)


Then solve the equation in general using this factor. Compare with the formula for a particular solution of an inhomogeneous, linear, first-order ordinary differential equation obtained from Duhamel's principle.

(3 Marks)

Exercise 2.4. Show that the solution of Clairaut's equation obtained from a slope parametrization of the integral curve is always the envelope of the straight-line solutions.

(3 Marks)

Exercise 2.5. Light strikes a plane curve C in such a manner that all beams L parallel to the y -axis are reflected to a single point O (see the diagram below). The objective of this exercise is to determine the differential equation for the function $y = f(x)$ describing the curve C .



- i) Show geometrically that $\phi = 2\theta$, $\tan \phi = \frac{x}{y}$ and $\tan(\pi/2 - \theta) = \frac{dy}{dx}$.
- ii) Use the identity $\tan(\pi/2 - x) = \frac{1}{\tan x}$ to show that $\tan \theta = \frac{dx}{dy}$.
- iii) Use the identity $\tan(2x) = 2 \tan x / (1 - \tan^2 x)$ to derive the ODE

$$x \left(\frac{dx}{dy} \right)^2 + 2y \frac{dx}{dy} = x. \quad (*)$$

- iv) Substitute $w = x^2$ in $(*)$ to obtain a differential equation of Clairaut type. Solve this equation, and resubstitute $w = x^2$ to obtain a solution of $(*)$.

- v) What do the above calculations tell you about the types of curves that can be used to focus rays into a single point?

(3 × 1/2 + 1 × 1/2 + 2 + 2 + 2 Marks)

Exercise 2.6. Consider the initial value problem

$$y' = y^2 + x^2, \quad y(0) = 0. \quad (**)$$

- i) Use Picard iteration to find a succession of approximate solutions y_1, y_2, y_3, y_4 , starting from $y_0(x) = 0$. You may use Mathematica to help perform the integrations.
- ii) Use Mathematica to obtain a numerical solution to $(**)$. Plot the numerical solution as well as y_1, y_2, y_3, y_4 in a single graph.

(2 + 3 Marks)