



A Bachelor of Science thesis

Super Learners

and their oracle properties

Jinyang Liu

Supervised by Prof. Thomas Gerds
Co-supervised by Prof. Niels Richard Hansen
Department of Mathematical Sciences
University of Copenhagen, Denmark

Submitted: April 22, 2023

Contents

1	Introduction	3
2	Background	3
3	The discrete super learner, dSL	6
3.1	Risks and selectors	7
3.2	Oracle inequalities	8
3.3	Example: Binary Regression	9
4	The ensemble super learner, eSL	11
5	Simulation results	11
6	Discussion	11

1 Introduction

In the context of prediction, the goal is to estimate the conditional expectation $E(Y \mid X = x)$ for i.i.d. observation pairs $(Y_1, X_1), \dots, (Y_n, X_n)$. Depending on the data structure, a variety of standard statistical models can be employed. For instance, if Y is binary, a parametric model like logistic regression might be suitable. The task of identifying true statistical model \mathcal{P} for which $(Y, X) \sim P \in \mathcal{P}$, is challenging, and perhaps infeasible when we only have a limited amount data. It may therefore be motivating to utilize non-parametric and data-driven regression methods, such as tree-based algorithms like XGBoost or random forests to estimate the conditional mean. However, the underlying assumptions of tree-ensemble methods regarding the data generating process are not explicit, and they may not have probabilistic interpretations. We can nevertheless incorporate these methods as a part of our repertoire, but it is important that we can compare and choose the best method that most effectively accomplishes our goal, for example prediction.

The 'super learner' is the answer to how we can effectively select the 'best' learner (method or algorithm) among the learners that we have in our library of learners. The cross-validation selector, which evaluates learners on their cross-validated (empirical) risk and chooses the one with the lowest risk, is asymptotically equivalent to the oracle selector. The oracle selector finds the learner with the lowest true risk – the risk obtained by knowing the true distribution.

The discrete super learner is obtained by applying the cross-validation selector on our data. The asymptotic result shows that the cross-validation selector will select the same learner as the oracle selector when the number of observations goes to ∞ . The discrete super learner is not a fixed learner from our library, but rather depends on the available data. It represents the learner chosen by the cross-validation selector, which can vary depending on the amount of data at hand.

We first present the general theory and our goal, which is to estimate the conditional expectation $E(Y \mid X = x)$ for Y, X being some outcome-covariate pair. More specifically, we focus on the case where we regress on a binary outcome $Y \in \{0, 1\}$. The conditional expectation of Y given X exactly becomes the conditional probability $P(Y = 1 \mid X = x)$.

2 Background

Our setup closely models what is described in [VDL06] and [LD03]. Unless specified otherwise, our setup is as follows: Let statistical model \mathcal{P} be given on the measurable space $(\mathcal{O}, \mathcal{A})$ where $\mathcal{O} = \{0, 1\} \times \mathcal{X}$ is our sample space. We will consider the parameter set $\Theta = \{\theta \mid \theta : \mathcal{X} \rightarrow [0, 1]\}$, which represents the set of regression functions that map from our covariates to the probability interval. We will consider the quadratic loss function

Definition 1 (Quadratic loss). Let \mathcal{O} be our sample space, and Θ be the set of regression functions. Then the quadratic loss or L^2 -loss, $L : \mathcal{O} \times \Theta \rightarrow [0, \infty)$, for an observation $o \in \mathcal{O}$ and a regression function $\theta \in \Theta$ is defined as

$$L(o, \theta) = L((y, x), \theta) = (y - \theta(x))^2.$$

We aim to find the optimal parameter value $\theta^* \in \Theta$ that minimizes the L^2 -risk function $R : \Theta \rightarrow \mathbb{R}$

$$R(\theta) = \int L(o, \theta) dP(o) = EL(O_1, \theta),$$

but this task is challenging as it requires us to know the data-generating process, P , which we do not have access to. It can be shown that the minimum risk is achieved by the conditional probability $x \mapsto P(Y = 1 \mid X = x)$.

Theorem 2. *Let $(\mathcal{O}, \mathcal{A}, P)$ be a probability space for some probability measure $P \in \mathcal{P}$. Let Θ be the set of regression functions of the form $\theta : \mathcal{X} \rightarrow [0, 1]$. Let the loss function be the L^2 -loss $L(o, \theta) = (y - \theta(x))^2$, then for the optimum θ^* defined as*

$$\theta^* := \arg \min_{\theta \in \Theta} R(\theta) = \arg \min_{\theta \in \Theta} \int L(o, \theta) dP(o),$$

it holds for an observation $O = (Y, X) \sim P$ that

$$\theta^*(x) = E(Y \mid X = x)$$

Proof. See [Gyö+02][ch. 1] □

It follows immediately that if Y is binary, then $E(Y \mid X = x) = P(Y = 1 \mid X = x)$. In the case where we do not have access to the data-generating process, we would instead want to provide an estimate $\hat{\theta}$ of θ^* as a function of the data that we have observed. Let $O_1 = (Y_1, X_1), \dots, O_n = (Y_n, X_n) \in \mathcal{O}$ be i.i.d. observations distributed according to some $P \in \mathcal{P}$. Denote our data as $D_n = (O_1, \dots, O_n)$. An estimate is the outcome of applying an estimator to the data

$$D_n \mapsto \hat{\theta}(D_n) \in \Theta,$$

The estimate is a regression function, that is

$$\mathcal{X} \ni x \mapsto \hat{\theta}(D_n)(x) \in [0, 1].$$

In the context of super learners we will refer the estimators as learners, in the sense that they learn from the data, the resulting regression function is then the fit of the learner to the data. We formalize these notions in the subsequent section.

Example 1 (Parametric statistical model). In the case where we have n -i.i.d. observations distributed according to some data-generating process $P \in \mathcal{P}$, we have that each $O_i = (Y_i, X_i)$, and X_i is a stochastic variable. The distribution P factorizes essentially into two parts, the conditional distribution of Y given X and the background distribution of X , so $P = P_{Y|X=x} \cdot P_X$. In this setup we are doing estimation with random design [Gyö+02].

We can formalize the setup as follows: $O = (Y, X) \sim P$, if Y is $\mathcal{B}(\mathbb{R})$ - $\mathcal{B}(\mathbb{R})$ measurable and X is \mathcal{F} - $\mathcal{B}(\mathbb{R})$ measurable for some sigma-algebra \mathcal{F} on \mathcal{X} , then a **generalized regression model** could be considered as parametrized family of distributions, $\mathcal{P} = \{P_\theta \mid \theta \in \Theta\}$, given that Θ is finite-dimensional.

We can parametrize the conditional probability distributions for Y_1 given $X_1 = x$ as $\mathcal{Q} = \{Q_{\theta(x)} \mid \theta \in \Theta\}$ such that $Q_{\theta(x)}$ is a valid probability distribution on $\mathcal{B}(\mathbb{R})$ for each $x \in \mathcal{X}$ and $\theta \in \Theta$. For a given $P_\theta \in \mathcal{P}$ there will exist a $Q_\theta \in \mathcal{Q}$ such that

$$P_\theta(Y \in A \mid X = x) = Q_{\theta(x)}(A) \quad \text{for all } A \in \mathcal{B}(\mathbb{R}).$$

If we assume that X_1 is distributed according to some H_0 on \mathcal{X} , then the distribution P_θ over our observations (the joint over Y and X) will be

$$P_\theta(X \in A, Y \in B) = \int_A Q_{\theta(x)}(B) dH_0(x)$$

for every $A \in \mathcal{F}$ and $B \in \mathcal{B}(\mathbb{R})$.

Example 2 (Logistic regression model). Let $O_1 = (Y_1, X_1), \dots, O_n = (Y_n, X_n) \in \mathcal{O} = \{0, 1\} \times \mathcal{X}$ be i.i.d. observations from some distribution $P_{\theta_0} \in \mathcal{P}$, where Y_i is binary and $\mathcal{X} \subseteq \mathbb{R}^k$. We would like to estimate the parameter function $\theta_0 \in \Theta$

$$\theta_0(x) = E(Y_1 \mid X_1 = x) = P_{\theta_0}(Y_1 = 1 \mid X_1 = x),$$

In logistic regression we assume that $\Theta = \{x \mapsto \text{expit}(\beta x) \mid \beta \in \mathbb{R}^k\}$, so $\theta_0(x) = \text{expit}(\beta_0 x)$, then the goal becomes to estimate the k -dimensional parameter β_0 , in this case the \mathbb{R}^k parameter β_0 completely determines θ_0 , so Θ is also k -dimensional. The conditional distributions of Y_1 given $X_1 = x$ are Bernoulli distributions and can be parametrized as $\mathcal{Q} = \{\text{Ber}(\text{expit}(\beta x)) \mid \beta \in \mathbb{R}^k\}$. Now from example 1 we know that the statistical model, \mathcal{P} , can be parametrized through β , in particular we have

$$\begin{aligned} P_\beta(Y_1 = 1, X_1 \in A) &= \int_A Q_{\theta(x)}(\{1\}) dH_0(x) \\ &= \int_A \text{expit}(\beta x) dH_0(x) \end{aligned}$$

If H_0 has density f w.r.t. Lebesgue measure, we can write

$$P_\beta(Y_1 = 1, X_1 \in A) = \int_A \text{expit}(\beta x) f(x) dm(x)$$

We will now turn our attention to statistical estimators. Statistical literature commonly write that an estimator is stochastic variable taking values in our parameter space $\hat{\theta} \in \Theta$. An estimator is achieved by considering i.i.d. observations $O_1, \dots, O_n \in \mathcal{O}$ distributed

according to some measure P from some statistical model \mathcal{P} . We leave the model unspecified as it can be both parametric or nonparametric. Now let $h : \mathcal{O}^n \rightarrow \Theta$ be a measurable map, an estimator created from h is the random variable $T = h(O_1, \dots, O_n)$. For $\Theta \subseteq \mathbb{R}^k$ the canonical σ -algebra on Θ is the Borel algebra, but when the parameter set is a set of functions, the σ -algebra can only be chosen after careful consideration of constraints on Θ .

3 The discrete super learner, dSL

In the following section we introduce the terminology “estimator algorithm” which corresponds to the measurable map h from our finite sample observation space to our parameter space.

Definition 3 (Estimator algorithm h). An estimator algorithm is a measurable map $h : \mathcal{O}^n \rightarrow \Theta$ for $n \in \mathbb{N}$.

Definition 4 (Statistical Estimator $\hat{\theta}$). Let $O_1, \dots, O_n \in \mathcal{O}$ be i.i.d. observations distributed according to some $P \in \mathcal{P}$ for a statistical model \mathcal{P} on \mathcal{O} . Let $h : \mathcal{O}^n \rightarrow \Theta$ be an estimator algorithm. An estimator is the random variable $\hat{\theta} = h(O_1, \dots, O_n) \in \Theta$.

There is a one-to-one correspondence between the tuples of i.i.d. observations $(O_1, \dots, O_n) \in \mathcal{O}^n$ and the empirical measures over n observations on $(\mathcal{O}, \mathcal{A})$ defined as

$$P_n(A) = \frac{1}{n} \sum_{i=1}^n \delta_{O_i}(A) \quad \text{for } A \in \mathcal{A}.$$

Note that the empirical measure is a random variable. Thus, we can write $h(P_n)$ as an alternative representation of the estimator $h(O_1, \dots, O_n)$, by adjusting the notation without introducing ambiguity.

Example 3 (Prediction algorithm). Consider the setup from example 2, where we have i.i.d. observations $O_1 = (Y_1, X_1), \dots, O_n = (Y_n, X_n)$ such that $Y_i \in \{0, 1\}$ and $\mathcal{X} \in \mathbb{R}^k$ and our goal is to estimate the probability $\theta(x) = P_\theta(Y_1 = 1 \mid X_1 = x)$...

We would now like to consider the scenario where we have a library (set) of learner algorithms, h_1, \dots, h_p . From these algorithms, we can define the set of learners $\{\hat{\theta}_q = h_q(P_n) \mid 1 \leq q \leq p\}$, where our goal is to find $\hat{\theta}_{\hat{q}}(P_n)$, which denotes the learner that minimizes R and \hat{q} may depend on the observations.

In order to find \hat{q} we have to proceed via cross validation. In cross validation, we randomly split our data into a training set and a test set. Let the random binary vector $S = (S_1, \dots, S_n) \in \{0, 1\}^n$ be independent of X_1, \dots, X_n such that $S_i = 0$ indicates that X_i should be in the training set and $S_i = 1$ indicates that X_i belongs to the test set. We can define the empirical distributions over these two subsets, $P_{n,S}^0$ and $P_{n,S}^1$ as

$$P_{n,S}^0 = \frac{1}{n_0} \sum_{i:S_i=0} \delta_{X_i}$$

$$P_{n,S}^1 = \frac{1}{1 - n_0} \sum_{i:S_i=1} \delta_{X_i}$$

Where n_0 would be the number of S_i 's that are marked 0.

Example 4 (Random splits). For $n = 9$ observations one could for example define the distribution of the random vector S as

$$\begin{aligned} P(S = (0, 0, 0, 0, 0, 0, 1, 1, 1)) &= \frac{1}{3} \\ P(S = (0, 0, 0, 1, 1, 1, 0, 0, 0)) &= \frac{1}{3} \\ P(S = (1, 1, 1, 0, 0, 0, 0, 0, 0)) &= \frac{1}{3}, \end{aligned}$$

i.e. 3-fold cross-validation.

In general for n observations we have 2^n ways of choosing which observations should be in the training set and in the validation set. It might not be desirable to define the discrete probabilities for S over $\{0, 1\}^n$ simply as $\frac{1}{2^n}$ for each possible combination of training/validation data, since that would also include the combination where $n_1 = 0$. To ensure that we always have $n_1 > 0$, then let n_1 be given, then we see that there are $\binom{n}{n_1}$ ways of choosing both the validation and training set. We can therefore define the distribution of S as

$$P(S = s) = \binom{n}{n_1}^{-1} \quad \text{for each } s \in \{0, 1\}^n \text{ where } \sum_i s_i = n_1$$

3.1 Risks and selectors

Definition 5 (True risk of q 'th learner averaged over splits). Given the data $O_1, \dots, O_n \in \mathcal{O}$ and a set of learners $\{\theta_q(P_{n,S}^0) \mid 1 \leq q \leq k\}, k \in \mathbb{N}$ applied to our training data $P_{n,S}^0$. The risks of these learners averaged over some split-variable S is given as a function of q

$$q \mapsto E_S \int L(o, \theta_q(P_{n,S}^0)) dP(o) = E_S R(\theta_q(P_{n,S}^0))$$

Where P is the true distribution for our data X .

Definition 6 (Oracle selector). The oracle selector is a function $\tilde{q} : \mathcal{O}^n \rightarrow \{1, \dots, k\}$ which finds the learner that minimizes the true risk given our data $O_1, \dots, O_n \in \mathcal{O}$.

$$\tilde{q}(O_1, \dots, O_n) = \arg \min_{1 \leq q \leq k} E_S R(\theta_q(P_{n,S}^0))$$

Where $P_{n,S}^0$ is the empirical distribution over the training set of O_1, \dots, O_n as specified by some split-variable S .

In similar manner to the above the definitions, we can define the cross-validation risk and the cross-validation selector for our learners

Definition 7 (Cross-validation risk of q 'th learner averaged over splits). Given the data $O_1, \dots, O_n \in \mathcal{O}$ and a set of learners $\{\theta_q(P_{n,S}^0) \mid 1 \leq q \leq k\}, k \in \mathbb{N}$. The cross-validation risks of these learners averaged over some split-variable S is given as a function of q

$$q \mapsto E_S \int L(o, \theta_q(P_{n,S}^0)) dP_{n,S}^1(o) = E_S \hat{R}(\theta_q(P_{n,S}^0))$$

Where $P_{n,S}^1$ is the empirical distribution over the validation of O_1, \dots, O_n . We write \hat{R} for the empirical risk over the validation set.

Definition 8 (Cross-validation selector). The cross-validation selector is a function $\hat{q} : \mathcal{O}^n \rightarrow \{1, \dots, k\}$ which finds the learner that minimizes the cross-validation risk given our data $O_1, \dots, O_n \in \mathcal{O}$.

$$\hat{q}(O_1, \dots, O_n) = \arg \min_{1 \leq q \leq k} E_S \hat{R}(\theta_q(P_{n,S}^0))$$

Where \hat{R} is the empirical risk over the validation set and $P_{n,s}^0$ is the empirical distribution over the training set of O_1, \dots, O_n as specified by some split-variable S .

We are interested in the risk difference between the cross-validation selector and the oracle selector, we remark that the optimal risk is attained at the true value θ_0

$$R(\theta_0) = \int L(o, \theta_0) dP(o),$$

and clearly it is the case that $R(\theta_0) \leq R(\theta)$ for any learner θ of θ_0 . Given a set of learners we define the centered conditional risk as the difference

$$\begin{aligned} \Delta_S(\theta_{\hat{q}}, \theta_0) &= R(\theta_{\hat{q}}(P_{n,S}^0)) - R(\theta_0) \\ &= E_S \int L(o, \theta_{\hat{q}}(P_{n,S}^0)) - L(o, \theta_0) dP(o) \end{aligned}$$

3.2 Oracle inequalities

We introduce the notation Pf for the integral $\int f dP$ of an integrable function f with respect to P . Additionally, if P_n represents the empirical measure of O_1, \dots, O_n , we denote the empirical process indexed over an appropriate class of functions \mathcal{F} as $G_n f = \sqrt{n}(P_n f - Pf)$. Furthermore, we extend this notation to $G_{n,S}^i f = \sqrt{n}(P_{n,S}^i - Pf)$ for the empirical processes that correspond to applying the empirical measure over either the training sample or validation sample.

In the following results we assume that a proper loss function $L : \mathcal{O} \times \Theta \rightarrow \mathbb{R}$ has been given.

Lemma 9 (Lemma 2.1 in [VDL06]). *Let G_n be the empirical process of an i.i.d. sample of size n from the distribution P . For $\delta > 0$ it holds that*

$$\begin{aligned} E_S \int L(o, \theta_{\hat{q}}(P_{n,S}^0)) dP(o) &\leq (1 + 2\delta) E_S \int L(o, \theta_{\hat{q}}(P_{n,S}^0)) dP(o) \\ &\quad + \frac{1}{\sqrt{n_1}} E_S \max_{1 \leq q \leq k} \int L(o, \theta_q(P_{n,S}^0)) d((1 + \delta)G_{n,S}^1 - \delta\sqrt{n_1}P)(o) \\ &\quad + \frac{1}{\sqrt{n_1}} E_S \max_{1 \leq q \leq k} \int -L(o, \theta_q(P_{n,S}^0)) d((1 + \delta)G_{n,S}^1 + \delta\sqrt{n_1}P)(o) \end{aligned}$$

Proof. See appendix □

Lemma 10 (Lemma 2.2 in [VDL06]). *Let G_n be the empirical process of an i.i.d. sample of size n from the distribution P and assume that $Pf \geq 0$ for every $f \in \mathcal{F}$. Then, for any Bernstein pairs $(M(f), v(f))$ and for any $\delta > 0$ and $1 \leq p \leq 2$,*

$$E \max_{f \in \mathcal{F}} (G_n - \delta\sqrt{n}P)f \leq \frac{8}{n^{1/p-1/2}} \log(1 + \#\mathcal{F}) \max_{f \in \mathcal{F}} \left[\frac{M(f)}{n^{1-1/p}} + \left(\frac{v(f)}{(\delta Pf)^{2-p}} \right)^{1/p} \right].$$

The same upper bound is valid for $E \max_{f \in \mathcal{F}} (G_n + \delta\sqrt{n}P)(-f)$

Theorem 11 (Finite Sample Result: Theorem 2.3 in [VDL06]). *For $\theta \in \Theta$ let $(M(\theta), v(\theta))$ be a Bernstein pair for the function $o \mapsto L(o, \theta)$ and assume that $R(\theta) = \int L(o, \theta) dP(o) \geq 0$ for every $\theta \in \Theta$. Then for $\delta > 0$ and $1 \leq p \leq 2$ it holds that*

$$ER(\theta_{\hat{q}}(P_{n,S}^0)) \leq (1 + 2\delta)ER(\theta_{\hat{q}}(P_{n,S}^0)) + (1 + \delta)E \left(\frac{16}{n_1^{1/p}} \log(1 + k) \sup_{\theta \in \Theta} \left[\frac{M(\theta)}{n_1^{1-1/p}} + \left(\frac{v(\theta)}{R(\theta)^{2-p}} \right)^{1/p} \left(\frac{1 + \delta}{\delta} \right)^{2/p-1} \right] \right),$$

where k is the number of learners in our library $\{\theta_q(P_{n,S}^0) \mid 1 \leq q \leq k\}$.

Proof. We will apply lemma 10 to the second and third terms on the left hand side of the inequality in lemma 9. Let $\mathcal{F} = \{o \mapsto L(o, \theta_q(P_{n,S}^0)) \mid 1 \leq q \leq k\}$, be the collection of functions obtained by applying the loss L to each learner in our library of k learners. Note that $\mathcal{F} \subseteq \{o \mapsto L(o, \theta) \mid \theta \in \Theta\}$, and since $R(\theta) \geq 0$ for every $\theta \in \Theta$ it follows that $Pf \geq 0$ for every $f \in \mathcal{F}$.

For the second term we have

$$\begin{aligned} & \frac{1}{\sqrt{n_1}} E_S \max_{1 \leq q \leq k} \int L(o, \theta_q(P_{n,S}^0)) d((1 + \delta)G_{n,S}^1 - \delta\sqrt{n_1}P)(o) \\ &= \frac{1 + \delta}{\sqrt{n_1}} E_S \max_{1 \leq q \leq k} \int L(o, \theta_q(P_{n,S}^0)) d(G_{n,S}^1 - \frac{\delta}{1 + \delta}\sqrt{n_1}P)(o), \end{aligned}$$

applying lemma 10 to the expression above yields

$$\begin{aligned} & \frac{1 + \delta}{\sqrt{n_1}} E_S \max_{1 \leq q \leq k} \int L(o, \theta_q(P_{n,S}^0)) d(G_{n,S}^1 - \frac{\delta}{1 + \delta}\sqrt{n_1}P)(o) \\ & \leq \frac{1 + \delta}{\sqrt{n_1}} \left(\frac{8}{n_1^{1/p-1/2}} \log(1 + k) \max_{1 \leq q \leq k} \left[\frac{M(\theta_q(P_{n,S}^0))}{n_1^{1-1/p}} + \left(\frac{v(\theta_q(P_{n,S}^0))}{(\frac{\delta}{1+\delta})^{2-p} R(\theta_q(P_{n,S}^0))^{2-p}} \right)^{1/p} \right] \right) \\ & \leq \frac{1 + \delta}{\sqrt{n_1}} \left(\frac{8}{n_1^{1/p-1/2}} \log(1 + k) \sup_{\theta \in \Theta} \left[\frac{M(\theta)}{n_1^{1-1/p}} + \left(\frac{v(\theta)}{(\frac{\delta}{1+\delta})^{2-p} R(\theta)^{2-p}} \right)^{1/p} \right] \right) \\ & = (1 + \delta) \frac{8}{n_1^{1/p}} \log(1 + k) \sup_{\theta \in \Theta} \left[\frac{M(\theta)}{n_1^{1-1/p}} + \left(\frac{v(\theta)}{R(\theta)^{2-p}} \right)^{1/p} \left(\frac{1 + \delta}{\delta} \right)^{2/p-1} \right] \end{aligned}$$

Where for the third inequality we take the sup over Θ . We can also bound the third term with the same expression above. It is now immediate from lemma 9 that

$$\begin{aligned} E_S \int L(o, \theta_{\hat{q}}(P_{n,S}^0)) dP(o) & \leq (1 + 2\delta) E_S \int L(o, \theta_{\hat{q}}(P_{n,S}^0)) dP(o) \\ & \quad + 2 \cdot (1 + \delta) \frac{8}{n_1^{1/p}} \log(1 + k) \sup_{\theta \in \Theta} \left[\frac{M(\theta)}{n_1^{1-1/p}} + \left(\frac{v(\theta)}{R(\theta)^{2-p}} \right)^{1/p} \left(\frac{1 + \delta}{\delta} \right)^{2/p-1} \right], \end{aligned}$$

which was exactly what we set out to prove. \square

3.3 Example: Binary Regression

Consider the case where we have i.i.d. observations $O_1 = (Y_1, X_1), \dots, O_n = (Y_n, X_n)$ such that $Y_i \in \{0, 1\}$ and $X \in \mathbb{R}^d$ distributed according some $P \in \mathcal{P}$. We would like to estimate the conditional expectation $\theta_0(x) = E(Y \mid X = x) = P(Y = 1 \mid X = x)$. Let

$\Theta = \{\theta \mid \theta : \mathcal{X} \rightarrow [0, 1] \text{ measurable}\}$ and choose the quadratic loss function $L((Y, X), \theta) = (Y - \theta(X))^2$.

We observe that the quadratic loss is bounded by 1 for all choices of $\theta \in \Theta$ and $O \in \mathcal{O}$. It is stated in [VDL06, p. 7] that $M(\theta) = 1$ and $v(\theta) = \frac{3}{2} \int L(o, \theta)^2 dP(o)$ is a valid Bernstein pair for the function $o \mapsto L(o, \theta)$. It is also clear that $R(\theta) = \int L(o, \theta) dP(o) \geq 0$ since the loss function is positive. If we plug these numbers in theorem 11, then by using $p = 1$ and

$$ER(\theta_{\hat{q}}(P_{n,S}^0)) \leq (1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0)) + (1 + \delta)E \left(\frac{16}{n_1} \log(1 + k) \sup_{\theta \in \Theta} \left[M(\theta) + \frac{v(\theta)}{R(\theta)} \frac{1 + \delta}{\delta} \right] \right)$$

In the equation provided, we observe that we can manipulate the following variables: sample size n , validation set size n_1 , parameter $\delta > 0$, and the number of learners k . Assuming k remains constant, the validation set size n_1 could be either stochastic, depending on the split variable S , or fixed as a constant, as illustrated in example 4. For instance, we can set $n_1 = n/2$. By establishing a fixed value for n_1 , we can drop the expectation in the second term.

The supremum on the left side of the equation might increase significantly because $R(\theta)$ could be very small. However, by carefully selecting the value of $v(\theta)$, we can avoid the fraction from growing too large. Note that for any $\theta \in \Theta$:

$$\frac{v(\theta)}{R(\theta)} = \frac{3 \int L(o, \theta)^2 dP(o)}{2 \int L(o, \theta) dP(o)} \leq \frac{3 \int L(o, \theta) \cdot 1 dP(o)}{2 \int L(o, \theta) dP(o)} = \frac{3}{2},$$

by using $0 \leq L(o, \theta) \leq 1$ almost surely. Since $M(\theta)$ is constant for all $\theta \in \Theta$, it is possible to drop the supremum. Combining all the information above we obtain

$$\begin{aligned} ER(\theta_{\hat{q}}(P_{n,S}^0)) &\leq (1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0)) + (1 + \delta) \frac{16}{n_1} \log(1 + k) \left[1 + \frac{3}{2} \frac{1 + \delta}{\delta} \right] \\ &= (1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0)) + \log(1 + k) \frac{3 + 5\delta}{2\delta} (1 + \delta) \frac{16}{n_1} \\ &= (1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0)) + \log(1 + k) \frac{3 + 8\delta + 5\delta^2}{2\delta} \frac{16}{n_1} \\ &= (1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0)) + \log(1 + k) \left(\frac{3}{2\delta} + 4 + \frac{5}{2}\delta \right) \frac{16}{n_1}, \end{aligned}$$

We can now adjust for the precision in our bound by choosing δ and n . Note that a small delta will mean that the first term $(1 + 2\delta)ER(\theta_{\bar{q}}(P_{n,S}^0))$ will become smaller, but this is at the expense that the remainder term becomes larger due to the $\frac{1+\delta}{\delta}$ fraction at the end. By choosing n to be large, we can partially compensate for a smaller delta.

We might, therefore, for each n , choose the δ_n that minimizes the left-hand side for the given n . By substituting $n_1 = n/2$ into the the left hand side expression and then expanding it we obtain

$$ER(\theta_{\hat{q}}(P_{n,S}^0)) + 2\delta ER(\theta_{\bar{q}}(P_{n,S}^0)) + \frac{3}{2\delta} \log(1 + k) \frac{32}{n} + 4 \log(1 + k) \frac{32}{n} + \frac{5\delta}{2} \log(1 + k) \frac{32}{n},$$

we observe that when n is fixed, two terms remain constant, specifically the first and fourth terms, as they do not depend on δ . The optimal δ_n can be determined as follows:

$$\begin{aligned} \delta_n &= \arg \min_{\delta} \frac{1}{\delta} \cdot \frac{3 \cdot 32}{2n} \log(1 + k) + \delta \cdot \left(2ER(\theta_{\bar{q}}(P_{n,S}^0)) + \frac{5 \cdot 32}{2n} \log(1 + k) \right) \\ &= \arg \min_{\delta} \frac{1}{\delta} \cdot \frac{48}{n} \log(1 + k) + \delta \cdot \left(2ER(\theta_{\bar{q}}(P_{n,S}^0)) + \frac{80}{n} \log(1 + k) \right) \\ &= \arg \min_{\delta} \frac{1}{\delta} a(n) + \delta b(n), \end{aligned}$$

Essentially, solving for the minimum is a convex optimization problem, with the terms $a(n) = \frac{48}{n} \log(1+k)$ and $b(n) = 2ER(\theta_{\hat{q}}(P_{n,S}^0)) + \frac{80}{n} \log(1+k)$ remaining constant with respect to n . By differentiating the expression above and setting it equal to zero we obtain

$$0 = \left(\frac{1}{\delta} a(n) + \delta b(n) \right)' = -\frac{1}{\delta^2} a(n) + b(n),$$

and so we obtain the optimum by isolating δ

$$\delta_n = \sqrt{\frac{a(n)}{b(n)}},$$

4 The ensemble super learner, eSL

5 Simulation results

6 Discussion

Pellentesque tincidunt sodales risus, vulputate iaculis odio dictum vitae. Ut ligula tortor, porta a consequat ac, commodo non risus. Nullam sagittis luctus pretium. Integer vel nibh at justo convallis imperdiet sit amet ut lorem. Sed in gravida turpis. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Sed in massa vitae ligula pellentesque feugiat vitae in risus. Cras iaculis tempus mi, sit amet viverra nulla viverra pellentesque.

References

- [Gyö+02] László Györfi et al. *A distribution-free theory of nonparametric regression*. Vol. 1. Springer, 2002.
- [LD03] Mark Laan and Sandrine Dudoit. “Unified Cross-Validation Methodology For Selection Among Estimators and a General Cross-Validated Adaptive Epsilon-Net Estimator: Finite Sample Oracle Inequalities and Examples”. In: *UC Berkeley Division of Biostatistics Working Paper Series* (Jan. 2003).
- [VDL06] Aad W. van der Vaart, Sandrine Dudoit, and Mark J. van der Laan. In: *Statistics & Decisions* 24.3 (2006), pp. 351–371. DOI: doi:10.1524/stnd.2006.24.3.351. URL: <https://doi.org/10.1524/stnd.2006.24.3.351>.