

Nonnegative Ricci Curvature, Nilpotency, and Asymptotic Geometry

Jiayin Pan

UC Santa Cruz

SLMath, Recent Progress on
Geometric Analysis and Riemannian Geometry



SIM NS
FOUNDATION



SIMONS LAUFER
MATHEMATICAL
SCIENCES INSTITUTE



Curvature

Sectional curvature:

Let $p \in M$ and Σ be a plane in $T_p M$ through 0. Roughly speaking, $\sec_p(\Sigma) \geq 0$ means geodesics with initial data in Σ cannot (locally) spread wider than the ones in \mathbb{R}^2 .

$\sec \geq 0$ means $\sec_p(\Sigma) \geq 0$ for all p and all Σ .

Curvature

Sectional curvature:

Let $p \in M$ and Σ be a plane in $T_p M$ through 0. Roughly speaking, $\sec_p(\Sigma) \geq 0$ means geodesics with initial data in Σ cannot (locally) spread wider than the ones in \mathbb{R}^2 .

$\sec \geq 0$ means $\sec_p(\Sigma) \geq 0$ for all p and all Σ .

Ricci curvature:

Let $p \in M^n$ and v be a unit vector in $T_p M$. Extend v to an orthonormal basis $\{v, E_2, \dots, E_n\}$ at p . $\text{Ric}_p(v, v) = \sum_i \sec(v \wedge E_i)$.

$\text{Ric} \geq 0$ means $\text{Ric}_p(v, v) \geq 0$ for all p and all v .

Curvature

Sectional curvature:

Let $p \in M$ and Σ be a plane in $T_p M$ through 0. Roughly speaking, $\sec_p(\Sigma) \geq 0$ means geodesics with initial data in Σ cannot (locally) spread wider than the ones in \mathbb{R}^2 .

$\sec \geq 0$ means $\sec_p(\Sigma) \geq 0$ for all p and all Σ .

Ricci curvature:

Let $p \in M^n$ and v be a unit vector in $T_p M$. Extend v to an orthonormal basis $\{v, E_2, \dots, E_n\}$ at p . $\text{Ric}_p(v, v) = \sum_i \sec(v \wedge E_i)$.
 $\text{Ric} \geq 0$ means $\text{Ric}_p(v, v) \geq 0$ for all p and all v .

$\text{Ric} \geq 0$ controls the volume growth:

Theorem (Bishop 1964, Gromov 1980)

$$\text{vol}(B_r(p)) \leq \omega_n r^n, \quad \frac{\text{vol}(B_R(p))}{\text{vol}(B_r(p))} \leq \frac{R^n}{r^n}, \text{ where } 0 < r < R.$$

Nonnegative curvature and virtual abelianness

Theorem (Bieberbach 1912)

The fundamental group of a closed flat n -manifold contains a subgroup \mathbb{Z}^n with index $\leq C(n)$.

Nonnegative curvature and virtual abelianness

Theorem (Bieberbach 1912)

The fundamental group of a closed flat n -manifold contains a subgroup \mathbb{Z}^n with index $\leq C(n)$.

Theorem (Cheeger–Gromoll 1972)

The fundamental group of a closed manifold with $\text{Ric} \geq 0$ is virtually abelian ($\mathbb{Z}^k \triangleleft \pi_1$ of finite index, where $0 \leq k \leq n$).

Nonnegative curvature and virtual abelianness

Theorem (Bieberbach 1912)

The fundamental group of a closed flat n -manifold contains a subgroup \mathbb{Z}^n with index $\leq C(n)$.

Theorem (Cheeger–Gromoll 1972)

The fundamental group of a closed manifold with $\text{Ric} \geq 0$ is virtually abelian ($\mathbb{Z}^k \triangleleft \pi_1$ of finite index, where $0 \leq k \leq n$).

Theorem (Cheeger–Gromoll 1972)

Let M be an open (complete and non-compact) manifold of $\text{sec} \geq 0$. Then M contains a closed totally convex submanifold S (soul) s.t. M is diffeo to the normal bundle over S .
 $\Rightarrow \pi_1(M) = \pi_1(S)$ is virtually abelian.

Abelian/nilpotent groups

How about open manifolds of $\text{Ric} \geq 0$?

(Wei 1988) Examples of open M with $\text{Ric} > 0$ and torsion-free nilpotent π_1 . (In particular, not virtually abelian.)

(Wilking 2000) Every finitely generated virtually nilpotent group can be realized as π_1 of some open M with $\text{Ric} > 0$.

Abelian: $[G, G] = \{e\}$.

Nilpotent: $[G, G] = G^1, [G, G^i] = G^{i+1}$; eventually, $G^k = \{e\}$.

Abelian/nilpotent groups

How about open manifolds of $\text{Ric} \geq 0$?

(Wei 1988) Examples of open M with $\text{Ric} > 0$ and torsion-free nilpotent π_1 . (In particular, not virtually abelian.)

(Willing 2000) Every finitely generated virtually nilpotent group can be realized as π_1 of some open M with $\text{Ric} > 0$.

Abelian: $[G, G] = \{e\}$.

Nilpotent: $[G, G] = G^1, [G, G^i] = G^{i+1}$; eventually, $G^k = \{e\}$.

Example: Discrete Heisenberg 3-group $H^3(\mathbb{Z})$

$$N = \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in \mathbb{Z} \right\}.$$

$$N^1 = [N, N] = \left\{ \begin{pmatrix} 1 & 0 & c \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid c \in \mathbb{Z} \right\}.$$

$$N^2 = [N, N^1] = \{e\}.$$

Curvature and virtual abelianness/nilpotency

Let M be an open n -manifold of $\text{Ric} \geq 0$.

Theorem

Any finitely generated subgroup of $\pi_1(M)$

(1) has polynomial growth with degree $\leq n$ (Milnor 1968),

(2) is virtually nilpotent (Gromov 1981).

Curvature and virtual abelianness/nilpotency

Let M be an open n -manifold of $\text{Ric} \geq 0$.

Theorem

Any finitely generated subgroup of $\pi_1(M)$

(1) has polynomial growth with degree $\leq n$ (Milnor 1968),

(2) is virtually nilpotent (Gromov 1981).

Theorem (Kapovitch–Wilking 2011)

$\pi_1(M)$ is $C(n)$ -nilpotent (i.e., index $\leq C(n)$).

Curvature and virtual abelianness/nilpotency

Let M be an open n -manifold of $\text{Ric} \geq 0$.

Theorem

Any finitely generated subgroup of $\pi_1(M)$

(1) has polynomial growth with degree $\leq n$ (Milnor 1968),

(2) is virtually nilpotent (Gromov 1981).

Theorem (Kapovitch–Wilking 2011)

$\pi_1(M)$ is $C(n)$ -nilpotent (i.e., index $\leq C(n)$).

Counterexample to Milnor Conjecture (Brue–Naber–Semola 2023)

M^n ($n \geq 6$) with $\text{Ric} \geq 0$ and $\pi_1(M) = \mathbb{Q}/\mathbb{Z}$.

Curvature and virtual abelianness/nilpotency

Let M be an open n -manifold of $\text{Ric} \geq 0$.

Theorem

Any finitely generated subgroup of $\pi_1(M)$

- (1) has polynomial growth with degree $\leq n$ (Milnor 1968),
- (2) is virtually nilpotent (Gromov 1981).

Theorem (Kapovitch–Wilking 2011)

$\pi_1(M)$ is $C(n)$ -nilpotent (i.e., index $\leq C(n)$).

Counterexample to Milnor Conjecture (Brue–Naber–Semola 2023)

M^n ($n \geq 6$) with $\text{Ric} \geq 0$ and $\pi_1(M) = \mathbb{Q}/\mathbb{Z}$.

Question: Relationships between the virtual abelianness/nilpotency of $\pi_1(M)$ and the geometry of M ?

Escape phenomenon

$\sec \geq 0 \Rightarrow$ (by soul structure and metric retractions) All minimal representing loops in $\pi_1(M, p)$ are contained in a bounded set.

Escape phenomenon

$\sec \geq 0 \Rightarrow$ (by soul structure and metric retractions) All minimal representing loops in $\pi_1(M, p)$ are contained in a bounded set.

$\text{Ric} \geq 0$ & this boundedness condition \Rightarrow (by Cheeger–Gromoll's splitting theorem) $\pi_1(M)$ is virtually abelian.

Escape phenomenon


$\sec \geq 0 \Rightarrow$ (by soul structure and metric retractions) All minimal representing loops in $\pi_1(M, p)$ are contained in a bounded set.

$\text{Ric} \geq 0$ & this boundedness condition \Rightarrow (by Cheeger–Gromoll's splitting theorem) $\pi_1(M)$ is virtually abelian.

$\text{Ric} > 0$, $\pi_1(M)$ is infinite \Rightarrow (also by splitting theorem) These loops cannot be contained in any bounded set.

Example (Nabonnard 1980): a doubly warped product $\text{Ric} > 0$
 $M = [0, \infty) \times_f S^2 \times_h S^1$, $g = dr^2 + f(r)^2 ds_2^2 + h(r)^2 ds_1^2$.

$h(r)$ is decreasing \Rightarrow minimal representing loops of elements in $\pi_1(M, p)$ cannot be contained in any bounded sets.



Wei's examples (1988)

Let \tilde{N} be a simply connected (non-abelian) nilpotent Lie group and let Γ be a lattice in \tilde{N} . The nilmanifold $N = \tilde{N}/\Gamma$ admits a family of metrics $\{g_r\}_{r \geq 0}$ with

$$|\sec(g_r)| \rightarrow 0, \quad \text{diam}(N, g_r) \rightarrow 0 \text{ as } r \rightarrow \infty.$$

Wei's examples (1988)

Let \tilde{N} be a simply connected (non-abelian) nilpotent Lie group and let Γ be a lattice in \tilde{N} . The nilmanifold $N = \tilde{N}/\Gamma$ admits a family of metrics $\{g_r\}_{r \geq 0}$ with

$$|\sec(g_r)| \rightarrow 0, \quad \text{diam}(N, g_r) \rightarrow 0 \text{ as } r \rightarrow \infty.$$

For suitable $f(r)$ and g_r ,

$$M = [0, \infty) \times_f S^k \times N_r, \quad dr^2 + f^2(r)ds_k^2 + g_r$$

has $\text{Ric} > 0$ when k is large.

$\pi_1(M) = \pi_1(N) = \Gamma$ is not virtually abelian.

The escape phenomenon is also clear since $\text{diam}(N, g_r) \rightarrow 0$.

Escape rate

Quantify this escape phenomenon.

Definition (Pan 2020)

Let (M, p) be an open manifold with an infinite fundamental group. We define the escape rate of (M, p) , a scaling invariant, as

$$E(M, p) = \limsup_{|\gamma| \rightarrow \infty} \frac{\text{size}(c_\gamma)}{\text{length}(c_\gamma)},$$

where c_γ is a minimal representing geodesic loop of $\gamma \in \pi_1(M, p)$.

Convention: if $\pi_1(M)$ is finite, then we set $E(M, p) = 0$.

Escape rate

Quantify this escape phenomenon.

Definition (Pan 2020)

Let (M, p) be an open manifold with an infinite fundamental group. We define the escape rate of (M, p) , a scaling invariant, as

$$E(M, p) = \limsup_{|\gamma| \rightarrow \infty} \frac{\text{size}(c_\gamma)}{\text{length}(c_\gamma)},$$

where c_γ is a minimal representing geodesic loop of $\gamma \in \pi_1(M, p)$.

Convention: if $\pi_1(M)$ is finite, then we set $E(M, p) = 0$.

- $E(M, p) \leq 1/2$ always holds.
- (Sormani 2000) $E(M, p) < 1/2 \Rightarrow \pi_1(M)$ is finitely generated.
- The example by Brue–Naber–Semola has an infinitely generated $\pi_1(M) = \mathbb{Q}/\mathbb{Z}$, thus $E(M, p) = 1/2$.
- Wei's construction has $E(M, p) \geq c > 0$.

Small escape rate and virtual abelianness

Theorem (Pan 2020)

There is a constant $\epsilon(n) > 0$ s.t. if an open manifold M^n satisfies $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon(n)$, then $\pi_1(M)$ is virtually abelian.

Small escape rate and virtual abelianness

Theorem (Pan 2020)

There is a constant $\epsilon(n) > 0$ s.t. if an open manifold M^n satisfies $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon(n)$, then $\pi_1(M)$ is virtually abelian.

Equivariant asymptotic cone: $(r_i^{-1}\tilde{M}, \tilde{p}, \Gamma) \xrightarrow{GH} (Y, y, G)$.

$\Omega(\tilde{M}, \Gamma)$: the set of all equivariant asymptotic cones of (\tilde{M}, Γ) .

Small escape rate and virtual abelianness

Theorem (Pan 2020)

There is a constant $\epsilon(n) > 0$ s.t. if an open manifold M^n satisfies $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon(n)$, then $\pi_1(M)$ is virtually abelian.

Equivariant asymptotic cone: $(r_i^{-1}\tilde{M}, \tilde{p}, \Gamma) \xrightarrow{GH} (Y, y, G)$.

$\Omega(\tilde{M}, \Gamma)$: the set of all equivariant asymptotic cones of (\tilde{M}, Γ) .

$\text{sec} \geq 0 \Rightarrow$ unique $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$; Y is a metric cone with vertex y ; the orbit Gy is a Euclidean factor of Y .

$\text{Ric} \geq 0$: in general, Y may not be unique and may not be metric cones.

Small escape rate and virtual abelianness

Theorem (Pan 2020)

There is a constant $\epsilon(n) > 0$ s.t. if an open manifold M^n satisfies $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon(n)$, then $\pi_1(M)$ is virtually abelian.

Equivariant asymptotic cone: $(r_i^{-1}\tilde{M}, \tilde{p}, \Gamma) \xrightarrow{GH} (Y, y, G)$.

$\Omega(\tilde{M}, \Gamma)$: the set of all equivariant asymptotic cones of (\tilde{M}, Γ) .

$\text{sec} \geq 0 \Rightarrow$ unique $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$; Y is a metric cone with vertex y ; the orbit Gy is a Euclidean factor of Y .

$\text{Ric} \geq 0$: in general, Y may not be unique and may not be metric cones.

(Sketch) $E(M, p) = 0$

\Leftrightarrow (1) Gy is convex for all $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$;

\Leftrightarrow (2) Gy is a Euclidean factor of Y for all $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$.

Also, (2) \Rightarrow virtual abelianness.

Equivariant asymptotic geometry

Question: If M has $\text{Ric} \geq 0$ and $E(M, p) > 0$ (or $\pi_1(M)$ being torsion-free nilpotent), how is the equivariant asymptotic geometry?

It should be different from the ones with $\text{sec} \geq 0$.

Equivariant asymptotic geometry

Question: If M has $\text{Ric} \geq 0$ and $E(M, p) > 0$ (or $\pi_1(M)$ being torsion-free nilpotent), how is the equivariant asymptotic geometry?

It should be different from the ones with $\text{sec} \geq 0$.

Theorem (Pan–Wei 2021)

Examples of Ricci limits (as asymptotic cones) where Hausdorff dimension of the singular set exceeds that of the regular set.

Actually, they are isometric to the α -Grushin halfplane $\mathbb{G}^+(\alpha)$, a classical example from sub/almost Riemannian geometry.

$g = dr^2 + r^{-2\alpha} dv^2$ on $[0, \infty) \times \mathbb{R}$.

$\mathcal{S} = \{0\} \times \mathbb{R}$ has Hausdorff dimension $1 + \alpha$.

Asymptotic geometry of Wei's example

Open manifold M with $\text{Ric} > 0$ and $\pi_1(M) = H^3(\mathbb{Z})$ (Wei's construction). Then unique $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$ satisfies:

- G is the 3-dimensional simply connected Heisenberg group.
- The orbit Gy is not convex.
- Y is not a metric cone.
- Let $L(\simeq \mathbb{R})$ be the center of G . Then $\dim_{\mathcal{H}}(Ly) \geq 2$.

Asymptotic geometry of Wei's example

Open manifold M with $\text{Ric} > 0$ and $\pi_1(M) = H^3(\mathbb{Z})$ (Wei's construction). Then unique $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$ satisfies:

- G is the 3-dimensional simply connected Heisenberg group.
- The orbit Gy is not convex.
- Y is not a metric cone.
- Let $L(\simeq \mathbb{R})$ be the center of G . Then $\dim_{\mathcal{H}}(Ly) \geq 2$.

Questions

Does any of these conditions relate to abelianness/nilpotency?

- (1) G is (virtually) abelian for all $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$;
- (2) The orbit Gy is convex for all $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$;
- (3) (Y, y) is a metric cone with vertex y for all $(Y, y) \in \Omega(\tilde{M})$.
- (4) Hausdorff dimension of (closed) \mathbb{R} -orbits at y ?

Answers to (1), (2), and (3)

Remark: $\sec \geq 0 \Rightarrow (1,2,3)$ all hold; also, $\dim_{\mathcal{H}}(Ly) = 1$ for all $(Y, y) \in \Omega(\tilde{M}, \Gamma)$ and all closed \mathbb{R} -subgroup L of $\text{Isom}(Y)$.

Answers to (1), (2), and (3)

Remark: $\sec \geq 0 \Rightarrow (1,2,3)$ all hold; also, $\dim_{\mathcal{H}}(Ly) = 1$ for all $(Y, y) \in \Omega(\tilde{M}, \Gamma)$ and all closed \mathbb{R} -subgroup L of $\text{Isom}(Y)$.

We have seen that: $E(M, p) = 0 \Leftrightarrow (2) \Rightarrow$ virtual abelianness.

(1) \nRightarrow virtual abelianness.

In fact, one can modify Wei's example s.t. G is isomorphic to \mathbb{R}^3 for all asymptotic limits but $\Gamma = H^3(\mathbb{Z})$.

Answers to (1), (2), and (3)

Remark: $\sec \geq 0 \Rightarrow (1,2,3)$ all hold; also, $\dim_{\mathcal{H}}(Ly) = 1$ for all $(Y, y) \in \Omega(\tilde{M}, \Gamma)$ and all closed \mathbb{R} -subgroup L of $\text{Isom}(Y)$.

We have seen that: $E(M, p) = 0 \Leftrightarrow (2) \Rightarrow$ virtual abelianness.

(1) \nRightarrow virtual abelianness.

In fact, one can modify Wei's example s.t. G is isomorphic to \mathbb{R}^3 for all asymptotic limits but $\Gamma = H^3(\mathbb{Z})$.

We say that M is *conic at infinity*, if any asymptotic cone (Y, y) of M is a metric cone (may not be unique) with vertex y .

Theorem (Pan 2022)

Let (M, p) be an open n -manifold with $\text{Ric} \geq 0$ and $E(M, p) \neq \frac{1}{2}$.

(1) If \tilde{M} is conic at infinity, then $\pi_1(M)$ is virtually abelian.

(2) If \tilde{M} has Euclidean volume growth of constant at least L , then $\pi_1(M)$ is $C(n, L)$ -abelian.

Nilpotency step

Question (Pan–Wei 2021)

Is the nilpotency step of $\pi_1(M)$ related to the Hausdorff dimension of some isometric \mathbb{R} -action in some asymptotic cone of \tilde{M} ?

Nilpotency step

Question (Pan–Wei 2021)

Is the nilpotency step of $\pi_1(M)$ related to the Hausdorff dimension of some isometric \mathbb{R} -action in some asymptotic cone of \tilde{M} ?

Let N be a finitely generated torsion-free nilpotent group.

$N = N^0 \triangleright N^1 \triangleright \dots \triangleright N^k = \{e\}$, where $N^{j+1} = [N, N^j]$

Nilpotency step: $\text{step}(N) = \text{smallest integer } k \text{ s.t. } N^k = \{e\}$.

Example: $N = H^3(\mathbb{Z})$ has $\text{step}(N) = 2$.

Nilpotency step

Question (Pan–Wei 2021)

Is the nilpotency step of $\pi_1(M)$ related to the Hausdorff dimension of some isometric \mathbb{R} -action in some asymptotic cone of \tilde{M} ?

Let N be a finitely generated torsion-free nilpotent group.

$N = N^0 \triangleright N^1 \triangleright \dots \triangleright N^k = \{e\}$, where $N^{j+1} = [N, N^j]$

Nilpotency step: $\text{step}(N) = \text{smallest integer } k \text{ s.t. } N^k = \{e\}$.

Example: $N = H^3(\mathbb{Z})$ has $\text{step}(N) = 2$.

Let Γ be a finitely generated and virtual nilpotent group.

Then Γ has a torsion-free nilpotent subgroup N of finite index.

Define $\text{vir.step}(\Gamma) := \text{step}(N)$. (It is well-defined.)

In particular, this definition applies to any finitely generated $\pi_1(M)$ with $\text{Ric} \geq 0$.

Let M be a closed manifold with an infinite and virtually nilpotent fundamental group Γ (or equivalently, of polynomial growth).

(Gromov 1981, Pansu 1983) (\tilde{M}, \tilde{p}) has a unique asymptotic cone (G, e, d) , which is a simply connected stratified nilpotent Lie group G with a complete left-invariant subFinsler metric (Carnot group).

Let M be a closed manifold with an infinite and virtually nilpotent fundamental group Γ (or equivalently, of polynomial growth).

(Gromov 1981, Pansu 1983) (\tilde{M}, \tilde{p}) has a unique asymptotic cone (G, e, d) , which is a simply connected stratified nilpotent Lie group G with a complete left-invariant subFinsler metric (Carnot group).

It holds that:

$$\text{step}(G) = \text{vir.step}(\Gamma);$$

$\dim_{\mathcal{H}}(L, d) = \text{vir.step}(\Gamma)$, where L is any \mathbb{R} -subgroup in the last nontrivial subgroup of the lower central series of G .

Nilpotency step and Hausdorff dimension

Let (M, p) be an open n -manifold with $\text{Ric} \geq 0$, an infinite fundamental group Γ , and $E(M, p) \neq 1/2$.

Theorem (Pan 2023)

There exists $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$ and a closed \mathbb{R} -subgroup L of G s.t. $\dim_{\mathcal{H}}(Ly) \geq \text{vir.step}(\Gamma)$.

Theorem (Pan 2023)

For any $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$, the orbit Gy has a natural structure as a connected and simply connected nilpotent Lie group with $\text{step}(Gy) \leq \text{vir.step}(\Gamma)$.

Nilpotency step and Hausdorff dimension

Let (M, p) be an open n -manifold with $\text{Ric} \geq 0$, an infinite fundamental group Γ , and $E(M, p) \neq 1/2$.

Theorem (Pan 2023)

There exists $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$ and a closed \mathbb{R} -subgroup L of G s.t. $\dim_{\mathcal{H}}(Ly) \geq \text{vir.step}(\Gamma)$.

Theorem (Pan 2023)

For any $(Y, y, G) \in \Omega(\tilde{M}, \Gamma)$, the orbit Gy has a natural structure as a connected and simply connected nilpotent Lie group with $\text{step}(Gy) \leq \text{vir.step}(\Gamma)$.

Cf. When $\pi_1(M) = \mathbb{Q}/\mathbb{Z}$ (Brue-Naber-Semola), $E = 1/2$ and asymptotic orbit Gy could be compact or disconnected.

Nilpotency step and growth of length (upper bound)

Discrete Heisenberg 3-group $\Gamma = H^3(\mathbb{Z})$:

Let γ in the center of Γ . γ^m has word length

$$C_1 m^{1/2} \leq d_W(e, \gamma^m) \leq C_2 m^{1/2}.$$

Nilpotency step and growth of length (upper bound)

Discrete Heisenberg 3-group $\Gamma = H^3(\mathbb{Z})$:

Let γ in the center of Γ . γ^m has word length

$C_1 m^{1/2} \leq d_W(e, \gamma^m) \leq C_2 m^{1/2}$. Then

$$|\gamma^m| := d(\gamma^m \tilde{p}, \tilde{p}) \leq C m^{1/2}.$$

To prove virtual abelianness, it suffices to find a lower bound for $|\gamma^m|$ that contradicts $|\gamma^m| \leq C m^{1/2}$.

Nilpotency step and growth of length (upper bound)

Discrete Heisenberg 3-group $\Gamma = H^3(\mathbb{Z})$:

Let γ in the center of Γ . γ^m has word length

$C_1 m^{1/2} \leq d_W(e, \gamma^m) \leq C_2 m^{1/2}$. Then

$$|\gamma^m| := d(\gamma^m \tilde{p}, \tilde{p}) \leq C m^{1/2}.$$

To prove virtual abelianness, it suffices to find a lower bound for $|\gamma^m|$ that contradicts $|\gamma^m| \leq C m^{1/2}$.

In general, let N be a torsion-free nilpotent group with nilpotency step k . Take $\gamma \in N^{k-1} - \{e\}$. Then

$$|\gamma^m| \leq C m^{1/k}.$$

Asymptotic geometry and growth of length (lower bound)

$\gamma \in N^{k-1} - \{e\}$, where N torsion-free nilpotent of step k .

Small escape rate

\Rightarrow Limit orbit is always “close” to a standard Euclidean subspace

\Rightarrow almost translation estimate at large: $\forall \epsilon > 0$,

$|\gamma^2| \geq (1.99 - \epsilon)|\gamma|$ holds for all γ with sufficiently large $|\gamma|$.

Asymptotic geometry and growth of length (lower bound)

$\gamma \in N^{k-1} - \{e\}$, where N torsion-free nilpotent of step k .

Small escape rate

\Rightarrow Limit orbit is always “close” to a standard Euclidean subspace

\Rightarrow almost translation estimate at large: $\forall \epsilon > 0$,

$|\gamma^2| \geq (1.99 - \epsilon)|\gamma|$ holds for all γ with sufficiently large $|\gamma|$.

\tilde{M} conic at infinity & $E \neq 1/2$

\Rightarrow almost linear growth estimate: $\forall \epsilon > 0$, \exists a constant C' s.t.

$|\gamma^m| \geq C' m^{1-\epsilon}$ for all m large.

Asymptotic geometry and growth of length (lower bound)

$\gamma \in N^{k-1} - \{e\}$, where N torsion-free nilpotent of step k .

Small escape rate

\Rightarrow Limit orbit is always “close” to a standard Euclidean subspace

\Rightarrow almost translation estimate at large: $\forall \epsilon > 0$,

$|\gamma^2| \geq (1.99 - \epsilon)|\gamma|$ holds for all γ with sufficiently large $|\gamma|$.

\tilde{M} conic at infinity & $E \neq 1/2$

\Rightarrow almost linear growth estimate: $\forall \epsilon > 0$, \exists a constant C' s.t.

$|\gamma^m| \geq C'm^{1-\epsilon}$ for all m large.

Define $\mathcal{D} := \sup\{\dim_{\mathcal{H}}(Ly) | (Y, y, L) \in \Omega(\tilde{M}, \langle \gamma \rangle)\}$.

(Rem. As a crucial intermediate step, Ly is homeomorphic to \mathbb{R} .)

$E \neq 1/2$

$\Rightarrow \forall \epsilon > 0$, \exists a constant C' s.t. $|\gamma^m| \geq C'm^{(1/\mathcal{D})-\epsilon}$ for all m large.

$\Rightarrow \mathcal{D} \geq l$.

Relation to previous results

This result generalizes the previous results on virtual abelianness.

Relation to metric cones

Let $Y = C(Z)$ be a metric cone with vertex y . Then $\dim_{\mathcal{H}}(Ly) = 1$ for any closed \mathbb{R} -subgroups $L \leq \text{Isom}(Y)$.

\Rightarrow If $E(M, p) \neq 1/2$ and \tilde{M} is conic at infinity, then $\text{vir.step}(\pi_1(M)) \leq \sup \dim_{\mathcal{H}}(Ly) = 1$.

Relation to previous results

This result generalizes the previous results on virtual abelianness.

Relation to metric cones

Let $Y = C(Z)$ be a metric cone with vertex y . Then $\dim_{\mathcal{H}}(Ly) = 1$ for any closed \mathbb{R} -subgroups $L \leq \text{Isom}(Y)$.

\Rightarrow If $E(M, p) \neq 1/2$ and \tilde{M} is conic at infinity, then $\text{vir.step}(\pi_1(M)) \leq \sup \dim_{\mathcal{H}}(Ly) = 1$.

Relation to small escape rate

Let M be an open n -manifold with $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon$. Then $\dim_{\mathcal{H}}(Ly) \leq 1 + \delta(\epsilon|n)$, where $\delta(\epsilon|n) \rightarrow 0$ as $\epsilon \rightarrow 0$.

$\Rightarrow \pi_1(M)$ is virtually abelian when ϵ is small.

Open questions

Question (Finite generation v.s. $E < 1/2$)

If an open M has $\text{Ric} \geq 0$ and $\pi_1(M)$ is finitely generated, is it true that $E(M, p) < 1/2$?

Question (Escape rate hierarchy)

Given n , are there constants $0 < \epsilon(n, 1) < \dots < \epsilon(n, l(n)) < 1/2$ such that if an open M^n has $\text{Ric} \geq 0$ and $E(M, p) \leq \epsilon(n, k)$, then $\text{step}(\pi_1(M)) \leq k$?

Question (Minimal dimension with nilpotent π_1)

Let M be an open n -manifold with $\text{Ric} \geq 0$. Does $n \leq 12$ imply that any finitely generated subgroup of $\pi_1(M)$ is virtually abelian?

One more thing...

One more thing...

Let M be an open manifold with $\text{Ric} \geq 0$. We say that M has *linear (or minimal) volume growth* if

$$\limsup_{r \rightarrow \infty} \frac{\text{vol}(B_r(p))}{r} < \infty.$$

(Calabi/Yau ~ 1975) linear volume growth is minimal.

(Sormani 2000) $\pi_1(M)$ is finitely generated.

One more thing...

Let M be an open manifold with $\text{Ric} \geq 0$. We say that M has *linear (or minimal) volume growth* if

$$\limsup_{r \rightarrow \infty} \frac{\text{vol}(B_r(p))}{r} < \infty.$$

(Calabi/Yau ~ 1975) linear volume growth is minimal.

(Sormani 2000) $\pi_1(M)$ is finitely generated.

Theorem (Navarro–Pan–Zhu 2024)

Let M be an open manifold with $\text{Ric} \geq 0$ and linear volume growth. Then $\pi_1(M)$ is virtually abelian. (In fact, $E(M, p) = 0$.)

Theorem (Navarro–Pan–Zhu 2024)

Let M be an open manifold with $\text{Ric} > 0$ and linear volume growth. Then $\pi_1(M)$ is finite.