

Semi-local simple connectedness of non-collapsing Ricci limit spaces

Jiayin Pan (Joint work with Guofang Wei)

UC Santa Barbara

Gromov-Hausdorff distance

The Gromov-Hausdorff distance between two metric spaces measures how they look alike.

Gromov-Hausdorff distance

The Gromov-Hausdorff distance between two metric spaces measures how they look alike.

Gromov's precompactness theorem

Let $\{(M_i, p_i)\}_i$ be a sequence of complete Riemannian n -manifolds of

$$\operatorname{Ric}_{M_i} \geq -(n-1),$$

then $\{(M_i, p_i)\}_i$ has a convergent subsequence with respect to the (pointed) Gromov-Hausdorff distance.

Gromov-Hausdorff distance

The Gromov-Hausdorff distance between two metric spaces measures how they look alike.

Gromov's precompactness theorem

Let $\{(M_i, p_i)\}_i$ be a sequence of complete Riemannian n -manifolds of

$$\operatorname{Ric}_{M_i} \geq -(n-1),$$

then $\{(M_i, p_i)\}_i$ has a convergent subsequence with respect to the (pointed) Gromov-Hausdorff distance.

$(M_i, p_i) \xrightarrow{GH} (X, p)$...a useful tool to study a class of Riemannian manifolds with geometric conditions (curvature, volume, diameter, etc.)/to study infinitesimal or asymptotic geometry...important to understand the structure of the limit space X /relations between X and M_i for i large.

Topology of limit spaces

$(M_i^n, p_i) \xrightarrow{GH} (X, p)$, where M_i are complete.

Topology of limit spaces

$(M_i^n, p_i) \xrightarrow{GH} (X, p)$, where M_i are complete.

$\sec \geq -1$, X Alexandrov space (Burago-Gromov-Perelman)
 X is locally contractible (Perelman)

Topology of limit spaces

$(M_i^n, p_i) \xrightarrow{GH} (X, p)$, where M_i are complete.

$\sec \geq -1$, X Alexandrov space (Burago-Gromov-Perelman)
 X is locally contractible (Perelman)

$\text{Ric} \geq -(n-1)$, X Ricci limit space (Cheeger-Colding)
Even when non-collapsing, X may have
infinite local topological type (Menguy)

“Non-collapsing” means there is $\nu > 0$ such that $\text{vol}(B_1(p_i)) \geq \nu$
for all i .

Topology of Ricci limit spaces

Since Ric controls fundamental groups, it is natural to ask:

Question

Is X always (semi-)locally simply connected?

Topology of Ricci limit spaces

Since Ric controls fundamental groups, it is natural to ask:

Question

Is X always (semi-)locally simply connected?

Recall: Semi-locally simply connected \Leftrightarrow there exists a simply connected universal cover.

Topology of Ricci limit spaces

Since Ric controls fundamental groups, it is natural to ask:

Question

Is X always (semi-)locally simply connected?

Recall: Semi-locally simply connected \Leftrightarrow there exists a simply connected universal cover.

Theorem (Sormani-Wei, 2001)

Any Ricci limit space has a universal cover.

Remark: “Universal” in the sense of universal property of covering maps; unknown whether it is simply connected.

Topology of Ricci limit spaces

Since Ric controls fundamental groups, it is natural to ask:

Question

Is X always (semi-)locally simply connected?

Recall: Semi-locally simply connected \Leftrightarrow there exists a simply connected universal cover.

Theorem (Sormani-Wei, 2001)

Any Ricci limit space has a universal cover.

Remark: “Universal” in the sense of universal property of covering maps; unknown whether it is simply connected.



Theorem (Mondino-Wei, 2016)

Let (X, d, m) be an $\mathrm{RCD}^*(K, N)$ -space for some $K \in \mathbb{R}$, $N \in (1, +\infty)$. Then (X, d, m) admits a universal cover.

Examples

A Hawaii ring does not have a universal cover.

The Griffiths twin cone is one-point-join of two “cutted” cones over a Hawaii ring. It is not semi-locally simply connected, but it has a universal cover (as itself); also, in this example, a non-contractible loop may have infinite length.



Main Theorem (Pan-Wei, 2019)

Any non-collapsing Ricci limit space is semi-locally simply connected.

Methods for sectional curvature case

Methods for sectional curvature case

Method 1 : Tangent cones are metric cones (BGP);

Construct a homeomorphism from tangent cone to a local nbhd (Perelman)/Gradient flow of distance function

Methods for sectional curvature case

Method 1 : Tangent cones are metric cones (BGP);

True for non-collapsing Ricci limits (CC).

Construct a homeomorphism from tangent cone to a local nbhd (Perelman)/Gradient flow of distance function

Impossible for Ricci case, due to Menguy's example.

Methods for sectional curvature case

Method 1 : Tangent cones are metric cones (BGP);

True for non-collapsing Ricci limits (CC).

Construct a homeomorphism from tangent cone to a local nbhd (Perelman)/Gradient flow of distance function

Impossible for Ricci case, due to Menguy's example.

Method 2 : With the non-collapsing condition, control the local contractibility radius (Grove-Petersen);

then pass this property to the limit (Borsuk, Petersen).

Methods for sectional curvature case

Method 1 : Tangent cones are metric cones (BGP);

True for non-collapsing Ricci limits (CC).

Construct a homeomorphism from tangent cone to a local nbhd (Perelman)/Gradient flow of distance function

Impossible for Ricci case, due to Menguy's example.

Method 2 : With the non-collapsing condition, control the local contractibility radius (Grove-Petersen);

Not true for Ricci case on π_1 -level; examples by Otsu.

then pass this property to the limit (Borsuk, Petersen).

True (even without curvature conditions), but requires uniform control at all points.

Theorem (Grove-Petersen)

Given $n, \kappa, \nu > 0$, there exist positive constants $\epsilon(n, \kappa, \nu)$ and $C(n, \kappa, \nu)$ such that for any complete n -manifold (M, p) of

$$\sec_M \geq -\kappa, \quad \text{vol}(B_1(p)) \geq \nu,$$

$B_r(p)$ is contractible in $B_{Cr}(p)$, where $r \in [0, \epsilon)$.

Methods for sectional curvature case

Theorem (Grove-Petersen)

Given $n, \kappa, \nu > 0$, there exist positive constants $\epsilon(n, \kappa, \nu)$ and $C(n, \kappa, \nu)$ such that for any complete n -manifold (M, p) of

$$\sec_M \geq -\kappa, \quad \text{vol}(B_1(p)) \geq \nu,$$

$B_r(p)$ is contractible in $B_{Cr}(p)$, where $r \in [0, \epsilon)$.

Otsu's example

$S^3 \times \mathbb{R}P^2$ admits a sequence of Riemannian metrics g_i of positive Ricci curvature with a non-collapsing limit space as a metric suspension over $S^2 \times \mathbb{R}P^2$. At the “tip” point, there are shorter and shorter non-contractible loops.

Methods for sectional curvature case

Theorem (Grove-Petersen)

Given $n, \kappa, \nu > 0$, there exist positive constants $\epsilon(n, \kappa, \nu)$ and $C(n, \kappa, \nu)$ such that for any complete n -manifold (M, p) of

$$\sec_M \geq -\kappa, \quad \text{vol}(B_1(p)) \geq \nu,$$

$B_r(p)$ is contractible in $B_{Cr}(p)$, where $r \in [0, \epsilon)$.

Otsu's example

$S^3 \times \mathbb{R}P^2$ admits a sequence of Riemannian metrics g_i of positive Ricci curvature with a non-collapsing limit space as a metric suspension over $S^2 \times \mathbb{R}P^2$. At the “tip” point, there are shorter and shorter non-contractible loops.

Also, the Eguchi-Hanson metric on the tangent bundle of $\mathbb{R}P^2$ gives a Ricci flat example.

Definition (1-contractibility radius):

$$\rho(t, x) = \inf\{\infty, \rho \geq t \mid \text{any loop in } B_t(x) \text{ is contractible in } B_\rho(x)\}.$$

Remark: X is semi-locally simply connected if for any $x \in X$, there is $T > 0$ such that $\rho(T, x) < \infty$; X is locally simply connected if for any $x \in X$, there is $t_i \rightarrow 0$ such that $\rho(t_i, x) = t_i$.

Definition (1-contractibility radius):

$$\rho(t, x) = \inf\{\infty, \rho \geq t \mid \text{any loop in } B_t(x) \text{ is contractible in } B_\rho(x)\}.$$

Remark: X is semi-locally simply connected if for any $x \in X$, there is $T > 0$ such that $\rho(T, x) < \infty$; X is locally simply connected if for any $x \in X$, there is $t_i \rightarrow 0$ such that $\rho(t_i, x) = t_i$.

Local version of the main theorem with an estimate on $\rho(t, x)$

Let (M_i, p_i) be a sequence of Riemannian n -manifolds (not necessarily complete) converging to (X, p) such that for all i ,

- (1) $B_2(p_i) \cap \partial M_i = \emptyset$ and the closure of $B_2(p_i)$ is compact;
- (2) $\text{Ric} \geq -(n-1)$ on $B_2(p_i)$, $\text{vol}(B_1(p_i)) \geq \nu > 0$.

Then

$$\lim_{t \rightarrow 0} \frac{\rho(t, x)}{t} = 1$$

holds for any $x \in B_1(p)$.

Approach

Key: show $\lim_{t \rightarrow 0} \rho(t, x) = 0$.

After this, we can improve the result to $\lim_{t \rightarrow 0} \rho(t, x)/t = 1$ by using the structure of tangent cones and Sormani's uniform cut techniques.

Key: show $\lim_{t \rightarrow 0} \rho(t, x) = 0$.

After this, we can improve the result to $\lim_{t \rightarrow 0} \rho(t, x)/t = 1$ by using the structure of tangent cones and Sormani's uniform cut techniques.

Classification: We classify the points in X according to the local 1-contractibility radius on manifolds: $x \in X$, let x_i in M_i to x ;

- x is of type I, if there is $r > 0$ such that the family of functions $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is equi-continuous at $t = 0$;
- x is of type II, if $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is not equi-continuous at $t = 0$;
- x is of type III, if it is not of type I nor type II. (In other words, $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is equi-continuous at $t = 0$, but the family $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is not equi-continuous for any $r > 0$.)

Key: show $\lim_{t \rightarrow 0} \rho(t, x) = 0$.

After this, we can improve the result to $\lim_{t \rightarrow 0} \rho(t, x)/t = 1$ by using the structure of tangent cones and Sormani's uniform cut techniques.

Classification: We classify the points in X according to the local 1-contractibility radius on manifolds: $x \in X$, let x_i in M_i to x ;

- x is of type I, if there is $r > 0$ such that the family of functions $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is equi-continuous at $t = 0$;
- x is of type II, if $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is not equi-continuous at $t = 0$;
- x is of type III, if it is not of type I nor type II. (In other words, $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is equi-continuous at $t = 0$, but the family $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is not equi-continuous for any $r > 0$.)

Remark: Type II point may exist due to Otsu's example.

Key: show $\lim_{t \rightarrow 0} \rho(t, x) = 0$.

After this, we can improve the result to $\lim_{t \rightarrow 0} \rho(t, x)/t = 1$ by using the structure of tangent cones and Sormani's uniform cut techniques.

Classification: We classify the points in X according to the local 1-contractibility radius on manifolds: $x \in X$, let x_i in M_i to x ;

- x is of type I, if there is $r > 0$ such that the family of functions $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is equi-continuous at $t = 0$;
- x is of type II, if $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is not equi-continuous at $t = 0$;
- x is of type III, if it is not of type I nor type II. (In other words, $\{\rho(x_i, t)\}_{i \in \mathbb{N}}$ is equi-continuous at $t = 0$, but the family $\{\rho(q, t) | q \in B_r(x_i), i \in \mathbb{N}\}$ is not equi-continuous for any $r > 0$.)

Remark: Type II point may exist due to Otsu's example.

Spoiler: Type III points are the most difficult ones to deal with.

Type I

For a family of functions $\{\rho_\alpha(t)\}_{\alpha \in A}$ with $\rho_\alpha(0) = 0$, the family is equi-continuous at $t = 0$ if and only if there is a continuous function $\lambda(t)$ defined on $[0, T)$ with $\lambda(0) = 0$ such that $\rho_\alpha(t) \leq \lambda(t)$ for all $t \in [0, T)$ and all $\alpha \in A$.

Type I

For a family of functions $\{\rho_\alpha(t)\}_{\alpha \in A}$ with $\rho_\alpha(0) = 0$, the family is equi-continuous at $t = 0$ if and only if there is a continuous function $\lambda(t)$ defined on $[0, T)$ with $\lambda(0) = 0$ such that $\rho_\alpha(t) \leq \lambda(t)$ for all $t \in [0, T)$ and all $\alpha \in A$.

For **type I** points, we can pass the local 1-contractibility control on local balls around x_i to that around x in the limit space:

Type I

For a family of functions $\{\rho_\alpha(t)\}_{\alpha \in A}$ with $\rho_\alpha(0) = 0$, the family is equi-continuous at $t = 0$ if and only if there is a continuous function $\lambda(t)$ defined on $[0, T)$ with $\lambda(0) = 0$ such that $\rho_\alpha(t) \leq \lambda(t)$ for all $t \in [0, T)$ and all $\alpha \in A$.

For **type I** points, we can pass the local 1-contractibility control on local balls around x_i to that around x in the limit space:

Proposition (without curvature conditions)

Let (X_i, x_i) be a sequence of length metric spaces with the conditions below:

- (1) the closure of $B_1(x_i)$ is compact;
- (2) there is a nice function λ on $[0, T)$ such that for all i and all $q \in B_2(x_i)$, $\rho(t, q) \leq \lambda(t) < 1/2$ holds on $[0, T)$;
- (3) $(X_i, x_i) \xrightarrow{GH} (Y, y)$.

Then $\rho(t, q) \leq \lambda(t)$ for all $t \in [0, T)$ and all $q \in B_1(y)$.

Sketch: (Borsuk's method)

Sketch: (Borsuk's method)

For a loop c in the limit space, we can find a sequence of loops c_i in M_i that converges uniformly to c .

Sketch: (Borsuk's method)

For a loop c in the limit space, we can find a sequence of loops c_i in M_i that converges uniformly to c .

For c in a sufficiently small ball, we can contract c_i for a large i . Take a skeleton of this homotopy, then we transfer this skeleton to the next manifold in the sequence and fill in the skeleton there. This allows us to transfer the nullhomotopy of c_i along the sequence and pass it to the limit space by uniform convergence. In this construction, we need to control the distance between nullhomotopies to assure uniform convergence.

Type II

A point x is of **Type II** means there are $\epsilon > 0$ and $t_i \rightarrow 0$ such that $\rho(t_i, x_i) \geq \epsilon$. We consider the universal covering space of $B_\epsilon(x_i)$. The local fundamental group Γ_i has a subgroup generated by these non-contractible “small” loops H_i . In Otsu’s example, $\Gamma_i = \mathbb{Z}_2$. We consider the equivariant Gromov-Hausdorff convergence:

$$\begin{array}{ccc} (U_i, y_i, \Gamma_i, H_i) & \xrightarrow{GH} & (Y, y, G, H) \\ \downarrow \pi_i & & \downarrow \pi \\ (B_\epsilon(x_i), x_i) & \xrightarrow{GH} & (B_\epsilon(x) = Y/G, x), \end{array}$$

Type II

A point x is of **Type II** means there are $\epsilon > 0$ and $t_i \rightarrow 0$ such that $\rho(t_i, x_i) \geq \epsilon$. We consider the universal covering space of $B_\epsilon(x_i)$. The local fundamental group Γ_i has a subgroup generated by these non-contractible “small” loops H_i . In Otsu’s example, $\Gamma_i = \mathbb{Z}_2$. We consider the equivariant Gromov-Hausdorff convergence:

$$\begin{array}{ccc} (U_i, y_i, \Gamma_i, H_i) & \xrightarrow{GH} & (Y, y, G, H) \\ \downarrow \pi_i & & \downarrow \pi \\ (B_\epsilon(x_i), x_i) & \xrightarrow{GH} & (B_\epsilon(x) = Y/G, x), \end{array}$$

H_i has a uniform order upper bound, $\#H_i \leq N$ (Anderson). This also implies that H -action fixes y .

Type II

A point x is of **Type II** means there are $\epsilon > 0$ and $t_i \rightarrow 0$ such that $\rho(t_i, x_i) \geq \epsilon$. We consider the universal covering space of $B_\epsilon(x_i)$. The local fundamental group Γ_i has a subgroup generated by these non-contractible “small” loops H_i . In Otsu’s example, $\Gamma_i = \mathbb{Z}_2$. We consider the equivariant Gromov-Hausdorff convergence:

$$\begin{array}{ccc} (U_i, y_i, \Gamma_i, H_i) & \xrightarrow{GH} & (Y, y, G, H) \\ \downarrow \pi_i & & \downarrow \pi \\ (B_\epsilon(x_i), x_i) & \xrightarrow{GH} & (B_\epsilon(x) = Y/G, x), \end{array}$$

H_i has a uniform order upper bound, $\#H_i \leq N$ (Anderson). This also implies that H -action fixes y .

Roughly speaking, covering group increases the volume around y_i when compared with the volume around x_i . Together with volume convergence (Colding, CC), local volume of y is at least double of the local volume of x : $\mathcal{H}^n(B_s(y)) \geq 2 \cdot \mathcal{H}^n(B_s(x))$.

Half-volume lower bounds

Recall volume comparison: $\mathcal{H}^n(B_s(y))$ is no greater than the volume s -ball in the space form.

Half-volume lower bounds

Recall volume comparison: $\mathcal{H}^n(B_s(y))$ is no greater than the volume s -ball in the space form.

Therefore, if x has local volume strictly larger than half of corresponding volume in the space form, then the above diagram cannot happen. In fact, this implies such a point x must be type I.

Half-volume lower bounds

Recall volume comparison: $\mathcal{H}^n(B_s(y))$ is no greater than the volume s -ball in the space form.

Therefore, if x has local volume strictly larger than half of corresponding volume in the space form, then the above diagram cannot happen. In fact, this implies such a point x must be type I.

Theorem (By-product for half-volume lower bounds)

Given $n \geq 2$, $\kappa \geq 0$, and $\omega > 1/2$, there exist positive constants $\epsilon(n, \kappa, \omega)$ and $C(n, \kappa, \omega)$ such that the following holds.

Let (M, p) be a complete n -manifold satisfying

$$\text{Ric} \geq -(n-1)\kappa, \quad \text{vol}(B_1(p)) \geq \omega \cdot \text{vol}(B_1^n(-\kappa)).$$

Then every loop in $B_r(p)$ is contractible in $B_{Cr}(p)$, where $r \in [0, \epsilon)$.

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, $=$ holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

If $\omega(x) > 1/4$ for all $x \in B_1(p)$:

- Case x is of type I, we are good;

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

If $\omega(x) > 1/4$ for all $x \in B_1(p)$:

- Case x is of type I, we are good;
- Case x is of type II...go to local covers...the corresponding limit point y has $\omega(y) > 1/2$...lift the loop in $B_\varepsilon(x)$ to a loop in $B_\varepsilon(y)$...project the homotopy down...we are good;

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

If $\omega(x) > 1/4$ for all $x \in B_1(p)$:

- Case x is of type I, we are good;
- Case x is of type II...go to local covers...the corresponding limit point y has $\omega(y) > 1/2$...lift the loop in $B_\varepsilon(x)$ to a loop in $B_\varepsilon(y)$...project the homotopy down...we are good;
- Case x is of type III...main technical part...no local control around x_i to utilize...no local covers to build...

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

If $\omega(x) > 1/4$ for all $x \in B_1(p)$:

- Case x is of type I, we are good;
- Case x is of type II...go to local covers...the corresponding limit point y has $\omega(y) > 1/2$...lift the loop in $B_\varepsilon(x)$ to a loop in $B_\varepsilon(y)$...project the homotopy down...we are good;
- Case x is of type III...main technical part...no local control around x_i to utilize...no local covers to build...

...

Induction on local volume

Define $\omega(x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r(x))}{\text{vol}(B_r^n(0))}$.

Remark: $0 < \omega(x) \leq 1$, = holds iff x is a regular point. $\omega(x)$ has a uniform lower bound for all $x \in B_1(p)$.

We use an induction argument on $\omega(x)$:

If $\omega(x) > 1/2$ for all $x \in B_1(p)$, then every point $x \in B_1(p)$ is of type I...we are good.

If $\omega(x) > 1/4$ for all $x \in B_1(p)$:

- Case x is of type I, we are good;
- Case x is of type II...go to local covers...the corresponding limit point y has $\omega(y) > 1/2$...lift the loop in $B_\varepsilon(x)$ to a loop in $B_\varepsilon(y)$...project the homotopy down...we are good;
- Case x is of type III...main technical part...no local control around x_i to utilize...no local covers to build...

...We need a technical lemma to deal with type III points.

Technical lemma (without curvature conditions)

(Roughly speaking,) If all type II points are good, then all type III points are good as well.

Technical lemma (without curvature conditions)

(Roughly speaking,) If all type II points are good, then all type III points are good as well.

Assuming this, we clear the quarter volume lower bound case and we can continue the induction argument.

Technical lemma (without curvature conditions)

(Roughly speaking,) If all type II points are good, then all type III points are good as well.

Assuming this, we clear the quarter volume lower bound case and we can continue the induction argument.

On Technical Lemma:

By definition, if x is of type III, then the family $\{\rho(t, x_i)\}_{i \in \mathbb{N}}$ is equi-continuous at $t = 0$.

Technical lemma (without curvature conditions)

(Roughly speaking,) If all type II points are good, then all type III points are good as well.

Assuming this, we clear the quarter volume lower bound case and we can continue the induction argument.

On Technical Lemma:

By definition, if x is of type III, then the family $\{\rho(t, x_i)\}_{i \in \mathbb{N}}$ is equi-continuous at $t = 0$.

A trivial, but important, observation: for any x , either $\rho(t, x)$ is good (type II case, by assumption), or $\{\rho(t, x_i)\}_{i \in \mathbb{N}}$ is good (type III case); for type I points, both $\rho(t, x)$ and $\{\rho(t, x_i)\}_{i \in \mathbb{N}}$ are good.


A very brief sketch of the proof of technical lemma

To better illustrate the proof, we first present an alternative proof for type I points.

A very brief sketch of the proof of technical lemma

To better illustrate the proof, we first present an alternative proof for type I points.

We construct the nullhomotopy gradually in the limit space through a sequence of refining skeletons. By controlling the extensions on the new skeletons at each step, these maps on the skeletons converge uniformly to a continuous map defined on the disk.



A very brief sketch of the proof of technical lemma

Sketch: Roughly speaking, if a sub-triangle is away from a point of type III, we can directly contract this sub-triangle; if not, we will use the local 1-contractibility from the sequence to extend the map on a finer 1-skeleton.

