

# Math 470 Assignment 8

Arnold Jiadong Yu

February 10, 2018

6.4.3. Using any test covered in this chapter so far, find out which of the following series converge absolutely, which converge conditionally, and which diverge.

a)  $\sum_{k=1}^{\infty} \frac{(-1)^k k^3}{(k+1)!}$

proof: It converges absolutely. Let  $a_k = \frac{(-1)^k k^3}{(k+1)!}$ , by Ratio Test

$$\lim_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} = \lim_{k \rightarrow \infty} \frac{\frac{(k+1)^3}{(k+2)!}}{\frac{k^3}{(k+1)!}} = \lim_{k \rightarrow \infty} \frac{(k+1)^3}{k^4 + 2k^3} = 0 < 1$$

by L'Hospital's Rule. Hence,  $\sum_{k=1}^{\infty} \frac{(-1)^k k^3}{(k+1)!}$  converges absolutely.

b)  $\sum_{k=1}^{\infty} \frac{(-1)(-3)\dots(1-2k)}{1 \cdot 4 \dots (3k-2)}$

proof: It converges absolutely. Let  $a_k = \frac{(-1)(-3)\dots(1-2k)}{1 \cdot 4 \dots (3k-2)}$ . By Ratio Test,

$$\lim_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} = \lim_{k \rightarrow \infty} \left| \frac{1 - 2(k+1)}{3(k+1) - 2} \right| = \lim_{k \rightarrow \infty} \left| \frac{-2k - 1}{3k + 1} \right| = \frac{2}{3} < 1$$

Hence,  $\sum_{k=1}^{\infty} \frac{(-1)(-3)\dots(1-2k)}{1 \cdot 4 \dots (3k-2)}$  converges absolutely.

c)  $\sum_{k=1}^{\infty} \frac{(k+1)^k}{p^k k!}, p > e$

proof: It converges absolutely. Let  $a_k = \frac{(k+1)^k}{p^k k!}$ . By Ratio Test,

$$\lim_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} = \lim_{k \rightarrow \infty} \left| \frac{\frac{(k+2)^{k+1}}{p^{k+1}(k+1)!}}{\frac{(k+1)^k}{p^k k!}} \right| = \lim_{k \rightarrow \infty} \frac{1}{p} \left( \frac{k+2}{k+1} \right)^{k+1} = \frac{e}{p} < 1$$

by  $p > e$ . Hence  $\sum_{k=1}^{\infty} \frac{(k+1)^k}{p^k k!}$  converges absolutely for  $p > e$ .

$$d) \sum_{k=1}^{\infty} \frac{(-1)^{k+1} \sqrt{k}}{k+1}$$

proof: It converges conditionally. Let  $f(x) = \frac{\sqrt{x}}{x+1}$ , then  $f'(x) = \frac{1-x}{2\sqrt{x}(x+1)^2} < 0$  for all  $x > 1$ . Thus  $f(x) \downarrow 0$  as  $x \rightarrow \infty$ , it converges by Alternating Series Test. Let  $a_k = \frac{\sqrt{k}}{k+1} > \frac{1}{k}$  for large  $k$ , it diverges by Comparison Test. Hence it converges conditionally.

$$e) \sum_{k=1}^{\infty} \frac{(-1)^k \sqrt{k+1}}{\sqrt{k} k^k}$$

proof: It converges absolutely. Considering series  $\sum_{k=1}^{\infty} \frac{\sqrt{k+1}}{\sqrt{k} k^k}$  and let  $a_k = \frac{\sqrt{k+1}}{\sqrt{k} k^k}$  and  $b_k = \frac{1}{k^k}$ , then

$$\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = \lim_{k \rightarrow \infty} \frac{\sqrt{k+1}}{\sqrt{k}} = 1.$$

Hence  $\sum_{k=1}^{\infty} \frac{(-1)^k \sqrt{k+1}}{\sqrt{k} k^k}$  converges absolutely by Limit Comparison Test.

6.4.4. [Abel's Test] Suppose that  $\sum_{k=1}^{\infty} a_k$  converges and that  $b_k \downarrow b$  as  $k \rightarrow \infty$ . Prove that  $\sum_{k=1}^{\infty} a_k b_k$  converges.

proof: Suppose that  $\sum_{k=1}^{\infty} a_k = A \in \mathbb{R}$ ,  $b_k \downarrow b$  as  $k \rightarrow \infty$ , then  $b_k - b \downarrow 0$  as  $k \rightarrow \infty$ . By Dirichlet's Test,  $\sum_{k=1}^{\infty} a_k (b_k - b)$  converges. Let  $\sum_{k=1}^{\infty} a_k (b_k - b) = B \in \mathbb{R}$ , and  $\sum_{k=1}^{\infty} a_k (b) = Ab$ . Thus  $\sum_{k=1}^{\infty} a_k (b) + \sum_{k=1}^{\infty} a_k (b_k - b) = \sum_{k=1}^{\infty} a_k b_k = Ab + B$  converges by Theorem 6.10.

Q: Does  $\sum_{k=1}^{\infty} (\frac{\pi}{2} - \arctan k)$  converges or diverges?

proof: Let  $\tan \alpha = x \in \mathbb{R}$ , then  $\arctan x = \alpha$ . This implies  $\frac{\pi}{2} - \arctan x = \frac{\pi}{2} - \alpha$ . By Cofunction Identities of Trigonometry

$$\tan \alpha = \cot(\frac{\pi}{2} - \alpha) \Rightarrow \frac{1}{\tan \alpha} = \frac{1}{\cot(\frac{\pi}{2} - \alpha)} \Rightarrow \tan(\frac{\pi}{2} - \alpha) = \frac{1}{x}.$$

Thus we are looking at convergence or divergence of  $\sum_{k=1}^{\infty} \arctan \frac{1}{k}$ . Let  $f(x) = \arctan \frac{1}{x}$  and  $g(x) = \frac{1}{x}$ , then  $f'(x) = \frac{-1}{x^2+1}$  and  $g'(x) = \frac{-1}{x^2}$ . Thus  $f'(x) = g'(x)$  for large  $x$ . This implies  $\arctan \frac{1}{x}$  has same behavior as  $\frac{1}{x}$  for

large  $x$ . Hence, by Comparison Test and Harmonic Series Test.  $\sum_{k=1}^{\infty} (\frac{\pi}{2} - \arctan k)$  diverges.