


Why pseudobands works?

Jinyuan Wu

December 14, 2023

A review at *GW*

GW in momentum space

To be very concise... $-i\Sigma \approx GW^{\text{RPA}} =$  .

Assuming quasiparticles... $v(\mathbf{q}+\mathbf{G}) = \frac{4\pi}{V|\mathbf{q}+\mathbf{G}|^2} \cdot \Sigma^{\text{contour}} = \Sigma^{\text{CH}} + \Sigma^{\text{SX}},$

$$\epsilon_{\mathbf{G}\mathbf{G}'}(\mathbf{q}, \omega) = \delta_{\mathbf{G}\mathbf{G}'} - v(\mathbf{q}+\mathbf{G})\chi_{\mathbf{G}\mathbf{G}'}(\mathbf{q}, \omega), \quad M_{nn'}(\mathbf{k}, \mathbf{q}, \mathbf{G}) = \langle n\mathbf{k} + \mathbf{q} | e^{i(\mathbf{q}+\mathbf{G}) \cdot \mathbf{r}} | n'\mathbf{k} \rangle$$

$$\begin{aligned} \langle n\mathbf{k} | \Sigma^{\text{SX}}(\omega) | n'\mathbf{k} \rangle &= - \sum_{n''}^{\text{occ}} \sum_{\mathbf{q}, \mathbf{G}, \mathbf{G}'} M_{n''n}^*(\mathbf{k}, -\mathbf{q}, -\mathbf{G}) M_{n''n'}(\mathbf{k}, -\mathbf{q}, -\mathbf{G}') \\ &\quad \times \epsilon_{\mathbf{G}\mathbf{G}'}^{-1}(\mathbf{q}, \omega - E_{n'', \mathbf{k}-\mathbf{q}}) v(\mathbf{q} + \mathbf{G}'), \end{aligned}$$

$$\begin{aligned} \langle n\mathbf{k} | \Sigma^{\text{CH}}(\omega) | n'\mathbf{k} \rangle &= \frac{i}{2\pi} \sum_{n''} \sum_{\mathbf{q}, \mathbf{G}, \mathbf{G}'} M_{n''n}^*(\mathbf{k}, -\mathbf{q}, -\mathbf{G}) M_{n''n'}(\mathbf{k}, -\mathbf{q}, -\mathbf{G}') \\ &\quad \times \int_0^\infty d\omega' \frac{[\epsilon_{\mathbf{G}\mathbf{G}'}^r]^{-1}(\mathbf{q}, \omega') - [\epsilon_{\mathbf{G}\mathbf{G}'}^a]^{-1}(\mathbf{q}, \omega')}{\omega - E_{n\mathbf{k}} - \omega' + i0^+ \text{sgn}(E_{n\mathbf{k}})} v(\mathbf{q} + \mathbf{G}'). \end{aligned}$$

$$\begin{aligned} \chi_{\mathbf{G}\mathbf{G}'}^{r/a}(\mathbf{q}, \omega) &= \sum_{\mathbf{k}} \sum_n^{\text{occ}} \sum_{n'}^{\text{emp}} M_{nn'}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}') \\ &\quad \times \left(\frac{1}{\omega + E_{n, \mathbf{k}+\mathbf{q}} - E_{n'\mathbf{k}} \pm i0^+} + \frac{1}{-\omega + E_{n, \mathbf{k}+\mathbf{q}} - E_{n'\mathbf{k}} \mp i0^+} \right). \end{aligned}$$

GW in momentum space

Note that χ and $\Sigma_{nn'}^{\text{CH}}$ involve summation over high energy states (with or without GPP)

One bottleneck of GW in momentum space

The problem of summing over many empty bands

One further simplification trick: pseudobands. For $E_{nk} \gg E_F$:

$$\{\phi_{nk}\}_{\text{adjacent energy block } n_b} \rightarrow \sum_{n \in \text{block } n_b} \phi_{nk},$$
$$\{E_{nk}\}_{\text{adjacent energy block } n_b} \rightarrow \frac{1}{|n_b|} \sum_{n \in \text{block } n_b} E_{nk}.$$

4000 bands $\rightarrow \sim 400$ to ~ 1000 bands.

...but is this snake oil? It makes no sense!!!

pseudobands, **and when it works**

Understanding pseudobands: energy averaging

Observation

- $\chi \sim MM^* \times \text{some-function}(E_{\text{emp}} - E_{\text{occ}})$;
- $\Sigma^{\text{CH}} \sim MM^* \times \text{some-function}(\omega - E)$, $\omega \sim E_{\text{F}}$.



👉 For $E_{n\mathbf{k}} \gg E_{\text{F}}$ (ω): energy-dependent factors $\sim \text{const.}$ for all \mathbf{k} .

Thus for both χ and Σ^{CH} involving summation over empty bands:

high emp. bands

$$\sum_{n''} M_{n''n}^* M_{n''n'} \times f(E_{n''\mathbf{k}}) = \sum_{\text{block } n_b} f(\bar{E}_{n_b}) \sum_{n''}^{\text{block } n_b} M_{n''n}^* M_{n''n'}.$$



Understanding pseudobands: energy averaging



Example High-energy DFT bands of WTe_2 monolayer (ONCV-SG15, 120 electrons; 80 Ry cutoff, $20 \times 20 \times 1$ grid). x axis = \mathbf{k} index in irreducible 1BZ; fastest varying coordinate = k_y

👉 $n \sim 1000$: bands are generally flat

Understanding pseudobands: energy averaging



Example High-energy DFT bands of WTe_2 monolayer (ONCV-SG15, 120 electrons; 80 Ry cutoff, $20 \times 20 \times 1$ grid). x axis = \mathbf{k} index in irreducible 1BZ; fastest varying coordinate = k_y

👉 High-lying bands are more dispersive but since they are high, $1/(E_{\text{emp}} - E_{\text{occ}})$ is still flat

Understanding pseudobands: wave function averaging?

For pseudobands to work, we need

$$\underbrace{\sum_{n'' \in \text{block } n_b} M_{n''n}^* M_{n''n'}}_{\text{normal}} \sim M_{\text{averaged band},n}^* M_{\text{averaged band},n'} \quad (1)$$
$$= \underbrace{\sum_{n_1'', n_2'' \in \text{block } n_b} M_{n_1''n'} M_{n_2''n}^*}_{\text{pseudobands}}$$

The question: is $[M_{n_1''n'} M_{n_2''n}^*]_{n_1''n_2''}$ **diagonal?** Easy to verify: it's not.

Then in which case is (1) correct in some sense?

Technical issues in calculating $M_{nn'}(\mathbf{k}, \mathbf{q}, \mathbf{G})$

The structure of $\phi_{n\mathbf{k}}$

Plane wave basis In BerkeleyGW WFN.h5,

$$\phi_{n\mathbf{k}}(\mathbf{r}, \sigma) = \frac{1}{\sqrt{V}} \sum_{\mathbf{G}} e^{i(\mathbf{k}+\mathbf{G})\cdot\mathbf{r}} c_{n\mathbf{k},\mathbf{G}\sigma}.$$

Thus

$$M_{nn'}(\mathbf{k}, \mathbf{q}, \mathbf{G}) = \langle n\mathbf{k} + \mathbf{q} | e^{i(\mathbf{q}+\mathbf{G})\cdot\mathbf{r}} | n'\mathbf{k} \rangle = \sum_{\mathbf{G}', \sigma} c_{n\mathbf{k}+\mathbf{q}, \mathbf{G}+\mathbf{G}'\sigma}^* c_{n\mathbf{k}, \mathbf{G}'\sigma}.$$

Cutoff Each \mathbf{k} has its own \mathbf{G} -grid (~ 30000 vectors for 80 Ry).

Procedure

Input

- indices of \mathbf{k}, \mathbf{q} in \mathbf{k} -grid;
- index of \mathbf{G} in \mathbf{G} -grid of \mathbf{k} (expect a \mathbf{G} in GW \mathbf{G} -grid, cutoff = say 30 Ry, not 80 Ry);
- n, n' .

Procedure

1. find index of \mathbf{k}
2. find index of $\mathbf{G} + \mathbf{G}'$ in \mathbf{G} -grid of $\mathbf{k} + \mathbf{q}$, for each \mathbf{G}' in \mathbf{G} -grid of \mathbf{k}
3. do summation $\sum_{\mathbf{G}', \sigma} c_{n\mathbf{k}+\mathbf{q}, \mathbf{G}+\mathbf{G}'}^* c_{n\mathbf{k}, \mathbf{G}'} \sigma$.

Performance Main bottleneck: finding $\mathbf{G} + \mathbf{G}'$. Using `StaticArrays.jl` helps a lot!

pseudobands **for** χ

Under GPP:

$$\chi_{\mathbf{G}\mathbf{G}'}(\mathbf{q}, \omega = 0)^{\text{high band terms}} \approx \sum_{\mathbf{k}} \sum_{\text{block } n_b}^{\text{emp}} \frac{2}{E_{n,\mathbf{k}+\mathbf{q}} - E_{\text{average in block } n_b}} \\ \times \sum_{n' \in \text{block } n_b} \sum_n^{\text{occ}} M_{nn'}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$$


Our goal *Finding how diagonal is*

$$\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$$

It should be diagonal for pseudobands to work.

Numerical characterization of $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$


Case 1: $\mathbf{G} = \mathbf{G}'$ In this case $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ is large and fairly diagonal



Parameters $\mathbf{G} = (0, 1, -14)$, $\mathbf{k} = \mathbf{k}_2 = (0, 0.00, 0)$, $\mathbf{q} = (0, 0.00, 0)$

Numerical characterization of $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$


Case 1: $\mathbf{G} = \mathbf{G}'$ In this case $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ is large and fairly diagonal



Parameters $\mathbf{G} = (0, 1, -14)$, $\mathbf{k} = \mathbf{k}_2 = (0, 0.05, 0)$, $\mathbf{q} = (0, 0.10, 0)$

Numerical characterization of $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$


Case 1: $\mathbf{G} = \mathbf{G}'$ In this case $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ is large and fairly diagonal



Strikingly, $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ is still very diagonal for bands near Fermi surface!

Numerical characterization of $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$

Case 2: $\mathbf{G} \neq \mathbf{G}'$ $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ is very non-diagonal, but since the terms's random phases cancel each other so the overall sum after $\sum_{n'}^{\text{emp}}$ is small




Half-way generalization about pseudobands in χ

- 👉 pseudobands works when it's necessary to do so
- 👉 What prevents pseudobands from working around Fermi surface is the energy dispersion

...but do we have any theoretical explanation for this?

Cancellation of cross-terms in χ

$M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$ and n (occupied band index), when $n'_1 \neq n'_2$:



Although for a single n , the value can be large, as we sum over n the terms cancel each other \Rightarrow diagonal $\sum_n^{\text{occ}} M_{nn'_1}(\mathbf{k}, \mathbf{q}, \mathbf{G}) M_{nn'_2}^*(\mathbf{k}, \mathbf{q}, \mathbf{G}')$

pseudobands **for** Σ

Only Σ^{CH} involves Σ^{emp} :

$$\langle n\mathbf{k} | \Sigma^{\text{CH, GPP}} | n'\mathbf{k} \rangle = \frac{1}{2} \sum_{n''} \sum_{\mathbf{q}\mathbf{G}\mathbf{G}'} M_{n''n}^*(\mathbf{k}, -\mathbf{q}, -\mathbf{G}) M_{n''n'}(\mathbf{k}, -\mathbf{q}, -\mathbf{G}') \times$$


something about $\mathbf{q}, \mathbf{k}, \mathbf{G}, \mathbf{G}'$

Problems

- No summation over occupied states \Rightarrow ~~X~~the aforementioned cancellation mechanism
- Summation over \mathbf{G} very complicated \Rightarrow analysis based on \mathbf{G} very hard.

Some tentative ideas

- It has been verified that a single $M_{n_1''n}^* M_{n_2''n}$ term is never diagonal
- But it seems naively summing over $\mathbf{G} = \mathbf{G}'$ (without considering the weight factor) makes it diagonal. . .



Some tentative ideas

- $M_{n_1''n}^*(\mathbf{G}_1)M_{n_2''n}(\mathbf{G}_2)$ is not diagonal at all when $\mathbf{G}_1 \neq \mathbf{G}_2$;
- but probably only the $\mathbf{G}_1 = \mathbf{G}_2$ terms matter?

Comments from Felipe H. Jornada's recent paper

“Stochastic” pseudobands

Energy averaging is also done in their paper:

$$G = \sum_{n \text{ near F.S.}} \frac{|\phi_{n\mathbf{k}}\rangle\langle\phi_{n\mathbf{k}}|}{\omega - E_{n\mathbf{k}}} + \sum_{\text{pseudobands block}} \frac{1}{\omega - \bar{E}} \sum_{n \in \text{block}} |\phi_{n\mathbf{k}}\rangle\langle\phi_{n\mathbf{k}}|.$$

Their procedure Replace each pseudobands block by *several bands*:

$$\{\phi_{n\mathbf{k}}\}_{n \text{ in block}} \longrightarrow \left\{ |\phi_{\xi\mathbf{k}}\rangle = \sum_{n \text{ in block}} e^{i\theta_{n\xi}} |\phi_{n\mathbf{k}}\rangle \right\}_{\xi}$$

Justification When $N_{\xi} \rightarrow \infty$, $\langle e^{-i\theta_{n'\xi}} e^{i\theta_{n\xi}} \rangle_{\xi} = \delta_{nn'}$

Comparison with the current script When $N_{\xi} = 1$, $e^{i\theta_{n\xi}}$ comes from DFT diagonalization

FHJ's justification of pseudobands

Convergence conditions When $N_{\text{pseudobands blocks}} \rightarrow \infty$ and in each block $N_{\xi} \rightarrow \infty$,

- the expectations of χ and Σ goes to the true value;
- the standard error goes to zero.

Is this result enough as a justification? My answer: no; because the limits = not doing pseudobands at all. (And their recommended values are not large enough anyway.)

FHJ's justification of pseudobands

Convergence conditions When $N_{\text{pseudobands blocks}} \rightarrow \infty$ and in each block $N_{\xi} \rightarrow \infty$,

- the expectations of χ and Σ goes to the true value;
- the standard error goes to zero.

Is this result enough as a justification? My answer: no; because the limits = not doing pseudobands at all. (And their recommended values are not large enough anyway.)

Additional justification “the polarizability tends to converge much faster than G , partially due to the *rapidly oscillating nature* of the matrix elements involving Kohn-Sham states used in the evaluation of the polarizability”

Compressing the bands even harder: size of blocks

Recall that band dispersion is the main problem...

$$\Delta\chi \sim \frac{\Delta E}{E^2}, \quad \chi \sim \frac{1}{E} \Rightarrow \frac{\Delta\chi}{\chi} \simeq \frac{\Delta E}{E}$$
$$\Rightarrow \frac{\Delta\chi}{\chi} \lesssim \text{const} \Leftrightarrow \boxed{\frac{\Delta E}{E} \lesssim \text{const}}$$

Exponential growth of the energy spread of blocks!

Compressing the bands even harder: pseudobands near E_F ?

They claim that for ϵ , bands near Fermi surface can also be pseudo-ized

It somehow goes against my own numerical experiments; but they are quite sure about that hmm. . .

They also claim that for σ , only the bands near Fermi surface shouldn't be pseudo-ized.

Important claim: protection window is not a convergence parameter.