

Homework 3

Jinyuan Wu

November 2, 2022

Problem 2

Solution

(a) Repeating the procedure used in ordinary superfluid, we do the decomposition

$$\varphi = \sqrt{\rho} e^{i\theta} = \sqrt{\rho_0 + \delta\rho} e^{i\theta}, \quad (1)$$

and therefore

$$-\frac{\varphi^* \nabla^2 \varphi}{2m} = \frac{\rho}{2m} (\nabla \theta)^2 + \frac{(\nabla \rho)^2}{8\rho m}, \quad (2)$$

$$\varphi^* \partial_\tau \varphi = \underbrace{\frac{1}{2} \partial_\tau \rho}_{\text{time derivative, ignored}} + i\rho \partial_\tau \theta, \quad (3)$$

$$|\varphi(\mathbf{x})| U(\mathbf{x} - \mathbf{y}) |\varphi(\mathbf{y})| = \rho(\mathbf{x}) U(\mathbf{x} - \mathbf{y}) \rho(\mathbf{y}), \quad (4)$$

the theory is now

$$S = \int d\tau \left(\int d^d \mathbf{x} \left(i\rho \partial_\tau \theta + \frac{\rho}{2m} (\nabla \theta)^2 + \frac{(\nabla \rho)^2}{8\rho m} - \mu\rho \right) + \frac{1}{2} \int d^d \mathbf{x} \int d^d \mathbf{y} \rho(\mathbf{x}) U(\mathbf{x} - \mathbf{y}) \rho(\mathbf{y}) \right). \quad (5)$$

Around the ground state, we have (note that since we are around a saddle point, the sum of all terms containing $\delta\rho$ only is always zero; the resulting theory has the form of $c_1 \delta\rho \partial_\tau \theta + c_2 \delta\rho^2$; the chemical potential term is therefore missing in the theory around the saddle point)

$$i\rho \partial_\tau \theta = \underbrace{i\rho_0 \partial_\tau \theta}_{\text{time derivative}} + i\delta\rho \partial_\tau \theta,$$

and since $\nabla \rho = \nabla \delta\rho$, we have

$$\frac{(\nabla \rho)^2}{8\rho m} \approx \frac{(\nabla \delta\rho)^2}{8\rho_0 m},$$

ignoring the fluctuation of the ρ in the denominator. Similarly, since we are working on a low energy theory, the fluctuation of θ shouldn't be large, and we have

$$\frac{\rho}{2m} (\nabla \theta)^2 \approx \frac{\rho_0}{2m} (\nabla \theta)^2.$$

The theory is then

$$S = \int d^{d+1}x \left(\frac{\rho_0}{2m} (\nabla \theta)^2 + i\delta\rho \partial_\tau \theta + \frac{(\nabla \delta\rho)^2}{8\rho_0 m} + \frac{1}{2} \delta\rho(\mathbf{x}) \int d^d \mathbf{y} U(\mathbf{x} - \mathbf{y}) \delta\rho(\mathbf{y}) \right) + S_{\text{saddle}}. \quad (6)$$

Integrating out $\delta\rho$, we get

$$\begin{aligned} S_{\text{eff}} &= \int d^{d+1}x \frac{\rho_0}{2m} (\nabla \theta)^2 - \frac{1}{2} \int d\tau \int d^d \mathbf{x} d^d \mathbf{y} i\partial_\tau \theta(\mathbf{x}, \tau) \frac{1}{\int d^d \mathbf{y} U(\mathbf{x} - \mathbf{y}) - \frac{1}{4\rho_0 m} \nabla^2} i\partial_\tau \theta(\mathbf{y}, \tau) \\ &= \int d^{d+1}x \frac{\rho_0}{2m} (\nabla \theta)^2 + \frac{1}{2} \int d\tau \int d^d \mathbf{x} d^d \mathbf{y} \partial_\tau \theta(\mathbf{x}, \tau) G(\mathbf{x} - \mathbf{y}) \partial_\tau \theta(\mathbf{y}), \end{aligned} \quad (7)$$

where

$$\int d^d \mathbf{y} U(\mathbf{x} - \mathbf{y}) G(\mathbf{y} - \mathbf{z}) - \frac{1}{4\rho_0 m} \nabla_{\mathbf{x}}^2 G(\mathbf{x} - \mathbf{z}) = \delta(\mathbf{x} - \mathbf{z}). \quad (8)$$

Similar to the procedure in dealing with ordinary superfluid, since we are only interested in the long wave length behaviors of θ , the ∇^2 term can be thrown away, and we have

$$\begin{aligned}\int \frac{d^d \mathbf{p}}{(2\pi)^d} e^{i\mathbf{p} \cdot (\mathbf{x} - \mathbf{z})} &= \int \frac{d^d \mathbf{p}}{(2\pi)^d} \int d^d \mathbf{y} U(\mathbf{x} - \mathbf{y}) G(\mathbf{p}) e^{i\mathbf{p} \cdot (\mathbf{y} - \mathbf{z})} \\ &= \int \frac{d^d \mathbf{p}}{(2\pi)^d} \int d^d \mathbf{r} U(\mathbf{r}) e^{-i\mathbf{p} \cdot \mathbf{r}} G(\mathbf{p}) e^{i\mathbf{p} \cdot (\mathbf{x} - \mathbf{z})} \quad (\mathbf{r} = \mathbf{x} - \mathbf{y}),\end{aligned}$$

so

$$G(\mathbf{r}) = \int \frac{d^d \mathbf{p}}{(2\pi)^d} e^{i\mathbf{p} \cdot \mathbf{r}} G(\mathbf{p}), \quad G(\mathbf{p}) = \frac{1}{U(\mathbf{p})} = \frac{1}{\int d^d \mathbf{r} U(\mathbf{r}) e^{-i\mathbf{p} \cdot \mathbf{r}}}. \quad (9)$$

To evaluate $G(\mathbf{p})$, we need to find

$$U(\mathbf{p}) = \int_0^\infty dr \frac{2\pi^{d/2}}{\Gamma(d/2)} r^{d-1} \frac{U_0}{r^{d-\epsilon}} \quad (10)$$

Problem 3

Solution

(a) The energy now can be exactly evaluated (N is the number of sites):

$$E = \frac{UN}{2}(M^2 - M) - \mu NM = \frac{N}{2}(UM^2 - (U + 2\mu)M). \quad (11)$$

At the ground state, E is minimized, and we have

$$M = \frac{U + 2\mu}{2U}, \quad (12)$$

and the energy gap is

$$\Delta E = E|_{M+1} - E|_M = \frac{N}{2}(2UM - 2\mu) = \frac{1}{2}NU. \quad (13)$$