

Electron Gas with Coulomb Interaction

Jinyuan Wu

January 21, 2022

It is said that for a simple demonstration of what happens in a metal, we can only work with the **jellium model**, ignoring the details of the lattice, which means we can just work with a non-relativistic electron gas with Coulomb interaction. This article is an extension of Section 6.2 in [this solid state physics note](#). We will discuss some early development of RPA.

Zhengzhong
Li, Sec. 4.2

1 Notations and basic facts about the jellium model

We define

$$\rho(\mathbf{r}) = \sum_{\sigma} \psi_{\sigma}^{\dagger}(\mathbf{r}) \psi_{\sigma}(\mathbf{r}) = \frac{1}{V} \sum_{\mathbf{k}, \mathbf{k}', \sigma} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}'\sigma} e^{-i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} = \frac{1}{V} \sum_{\mathbf{q}} \sum_{\mathbf{k}, \sigma} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}+\mathbf{q}, \sigma} e^{i\mathbf{q} \cdot \mathbf{r}}, \quad (1)$$

and the commutation relations are

$$[c_{\mathbf{k}}, c_{\mathbf{k}'}^{\dagger}] = \delta_{\mathbf{k}\mathbf{k}'}. \quad (2)$$

We can also write down $\rho(\mathbf{r})$ in the first quantization formulation, i.e.

$$\rho(\mathbf{r}) = \sum_i \delta(\mathbf{r} - \mathbf{r}_i) = \sum_i \frac{1}{V} \sum_{\mathbf{q}} e^{i\mathbf{q} \cdot (\mathbf{r} - \mathbf{r}_i)} = \frac{1}{V} \sum_{\mathbf{q}} \sum_i e^{i\mathbf{q} \cdot (\mathbf{r} - \mathbf{r}_i)}. \quad (3)$$

We have

$$\rho_{\mathbf{q}} := \sum_i e^{-i\mathbf{q} \cdot \mathbf{r}_i} = \sum_{\mathbf{k}, \sigma} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}+\mathbf{q}, \sigma}, \quad \rho(\mathbf{r}) = \frac{1}{V} \sum_{\mathbf{q}} \rho_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{r}}. \quad (4)$$

Easily, we find

$$\rho_{\mathbf{q}=0} = \int d^3\mathbf{r} \rho(\mathbf{r}) = N. \quad (5)$$

The Hamiltonian of electrons is

$$H_{\text{electron}} = \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + \frac{1}{2V} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}, \sigma, \sigma'} c_{\mathbf{k}+\mathbf{q}, \sigma}^{\dagger} c_{\mathbf{k}'-\mathbf{q}, \sigma'}^{\dagger} V(\mathbf{q}) c_{\mathbf{k}'\sigma'} c_{\mathbf{k}\sigma},$$

where

$$V(\mathbf{q}) = \int e^{-i\mathbf{q} \cdot \mathbf{r}} \frac{e^2}{r} = \frac{4\pi e^2}{q^2}. \quad (6)$$

The Hamiltonian of the lattice and the electron-lattice coupling is

$$H_{\text{lattice}} = \frac{1}{2V} \sum_{\mathbf{q}} \rho_{\text{ion}, -\mathbf{q}} V(\mathbf{q}) \rho_{\text{ion}, \mathbf{q}} - \frac{1}{V} \sum_{\mathbf{q}} \rho_{\text{ion}, \mathbf{q}} V(\mathbf{q}) \rho_{\mathbf{q}}.$$

When $|\mathbf{q}| = 0$, $V(\mathbf{q})$ is divergent, but this does not matter. In the jellium model, the only non-zero Fourier component of $\rho_{\text{ion}}(\mathbf{r})$ is $\rho_{\text{ion}, \mathbf{q}=0} = N$ (the solid is neutral so the number of positive charges must be equal to the number of negative charges), and we soon find

$$H_{\text{lattice}} + \frac{1}{2V} \sum_{\mathbf{q}=0, \mathbf{k}, \mathbf{k}', \sigma, \sigma'} c_{\mathbf{k}+\mathbf{q}, \sigma}^{\dagger} c_{\mathbf{k}'-\mathbf{q}, \sigma'}^{\dagger} V(\mathbf{q}) c_{\mathbf{k}'\sigma'} c_{\mathbf{k}\sigma} = 0.$$

So all divergences cancel with each other, and in the end, the total Hamiltonian is

$$\begin{aligned} H &= \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + \frac{1}{2V} \sum_{\mathbf{q} \neq 0} \sum_{\mathbf{k}, \mathbf{k}', \sigma, \sigma'} c_{\mathbf{k}+\mathbf{q}, \sigma}^{\dagger} c_{\mathbf{k}'-\mathbf{q}, \sigma'}^{\dagger} V(\mathbf{q}) c_{\mathbf{k}'\sigma'} c_{\mathbf{k}\sigma} \\ &= \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + \frac{1}{2V} \sum_{\mathbf{q} \neq 0} \rho_{\mathbf{q}}^{\dagger} V(\mathbf{q}) \rho_{\mathbf{q}} = \sum_{\mathbf{k}, \sigma} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + \frac{1}{2V} \sum_{\mathbf{q} \neq 0} \rho_{-\mathbf{q}} V(\mathbf{q}) \rho_{\mathbf{q}}. \end{aligned} \quad (7)$$

The contribution of the positive ion “jell” both constrains the electrons in the solid (or in other words, give a chemical potential) and regularize the singularity of $V(\mathbf{q} = 0)$.

Note that (7) differs with

$$H = \sum_i \frac{\mathbf{p}_i^2}{2m} + \frac{1}{2} \sum_{i \neq j} \frac{e^2}{|\mathbf{r}_i - \mathbf{r}_j|} \quad (8)$$

with the $i = j$ term. This is an infinite constant and does not matter.

2 Classical (or first quantized) theory of the collective oscillation of electrons

From (4), we have

Zhengzhong

Li, Sec. 4.3

$$\begin{aligned} \dot{\rho}_{\mathbf{q}} &= \sum_j (-i\mathbf{q}) \cdot \mathbf{v}_j e^{-i\mathbf{q} \cdot \mathbf{r}_j}, \\ \ddot{\rho}_{\mathbf{q}} &= \sum_j (-i\mathbf{q} \cdot \dot{\mathbf{v}}_j - (\mathbf{q} \cdot \mathbf{v}_j)^2) e^{-i\mathbf{q} \cdot \mathbf{r}_j}. \end{aligned} \quad (9)$$

Now $\dot{\mathbf{v}}_j$ can be derived from (7):

$$\begin{aligned} m\dot{\mathbf{v}}_j &= -\nabla_j \frac{1}{2V} \sum_{\mathbf{q} \neq 0} V(\mathbf{q}) \rho_{\mathbf{q}}^\dagger \rho_{\mathbf{q}} \\ &= -\frac{1}{2V} \nabla_j \sum_{\mathbf{q} \neq 0} V(\mathbf{q}) \sum_{i,k} e^{i\mathbf{q} \cdot (\mathbf{r}_i - \mathbf{r}_k)} \\ &= -\frac{1}{2V} \sum_{\mathbf{q} \neq 0} V(\mathbf{q}) \sum_k (i\mathbf{q}) e^{i\mathbf{q} \cdot (\mathbf{r}_j - \mathbf{r}_k)} + \sum_i (-i\mathbf{q}) e^{i\mathbf{q} \cdot (\mathbf{r}_j - \mathbf{r}_i)} \\ &= -\frac{1}{V} \sum_{\mathbf{q} \neq 0} i\mathbf{q} V(\mathbf{q}) \sum_i e^{i\mathbf{q} \cdot (\mathbf{r}_j - \mathbf{r}_i)} \\ &= -\frac{4\pi e^2}{V} \sum_{\mathbf{q} \neq 0} \frac{i\mathbf{q}}{q^2} e^{i\mathbf{q} \cdot \mathbf{r}_j} \rho_{\mathbf{q}}, \end{aligned}$$

and from (9), we have

$$\begin{aligned} \ddot{\rho}_{\mathbf{q}} &= -\sum_j \frac{4\pi e^2}{mV} \sum_{\mathbf{q}' \neq 0} \frac{\mathbf{q} \cdot \mathbf{q}'}{q'^2} e^{i\mathbf{q}' \cdot \mathbf{r}_j} \rho_{\mathbf{q}'} e^{-i\mathbf{q} \cdot \mathbf{r}_j} - \sum_j (\mathbf{q} \cdot \mathbf{v}_j)^2 e^{-i\mathbf{q} \cdot \mathbf{r}_j} \\ &= -\frac{4\pi e^2}{mV} \sum_{\mathbf{q}' \neq 0} \rho_{\mathbf{q}'} \rho_{\mathbf{q}-\mathbf{q}'} - \sum_j (\mathbf{q} \cdot \mathbf{v}_j)^2 e^{-i\mathbf{q} \cdot \mathbf{r}_j}. \end{aligned}$$

Now we make the **random phase approximation (RPA)** in its original form: We assume that only the $\mathbf{q} = \mathbf{q}'$ term in the first term is important, because in the high density limit, there are no position preference of electrons (when the density is low, there might be a Wigner crystal, and RPA fails), and when $\mathbf{q} \neq 0$, both $\rho_{\mathbf{q}'}$ and $\rho_{\mathbf{q}-\mathbf{q}'}$ are sums of almost random phase factors $e^{-i\mathbf{q} \cdot \mathbf{r}_j}$, and therefore are both small enough. So we get the EOM after RPA:

$$\begin{aligned} \ddot{\rho}_{\mathbf{q}} &= -\frac{4\pi e^2}{mV} \rho_{\mathbf{q}} \rho_0 - \sum_j (\mathbf{q} \cdot \mathbf{v}_j)^2 e^{-i\mathbf{q} \cdot \mathbf{r}_j} \\ &= -\frac{4\pi e^2 n}{m} \rho_{\mathbf{q}} - \sum_j (\mathbf{q} \cdot \mathbf{v}_j)^2 e^{-i\mathbf{q} \cdot \mathbf{r}_j}, \end{aligned} \quad (10)$$

where $n = N/V$ is the jellium density. Section (5.4.2) in [this solid state physics note](#) tells us that the electron-hole pair excitations are gapless, but from the EOM above we soon find that in the $\mathbf{q} \rightarrow 0$ case there is a finite ω solution, which is given by

$$\ddot{\rho}_{\mathbf{q}} + \omega_p^2 \rho_{\mathbf{q}} = 0, \quad \omega_p^2 = \frac{4\pi e^2 n}{m}. \quad (11)$$

We see that this term arises from the Coulomb interaction. In other words, long range interaction induces a collective modes in the metal, which is now known as **plasmon**. We know it is plasmon, or quantized plasma oscillation, because our derivation above also works for the oscillation of negative charges around positive charges in a plasma.

3 Second quantized EOM of density modes

Now we try to derive plasmon in the second quantization EOM framework. Generally speaking, a generalized hydrodynamic mode in an electron gas is made of a linear combination of

Zhengzhong
Li, Sec. 4.4

$$\rho_{\mathbf{k}\mathbf{q}}^\dagger =: \sum_{\sigma} c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{k}\sigma}. \quad (12)$$

By evaluating the EOM, we may be able to identify stable density modes.

We start from the free theory. It can be verified that

$$[c_1^\dagger c_2, c_3^\dagger c_4] = \delta_{23} c_1^\dagger c_4 - \delta_{14} c_3^\dagger c_2, \quad (13)$$

and therefore we have

$$\begin{aligned} i\dot{\rho}_{\mathbf{k}\mathbf{q}}^\dagger &= [\rho_{\mathbf{k}\mathbf{q}}^\dagger, H_0] = \sum_{\mathbf{p},\alpha} \frac{\mathbf{p}^2}{2m} \sum_{\sigma} [c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{k}\sigma}, c_{\mathbf{p}\alpha}^\dagger c_{\mathbf{p}\alpha}] \\ &= \sum_{\mathbf{p},\alpha,\sigma} \frac{\mathbf{p}^2}{2m} (c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{p}\alpha} \delta_{\mathbf{k}\mathbf{p}} \delta_{\sigma\alpha} - c_{\mathbf{p}\alpha}^\dagger c_{\mathbf{k}\sigma} \delta_{\mathbf{k}+\mathbf{q},\mathbf{p}} \delta_{\sigma\alpha}) \\ &= \sum_{\sigma} \left(\frac{\mathbf{k}^2}{2m} c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{k}\sigma} - \frac{(\mathbf{q}+\mathbf{k})^2}{2m} c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{k}\sigma} \right), \end{aligned}$$

so

$$i\dot{\rho}_{\mathbf{k}\mathbf{q}}^\dagger = [\rho_{\mathbf{k}\mathbf{q}}^\dagger, H_0] = -\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} \rho_{\mathbf{k}\mathbf{q}}^\dagger, \quad \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} = \frac{(\mathbf{q}+\mathbf{k})^2}{2m} - \frac{\mathbf{k}^2}{2m}. \quad (14)$$

Therefore, we find there are electron-hole pairs in the free theory. In the last section we find that in a plasmon mode, we have $\rho_{\mathbf{k}} = \sum_{\mathbf{q}} \rho_{\mathbf{k}\mathbf{q}} \sim e^{-i\omega_{\mathbf{p}} t}$, and $\omega_{\mathbf{q}=0} \neq 0$, but here when $\mathbf{q} = 0$, $\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}}$ vanishes, so there is no plasmon. This is expected, since in the last section we find the plasmon comes from the Coulomb interaction.

Now we move on to discuss the jellium model. We have

$$\begin{aligned} [\rho_{\mathbf{k}\mathbf{q}}^\dagger, H] &= [\rho_{\mathbf{k}\mathbf{q}}^\dagger, H_0 + H_{\text{Coulomb}}] = -\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} \rho_{\mathbf{k}\mathbf{q}}^\dagger + \frac{1}{2V} \sum_{\mathbf{q}' \neq 0} V(\mathbf{q}') [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{-\mathbf{q}'} \rho_{\mathbf{q}'}] \\ &= -\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} \rho_{\mathbf{k}\mathbf{q}}^\dagger + \frac{1}{2V} \sum_{\mathbf{q}' \neq 0} V(\mathbf{q}') (\rho_{-\mathbf{q}'} [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{\mathbf{q}'}] + [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{-\mathbf{q}'}] \rho_{\mathbf{q}'}) \\ &= -\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} \rho_{\mathbf{k}\mathbf{q}}^\dagger + \frac{1}{2V} \sum_{\mathbf{q}' \neq 0} V(\mathbf{q}') (\rho_{-\mathbf{q}'} [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{\mathbf{q}'}] + [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{\mathbf{q}'}] \rho_{-\mathbf{q}'}), \end{aligned}$$

where

$$\begin{aligned} [\rho_{\mathbf{k}\mathbf{q}}^\dagger, \rho_{\mathbf{q}'}] &= \sum_{\alpha} \sum_{\mathbf{k}',\beta} [c_{\mathbf{k}+\mathbf{q},\alpha}^\dagger c_{\mathbf{k}\alpha}, c_{\mathbf{k}'\beta}^\dagger c_{\mathbf{k}'+\mathbf{q}',\beta}] \\ &= \sum_{\alpha} (c_{\mathbf{k}+\mathbf{q},\alpha}^\dagger c_{\mathbf{k}+\mathbf{q}',\alpha} - c_{\mathbf{k}+\mathbf{q}-\mathbf{q}',\alpha}^\dagger c_{\mathbf{k}\alpha}). \end{aligned}$$

So the EOM of $\rho_{\mathbf{k}\mathbf{q}}^\dagger$ is finally

Zhengzhong
Li (4.4.9)

$$i\dot{\rho}_{\mathbf{k}\mathbf{q}}^\dagger = -\omega_{\mathbf{k}\mathbf{q}}^{\text{pair}} \rho_{\mathbf{k}\mathbf{q}}^\dagger - \frac{1}{2V} \sum_{\mathbf{q}' \neq 0, \alpha} V(\mathbf{q}') \{ \rho_{-\mathbf{q}'} [c_{\mathbf{k}+\mathbf{q}-\mathbf{q}',\alpha}^\dagger c_{\mathbf{k}\alpha} - c_{\mathbf{k}+\mathbf{q},\alpha}^\dagger c_{\mathbf{k}+\mathbf{q}',\alpha}] \}. \quad (15)$$

By taking all possible momenta in (15), we get a closed group of equations about density modes. In principle, this gives the (generalized) hydrodynamic modes in the jellium model

completely, though we all know it is highly difficult to solve EOM of operators. This can also be seen as a successful *bosonization* the theory, though unlike the case in a Luttinger liquid, this does not help us understand the behavior of the jellium model. A possible way to simplify (15) is to assume that $c_{\mathbf{k}+\mathbf{q}-\mathbf{q}',\alpha}^\dagger c_{\mathbf{k}\alpha} - c_{\mathbf{k}+\mathbf{q},\alpha}^\dagger c_{\mathbf{k}+\mathbf{q}',\alpha}$ does not fluctuate much, and we can replace it by the expectation value. if under this approximation we find a stable mode, then we have

$$\begin{aligned} (\omega - \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}}) \rho_{\mathbf{k}\mathbf{q}}^\dagger &= \frac{1}{2V} \sum_{\mathbf{q}' \neq 0, \alpha} V(q') \{ \rho_{-\mathbf{q}', \alpha} \langle c_{\mathbf{k}+\mathbf{q}-\mathbf{q}',\alpha}^\dagger c_{\mathbf{k}\alpha} - c_{\mathbf{k}+\mathbf{q},\alpha}^\dagger c_{\mathbf{k}+\mathbf{q}',\alpha} \rangle \} \\ &= \frac{1}{2V} \sum_{\mathbf{q}' \neq 0, \alpha} 2V(q') \rho_{\mathbf{q}'}^\dagger (n_{\mathbf{k}} \delta_{\mathbf{q}\mathbf{q}'} - n_{\mathbf{k}+\mathbf{q}} \delta_{\mathbf{q}\mathbf{q}'}) \\ &= \frac{2}{V} \rho_{\mathbf{q}}^\dagger (n_{\mathbf{k}} - n_{\mathbf{k}+\mathbf{q}}) V(q), \end{aligned}$$

where

$$n_{\mathbf{k}} := \langle c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} \rangle. \quad (\text{no summation over } \sigma) \quad (16)$$


From the equation above we find

$$\rho_{\mathbf{k}\mathbf{q}}^\dagger = \frac{2V(q)}{V} \rho_{\mathbf{q}}^\dagger \frac{n_{\mathbf{k}} - n_{\mathbf{k}+\mathbf{q}}}{\omega - \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}}}, \quad (17)$$

and summing over \mathbf{k} , we have

$$1 = \frac{2V(q)}{V} \sum_{\mathbf{k}} \frac{n_{\mathbf{k}} - n_{\mathbf{k}+\mathbf{q}}}{\omega - \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}}}. \quad (18)$$

What we are doing here is actually a mean field approximation. It is natural to guess that the approximation corresponds to one Green function resummation strategy, which we will discuss in detail in Section 4. An intuitive way to see what is going on is to note that (17) is actually



$$\text{Bubble diagram with shaded circle} = \text{Vertex diagram} + \text{Bubble diagram with shaded circle and wavy line}, \quad (19)$$

where the $(n_{\mathbf{k}} - n_{\mathbf{k}+\mathbf{q}})/(\omega - \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}})$ factor is “the propagator of an electron-hole pair”. The approximation we have made is also a *random phase approximation*, because after the approximation, all $\mathbf{q}' \neq \mathbf{q}$ terms in (15) disappear.

Now we can easily find the frequencies and the states of all density modes. This also shows a benefit of our operator EOM based approach: in a Green function based approach, we can only easily find the spectrum of the density modes by checking the singularities of $\langle nn \rangle$, but it is not that easy to identify their actual “shape”. We, however, will postpone the solution of (18) to Section 5, since (15) is actually just about the *real* part of the Green function: we can recognize factors like $1/(\omega - \omega_{\mathbf{k}\mathbf{q}}^{\text{pair}})$, which misses the important infinitesimal $i0^+$ imaginary part in the denominator, which can be used to determine the damping rate.

4 The Green function, linear response and Feynman diagram resummation

We have already discussed the density-density Green function in the free theory in (5.58) in [this solid state physics note](#). In this section we will provide a more detailed analysis based on Huaiyu Wang’s book.¹

First we estimate the magnitude of the kinetic energy and the Coulomb repulsion energy. By definition we have

$$\frac{V}{N} = \frac{4}{3} \pi r_0^3,$$

¹Which is accused of plagiarism of Eleftherios’ book about Green functions – but since I have only read the former, I will go on with the section index of the former.

and

$$N = \frac{V}{(2\pi)^3} \cdot \frac{4}{3} \pi k_F^3,$$

and therefore we have

$$r_s =: \frac{r_0}{a_0} = \left(\frac{9}{4} \pi \right)^{1/3} \frac{1}{k_F a_0}, \quad (20)$$

where

$$a_0 = \frac{1}{2me^2} \quad (21)$$

is the Bohr radius. The dimensionless r_s gives the ratio between the average distance between electrons and the Bohr radius. The high density condition $r_s \ll 1$ is therefore equivalent to

$$k_F a_0 \gg 1. \quad (22)$$

Huaiyu
Wang Eq.
(13.3.8)

The kinetic energy of a single electron is about

$$E_K \sim \frac{k_F^2}{2m}, \quad (23)$$

while the potential energy of a single electron is about

$$E_V \sim \frac{1}{r_s a_0}, \quad (24)$$

and the ratio between the two is

$$\frac{E_V}{E_K} \sim \frac{e^2}{r_s a_0} \frac{2m}{k_F^2} = \frac{1}{(k_F a_0)^2 r_s} \ll 1. \quad (25)$$

We reach a quite counter-intuitive result: in a *high* density electron gas, where the Coulomb repulsion is *strong*, the Coulomb energy is just a small perturbation! The explanation can be found in Section 6.1 in [this solid state physics note](#).

Note


We need to be cautious when comparing energies, since we can add an arbitrary large or small constant to any term in the Hamiltonian. What really matters is the *range* in which an energy term can vary. For example, an electron can take any of the momenta in the Fermi sea, and therefore its kinetic energy ranges from 0 to $k_F^2/2m$, and therefore we have (23).

Now we conduct loop diagram corrections. The first-order self-energy correction is done in (6.14) in [this solid state physics note](#), and we find it is finite until we go close to the Fermi surface. This is not really concerning, because if we replace the bare interaction line (“photon line”) by the *screened* version (which we are going to discuss soon), the divergence vanishes. Here we follow the approach in Section 13.3.2 and use a power counting method to estimate the divergence of corrections. It is often the case that a certain type of diagrams all diverges severely, but resummation of them removes the divergence. In our current study on the jellium model, divergence analysis is especially important. Suppose q is the four-momentum of an interaction line (i.e. a “virtual photon line” that carries the Coulomb repulsion), which of course we need to integrate over (since an interaction line cannot be an external line), we will show that only diagrams that are severely divergent in $|q|$ are important with the argument in the discussion around Eq. (5.25) in Altland. We know an interaction line introduces the following component into the diagram:

Huaiyu
Wang
Eq. (13.3.10)

Huaiyu
Wang
Sec. 13.3.2

Altland
Eq. (5.25)



and the two electron propagators result in something like (5.58) in [this solid state physics note](#). We find that when $\mathbf{q} \rightarrow 0$, \mathbf{k} is near the Fermi surface, and q_0 is also approaching zero, $\sum_{\mathbf{k}} G_{\mathbf{k}} G_{\mathbf{q}+\mathbf{k}}$ reaches a peak because there is a step between $f(\xi_{\mathbf{k}})$ and $f(\xi_{\mathbf{k}+\mathbf{q}})$, just as is the case in (6.27) in [this solid state physics note](#). Generally speaking, q_0 will not be zero because for example, we can assume that the two electron lines in the above diagram are external ones, and in this case q_0 is usually non-zero. However, if the diagram has algebraic divergence in \mathbf{q} , again we have something like

$$\frac{1}{|\mathbf{q}|} \sum_{\mathbf{k}} G_{\mathbf{k}} G_{\mathbf{q}+\mathbf{k}} = \frac{2}{V} \sum_{\mathbf{k}} \frac{1}{\omega + \xi_{\mathbf{k}} - \xi_{\mathbf{k}+\mathbf{q}}} \frac{f(\xi_{\mathbf{k}}) - f(\xi_{\mathbf{k}+\mathbf{q}})}{|\mathbf{q}|},$$

which evaluates to something proportion to the density of states at the Fermi surface. The denser the electron gas is, the larger this value is. We therefore conclude that in a high density electron system, the main contribution comes from Feynman diagrams with algebraic divergence in $|\mathbf{q}|$.

Note: Four momentum in condensed matter physics

In condensed matter physics, though we do not have Lorentz invariance, the four-momentum notation is still used to avoiding using too many letters and to connect the frequency and the momentum of a particle.

5 The electric susceptibility and bosonic modes

Now it is time to evaluate the analytic properties of the susceptibility. The dispersion relations are given by $\epsilon = 0$ (not $\epsilon \rightarrow \infty$ - see Section 2.2.1 in [this optics note](#)). Note that the imaginary part of ϵ indicates damping (see Section 8.1.1 in [this optics note](#)), and therefore if $\text{Im } \epsilon \neq 0$, even if on an (ω, \mathbf{k}) point we have $\text{Re } \epsilon = 0$, this is not an eigenmode. However, if $\text{Im } \epsilon$ is not large, we can still say that “there is a mode at (ω, \mathbf{k}) with a finite lifetime”. So what we are going to do is to solve the equation $\text{Re } \epsilon(\omega, \mathbf{k}) = 0$ and check the lifetime of each mode.

Zhengzhong
Li, Sec. 4.6