

Atomic physics: theories of laser beams and atoms

Jinyuan Wu

January 31, 2023

1 Overview of atoms

Atomic physics relies on the existence of atoms. The idea that matter consists of atoms was seen in ancient Greek, probably earlier in ancient India. In the West, people usually attribute the idea to Democritus (450 BC), who claimed that there are only atoms and “void”. The modern idea of atoms arose to explain the behavior of gases: if we assume the basic degrees of freedom of a gas systems are \mathbf{r}_i 's and \mathbf{p}_i 's, then everything works well – except we need to use a constant \hbar to decide the correct entropy of the gas:

$$d\Omega = \frac{d^3\mathbf{r} d^3\mathbf{p}}{h^3}, \quad h = 2\pi\hbar. \quad (1)$$

The origin of \hbar led to the discovery of quantum mechanics.

The quantum nature of atoms is best demonstrated by beam splitters. Beam splitters are widely used in optics: they can be used to make a Mach-Zehnder interferometer, and the observed intensity takes the form of

$$(\cos(\omega t + \phi_1) + \cos(\omega t + \phi_2))^2,$$

and by moving the mirrors, we change $\phi_{1,2}$, and thus peaks and valleys occur in the relation between I and $\phi_{1,2}$.

More generally, in Figure 1, we have

$$\begin{aligned} |\mathbf{E}_3|^2 + |\mathbf{E}_4|^2 &= (t\mathbf{E}_1 + r\mathbf{E}_2)(t\mathbf{E}_1 + r\mathbf{E}_2)^* + (t\mathbf{E}_2 + r\mathbf{E}_1)(t\mathbf{E}_2 + r\mathbf{E}_1)^* \\ &= (|\mathbf{E}_1|^2 + |\mathbf{E}_2|^2)(|t|^2 + |r|^2) + (\mathbf{E}_1^* \cdot \mathbf{E}_2 + \text{h.c.})(t^*r + r^*t). \end{aligned} \quad (2)$$

To ensure energy conservation, we get

$$|t|^2 + |r|^2 = 1, \quad (3)$$

which is expected, and

$$t^*r + r^*t = 0. \quad (4)$$

The last equation means there has to be a phase shift caused by the beam splitter. In the 50 %-50 % case, we have $r = te^{i\phi}$, and solving the above equation, we find $\phi = 90^\circ$. Thus a 50 %-50 % beam splitter introduces a 90° phase difference between the two output beams if one of the input beam is absent, i.e. the corresponding input port is a dark port.

A comment: when we deal with actual light, not just atom beams, polarization also influences the phase shift.




Figure 1: Beam splitter example

2 Blackbody radiation

Blackbody radiation is where atomic physics started. Both the candle and the incandescent light bulb give light when heated up. What makes it interesting is, if we heat up a black object, radiation also occurs: this can be demonstrated by putting a light mill before a piece of heated black object: the vanes rotate in the same way that they rotate when placed before a light bulb. When the temperature isn't too high, we can put an infrared screener between the light mill and the heated black object and we find the rotation rate of the light mill goes down. This means the relation between radiation and temperature of many objects is at least quantitatively modeled by the ideal black body, and therefore studying its behavior is important.

A blackbody can be modeled as a hole on a closed optical cavity: once something goes inside, it's highly unlikely to be reflected back. The possible optical modes between two walls are like

$$\sin\left(\frac{n\pi x}{L}\right) \quad \text{or} \quad \sin\left(\frac{n\pi y}{L}\right),$$

and $n = 0, 1, 2, 3, \dots$. In a cubic box, we have n_1, n_2, n_3 to label all the modes, and TODO

We will skip the tedious historical attempts to derive the correct blackbody radiation formula, and just jump to the final answer: the energy of each light mode has to be quantized. The first try is to assume

$$E_{\text{mode } \omega} = \hbar\omega n. \quad (5)$$

Thus, for each light mode, we have

$$Z = \sum_{n=0}^{\infty} e^{-n\hbar\omega/k_B T} = \frac{1}{1 - e^{-\hbar\omega/k_B T}}, \quad (6)$$

and therefore

$$\langle E \rangle = \frac{1}{Z} \sum_{n=0}^{\infty} n\hbar\omega e^{-n\hbar\omega/k_B T} = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}. \quad (7)$$

The high temperature limit would be

$$\langle E \rangle = \frac{\hbar\omega}{\frac{\hbar\omega}{k_B T} + \frac{1}{2} \left(\frac{\hbar\omega}{k_B T} \right)^2 + \dots} \rightarrow k_B T \left(1 - \frac{1}{2} \frac{\hbar\omega}{k_B T} \right) = k_B T - \frac{1}{2} \hbar\omega.$$

Thus, in order to go back to the classical prediction when $T \rightarrow \infty$, the energy of a single mode has to be

$$E = \hbar\omega \left(n + \frac{1}{2} \right), \quad (8)$$

and the correct expected energy is

$$\langle E \rangle = \frac{\hbar\omega}{\frac{\hbar\omega}{k_B T} + \frac{1}{2} \left(\frac{\hbar\omega}{k_B T} \right)^2 + \dots} \rightarrow k_B T \left(1 - \frac{1}{2} \frac{\hbar\omega}{k_B T} \right) = k_B T - \frac{1}{2} \hbar\omega + \frac{1}{2} \hbar\omega. \quad (9)$$

Now the total energy of the cavity can be found: it's

$$U = \int_0^{\infty} \langle E_{\omega} \rangle n(\omega) d\omega = L^3 \quad (10)$$

The energy jet can then be found:

$$J = \frac{1}{4} c U = \underbrace{\frac{\pi^2 k_B^4}{60 \hbar^3 c^3}}_{\sigma} T^4. \quad (11)$$

TODO: whether we need to multiply a 2 factor to J ; the direction of radiation; the vibrational 1/2 factor; the

Blackbody radiation is related to the temperature of the Earth, actually. The radiation power of the sun is

$$P_s = 4\pi r_s^2 \cdot \sigma T^4 \cdot \epsilon_s, \quad (12)$$




Figure 2: The energy level diagram of a two-level system

where ϵ_s is a (rather coarse) estimation of how much the Sun deviates from an ideal blackbody. The energy flow at the position of the Earth is

$$I_e = \frac{P_s}{4\pi R_e^2}, \quad (13)$$

and the power received by the Earth is

$$P_e = \pi r_e^2 I_e \epsilon_{e1}. \quad (14)$$

Here note that we are using the cross section of the Earth as the “surface”, because actually what we have is

$$P = \int dS \cdot \mathbf{I}.$$

The balance of energy, therefore, means

$$P_e = \pi r_e^2 I_e \epsilon_{e1} = 4\pi r_e^2 \sigma T_e^4 \epsilon_{e2}. \quad (15)$$

Albedo analysis means the absorption efficiency of the Earth is $\epsilon_{e1} = 0.65$, and if $\epsilon_{e2} \approx 1$, we find $T_e = 251$ K. This of course goes against the observed result 300 K. The point here is the Earth is not really a blackbody ideal enough when it comes to radiation, if we view the atmosphere as a part of the Earth: heat radiation is reflected back, and the amount of energy flowing into the space is reduced. This means the temperature of the Earth has to be higher than what is estimated using (15).

3 Spontaneous radiation

Einstein noticed one deficiency of the traditional picture of atomic radiation: because of the spatial rotational symmetry, what we get is a spherical wave, going out infinitely. Now this means an atom undergoes random walking because of this radiation.

Suppose n_2 is the occupation of the excited state, and n_1 is the occupation of the ground state. Then we have TODO: see my old notes

The time evolution – master equation indeed – of the two-level system is

$$\dot{n}_2 = -n_2 A_{21} - n_2 B_{21} \rho(\omega) d\omega + n_1 B_{12} \rho(\omega) d\omega, \quad (16)$$

and from the fact that $\dot{n}_2 = \dot{n}_1 = 0$ can be reached when we arrive at thermal equilibrium, i.e. when

$$\frac{n_2}{n_1} = e^{-\hbar\omega/k_B}, \quad (17)$$

we get

$$A = \rho(\text{zero point energy}) d\omega B = \frac{1}{2} \frac{\hbar\omega^3}{\pi^2 c^2}. \quad (18)$$

This means spontaneous radiation is necessary, or otherwise our model of the two-level system is unable to stay at the correct equilibrium state.