

8: Lab - Time Series

Environmental Data Analytics | John Fay and Luana Lima

Fall 2024

Objectives

1. Answer questions on M7
2. Explore the components of times series: trend, seasonal, random
3. Perform trend analysis on another dataset

Set up

```
library(tidyverse)
library(lubridate)
#install.packages("trend")
library(trend)
#install.packages("zoo")
library(zoo)
#install.packages("Kendall")
library(Kendall)
#install.packages("tseries")
library(tseries)
library(here)
here

## function (...)
## {
##   .root_env$root$f(...)
## }
## <bytecode: 0x12d530498>
## <environment: namespace:here>

# Set theme
mytheme <- theme_classic(base_size = 14) +
  theme(axis.text = element_text(color = "black"),
        legend.position = "top")
theme_set(mytheme)
```

Import Datasets

Today we will work with data from the Climate Change Knowledge Portal from the World Bank Group. More specifically historical rainfall and temperature averages for Brazil. You will find two new data files on

folder “/Data/Raw/”. One with rainfall named “pr_1901_2016_BRA.csv” and another with temperature named “tas_1901_2016_BRA.csv”. The data span the period from 1901 to 2016 in monthly steps. You can download the data [here][<https://climateknowledgeportal.worldbank.org/download-data>]

Research question: Can you see any changes on the rainfall regime or average temperature over time?

#Exercise 1: Import both datasets using the read.csv function.

```
Rainfall_BR <- read.csv(here("Data/Raw/pr_1901_2016_BRA.csv"), stringsAsFactors = TRUE)
```

```
Temp_BR <- read.csv(here("Data/Raw/tas_1901_2016_BRA.csv"), stringsAsFactors = TRUE)
```

#Exercise 2: Tidy the rainfall data sets.

#a Rename the column with Rainfall to get rid of the dots.

#b Note that on both data sets that is a column with the month name and average. Convert it to a Month

#c Now you should have a column with Month. Use the paste0() function to paste month and year together

#d Select only the columns of interest: Date and rainfall

```
Rainfall_BR_processed <-
```

```
  Rainfall_BR %>%
```

```
  rename( Rainfall = Rainfall...MM.) %>%
```

```
  separate(Statistics,c("Null","Month","Null2")," ") %>%
```

```
  mutate( Date = my(paste0(Month,"-",Year))) %>%
```

```
  select(Date,Rainfall)
```

#Exercise 3: Repeat exercise 2 to the temperature dataset.

```
Temp_BR_processed <-
```

```
  Temp_BR %>%
```

```
  rename( Temperature_C = Temperature...Celsius.) %>%
```

```
  separate(Statistics,c("Null","Month","Null2")," ") %>%
```

```
  mutate( Date = my(paste0(Month,"-",Year))) %>%
```

```
  select(Date,Temperature_C)
```

#Exercise 4: Join the temperature and rainfall into one tidy data frame with 3 columns: Date, Rainfall

```
BR_complete <- inner_join(Rainfall_BR_processed,Temp_BR_processed)
```

```
## Joining with 'by = join_by(Date)'
```

Initial plots

#Exercise 5: Check if there is any missing data for both temperature and rainfall series.

```
summary(BR_complete$Rainfall)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##  33.86   77.80  143.60  145.15  205.44  320.65
```

#no NAs

```
summary(BR_complete$Temperature_C)
```


```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##  21.73   24.27   25.20   24.96   25.67   27.35
```

```
#no NAs
```

```
#Exercise 6: Plot both series over time on separate plots. Add a trend line using geom_smooth(). Can you
```


```
ggplot(BR_complete, aes( x = Date, y = Rainfall)) +  
  geom_line() +  
  geom_smooth()
```

```
## 'geom_smooth()' using method = 'gam' and formula = 'y ~ s(x, bs = "cs")'
```



```
ggplot(BR_complete, aes( x = Date, y = Temperature_C)) +  
  geom_line() +  
  geom_smooth(method="lm")
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```



Creating time series objects

#Exercise 7: Create a time series objects for each series using ts(). Make sure you specify the starting

```
f_month <- month(first(BR_complete$Date))
f_year <- year(first(BR_complete$Date))

BR_Rain_ts <- ts(BR_complete$Rainfall,frequency = 12,start=c(1901,1))
BR_Temp_ts <- ts(BR_complete$Temperature_C,frequency = 12,start=c(f_year,f_month))
```

Decomposing a time series dataset


A given time series can be made up of several component series:

1. A **seasonal** component, which repeats over a fixed known period (e.g., seasons of the year, months, days of the week, hour of the day)
2. A **trend** component, which quantifies the upward or downward progression over time. The trend component of a time series does not have to be monotonic.
3. An **error** or **random** component, which makes up the remainder of the time series after other components have been accounted for. This component reflects the noise in the dataset.
4. (optional) A **cyclical** component, which repeats over periods greater than the seasonal component. A good example of this is El Niño Southern Oscillation (ENSO) cycles, which occur over a period of 2-8 years.


The `stl` function decomposes the time series object into its component parts. We must specify that the window for seasonal extraction is either “periodic” or a specific number of at least 7. The decomposition proceeds through a loess (locally estimated scatterplot smoothing) function.

*# Exercise 8: Use the `stl()` function to decompose both series.
Visualize the decomposed series using `plot()`. What components can you identify?*

```
BR_Rain_Decomposed <- stl(BR_Rain_ts, s.window = "periodic")  
plot(BR_Rain_Decomposed)
```



```
BR_Temp_Decomposed <- stl(BR_Temp_ts, s.window = "periodic")  
plot(BR_Temp_Decomposed)
```



Exercise 9: Use the times series object generated with the stl() to extract the components of each series and turn them into data frames.

```
BR_Rain_Components <- as.data.frame(BR_Rain_Decomposed$time.series[,1:3])
BR_Temp_Components <- as.data.frame(BR_Temp_Decomposed$time.series[,1:3])
```


```
BR_Rain_Components <- mutate(BR_Rain_Components,
  Observed = BR_complete$Rainfall,
  Date = BR_complete$Date)
```

```
BR_Temp_Components <- mutate(BR_Temp_Components,
  Observed = BR_complete$Temperature_C,
  Date = BR_complete$Date)
```


Exercise 10: Visualize how the trend maps onto the data for both series

```
ggplot(BR_Rain_Components) +
  geom_line(aes(y = Observed, x = Date), size = 0.25) +
  geom_line(aes(y = trend, x = Date), color = "#c13d75ff") +
  #geom_line(aes(y = seasonal, x = Date), color = "blue") +
  #geom_hline(yintercept = 0, lty = 2) +
  ylab("Rainfall")
```

```
## Warning: Using 'size' aesthetic for lines was deprecated in ggplot2 3.4.0.
## i Please use 'linewidth' instead.
## This warning is displayed once every 8 hours.
## Call 'lifecycle::last_lifecycle_warnings()' to see where this warning was
## generated.
```



```
ggplot(BR_Temp_Components) +  
  geom_line(aes(y = Observed, x = Date), size = 0.25) +  
  geom_line(aes(y = trend, x = Date), color = "#c13d75ff") +  
  #geom_line(aes(y = seasonal, x = Date), color = "blue") +  
  #geom_hline(yintercept = 0, lty = 2) +  
  ylab("Temperature")
```



Trend analysis

Monotonic trends are a gradual shift over time that is consistent in direction. Specific tests for monotonic trend analysis are listed below, with assumptions and tips:

- **linear regression:** no seasonality, fits the assumptions of a parametric test. Function: `lm` Regress time series over time. If significant relationship, then there is a trend.
- **Mann-Kendall:** no seasonality, non-parametric, missing data allowed. Function: `Kendall::MannKendall()` or `trend::mk.test()`

$H_0: S = 0$, i.e., no trend $H_1: S \neq 0$, i.e., follow a trend

- **Seasonal Mann-Kendall:** seasonality, non-parametric `Kendall::SeasonalMannKendall` or `trend::smk.test()`
- **Spearman Rho:** no seasonality, non-parametric, missing data allowed. Function: `stats::cor.test(method="spearmanr")`. Similar to `lm` but allows for non-linear trend.

Exercise 9: Apply one of the trend detection test to the original data set. Remember that the data has

```
Rain_trend <- trend::smk.test(BR_Rain_ts)
# Inspect results
Rain_trend
```



```
##
## Seasonal Mann-Kendall trend test (Hirsch-Slack test)
##
## data: BR_Rain_ts
## z = 4.8698, p-value = 1.117e-06
## alternative hypothesis: true S is not equal to 0
## sample estimates:
##      S      varS
## 7071 2107719
```

```
summary(Rain_trend)
```

```
##
## Seasonal Mann-Kendall trend test (Hirsch-Slack test)
##
## data: BR_Rain_ts
## alternative hypothesis: two.sided
##
## Statistics for individual seasons
##
## H0
##
##      S      varS      tau      z Pr(>|z|)
## Season 1: S = 0 662 175643.3 0.099 1.577 0.1147505
## Season 2: S = 0 828 175643.3 0.124 1.973 0.0484632 *
## Season 3: S = 0 486 175643.3 0.073 1.157 0.2471716
## Season 4: S = 0 1230 175643.3 0.184 2.932 0.0033626 **
## Season 5: S = 0 1148 175643.3 0.172 2.737 0.0062035 **
## Season 6: S = 0 756 175643.3 0.113 1.801 0.0716262 .
## Season 7: S = 0 1290 175643.3 0.193 3.076 0.0021004 **
## Season 8: S = 0 28 175643.3 0.004 0.064 0.9486326
## Season 9: S = 0 -110 175643.3 -0.016 -0.260 0.7948004
## Season 10: S = 0 -594 175643.3 -0.089 -1.415 0.1570853
## Season 11: S = 0 988 175643.3 0.148 2.355 0.0185199 *
## Season 12: S = 0 359 175642.3 0.054 0.854 0.3929845
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Temp_trend <- trend::smk.test(BR_Temp_ts)
# Inspect results
Temp_trend
```

```
##
## Seasonal Mann-Kendall trend test (Hirsch-Slack test)
##
## data: BR_Temp_ts
## z = 20.178, p-value < 2.2e-16
## alternative hypothesis: true S is not equal to 0
## sample estimates:
##      S      varS
## 29295 2107713
```

```
summary(Temp_trend)
```

```
##
## Seasonal Mann-Kendall trend test (Hirsch-Slack test)
##
## data: BR_Temp_ts
## alternative hypothesis: two.sided
##
## Statistics for individual seasons
##
## H0
##
```

		S	varS	tau	z	Pr(> z)	
## Season 1:	S = 0	2740	175641.3	0.411	6.535	6.3398e-11	***
## Season 2:	S = 0	2251	175642.3	0.338	5.369	7.9313e-08	***
## Season 3:	S = 0	2581	175642.3	0.387	6.156	7.4562e-10	***
## Season 4:	S = 0	2375	175642.3	0.356	5.665	1.4740e-08	***
## Season 5:	S = 0	2093	175642.3	0.314	4.992	5.9855e-07	***
## Season 6:	S = 0	1692	175643.3	0.254	4.035	5.4636e-05	***
## Season 7:	S = 0	1706	175643.3	0.256	4.068	4.7366e-05	***
## Season 8:	S = 0	2606	175643.3	0.391	6.216	5.1088e-10	***
## Season 9:	S = 0	2053	175642.3	0.308	4.896	9.7687e-07	***
## Season 10:	S = 0	3286	175643.3	0.493	7.838	4.5684e-15	***
## Season 11:	S = 0	2822	175643.3	0.423	6.731	1.6836e-11	***
## Season 12:	S = 0	3090	175643.3	0.463	7.371	1.6988e-13	***


```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Exercise 10: Now remove the seasonality and apply another test. Compare the results with what you obtained.

```
#removing seasonality for temperature
BR_Temp_Nonseas <- BR_Temp_ts - BR_Temp_Decomposed$time.series[,1]

BR_Temp_Components <- mutate(BR_Temp_Components,
                             Nonseasonal = BR_Temp_Nonseas)

#plotting resulting series
ggplot(BR_Temp_Components) +
  geom_line(aes(y = Observed, x = Date), size = 0.25) +
  geom_line(aes(y = Nonseasonal, x = Date), color = "#c13d75ff") +
  ylab("Temperature")
```



```
Temp_Nonseas_trend <- trend::mk.test(BR_Temp_Nonseas)
# Inspect results
Temp_Nonseas_trend
```


```
##
## Mann-Kendall trend test
##
## data: BR_Temp_Nonseas
## z = 19.941, n = 1392, p-value < 2.2e-16
## alternative hypothesis: true S is not equal to 0
## sample estimates:
##          S          varS          tau
## 3.453990e+05 3.000146e+08 3.567683e-01
```

```
#removing seasonality for rainfall
BR_Rain_Nonseas <- BR_Rain_ts - BR_Rain_Decomposed$time.series[,1]

BR_Rain_Components <- mutate(BR_Rain_Components,
                             Nonseasonal = BR_Rain_Nonseas)

#plotting resulting series
ggplot(BR_Rain_Components) +
  geom_line(aes(y = Observed, x = Date), size = 0.25) +
  geom_line(aes(y = Nonseasonal, x = Date), color = "#c13d75ff") +
  geom_smooth(aes(y = Nonseasonal, x = Date)) +
  ylab("Rainfall")
```

```
## 'geom_smooth()' using method = 'gam' and formula = 'y ~ s(x, bs = "cs")'
```



```
Rain_Nonseas_trend <- trend::mk.test(BR_Rain_Nonseas)
# Inspect results
Rain_Nonseas_trend
```

```
##
## Mann-Kendall trend test
##
## data: BR_Rain_Nonseas
## z = 4.9012, n = 1392, p-value = 9.523e-07
## alternative hypothesis: true S is not equal to 0
## sample estimates:
##      S      varS      tau
## 8.489500e+04 3.000146e+08 8.768917e-02
```

What would we conclude based on these findings?

Answer: